
Faculty of Science

Faculty Publications

Measurement of double-differential muon neutrino charged-current interactions on C_8H_8 without pions in the final state using the T2K off-axis beam

K. Abe, C. Andreopoulos, M. Antonova, S. Aoki, A. Ariga, D. Karlen, ... & J. Žmuda

June 2016

© 2016 K. Abe et al. This is an open access article distributed under the terms of the Creative Commons Attribution License. <https://creativecommons.org/licenses/by/3.0/>

This article was originally published at:

<https://doi.org/10.1103/PhysRevD.93.112012>

Citation for this paper:

Abe, K., Andreopoulos, C., Antonova, M., Aoki, S., Ariga, A., Karlen, D., ... Žmuda, A. (2016). Measurement of double-differential muon neutrino charged-current interactions on C_8H_8 without pions in the final state using the T2K off-axis beam. *Physical Review D*, 93(11), 1-25. <https://doi.org/10.1103/PhysRevD.93.112012>.

Measurement of double-differential muon neutrino charged-current interactions on C_8H_8 without pions in the final state using the T2K off-axis beam

K. Abe,⁴⁷ C. Andreopoulos,^{45,26} M. Antonova,²¹ S. Aoki,²³ A. Ariga,¹ S. Assylbekov,⁷ D. Autiero,²⁸ M. Barbi,³⁹ G. J. Barker,⁵⁵ G. Barr,³⁵ P. Bartet-Friburg,³⁶ M. Batkiewicz,¹² V. Berardi,¹⁷ S. Berkman,³ S. Bhadra,⁵⁹ A. Blondel,¹¹ S. Bolognesi,⁵ S. Bordini,¹⁴ S. B. Boyd,⁵⁵ D. Brailsford,^{25,16} A. Bravar,¹¹ C. Bronner,²² M. Buizza Avanzini,⁹ R. G. Calland,²² S. Cao,²⁴ J. Caravaca Rodríguez,¹⁴ S. L. Cartwright,⁴³ R. Castillo,¹⁴ M. G. Catanesi,¹⁷ A. Cervera,¹⁵ D. Cherdack,⁷ N. Chikuma,⁴⁶ G. Christodoulou,²⁶ A. Clifton,⁷ J. Coleman,²⁶ G. Collazuol,¹⁹ L. Cremonesi,³⁸ A. Dabrowska,¹² G. De Rosa,¹⁸ T. Dealtry,²⁵ P. F. Denner,⁵⁵ S. R. Dennis,^{55,45} C. Densham,⁴⁵ D. Dewhurst,³⁵ F. Di Lodovico,³⁸ S. Di Luise,¹⁰ S. Dolan,³⁵ O. Drapier,⁹ K. E. Duffy,³⁵ J. Dumarchez,³⁶ S. Dytman,³⁷ M. Dziewiecki,⁵⁴ S. Emery-Schrenk,⁵ A. Ereditato,¹ T. Feusels,³ A. J. Finch,²⁵ G. A. Fiorentini,⁵⁹ M. Friend,^{13,*} Y. Fujii,^{13,*} D. Fukuda,³³ Y. Fukuda,³⁰ A. P. Furmanski,⁵⁵ V. Galymov,²⁸ A. Garcia,¹⁴ S. G. Giffin,³⁹ C. Giganti,³⁶ F. Gizzarelli,⁵ M. Gonin,⁹ N. Grant,²⁵ D. R. Hadley,⁵⁵ L. Haegel,¹¹ M. D. Haigh,⁵⁵ P. Hamilton,¹⁶ D. Hansen,³⁷ T. Hara,²³ M. Hartz,^{22,51} T. Hasegawa,^{13,*} N. C. Hastings,³⁹ T. Hayashino,²⁴ Y. Hayato,^{47,22} R. L. Helmer,⁵¹ M. Hierholzer,¹ A. Hillairet,⁵² A. Himmel,⁸ T. Hiraki,²⁴ S. Hirota,²⁴ M. Hogan,⁷ J. Holeczek,⁴⁴ S. Horikawa,¹⁰ F. Hosomi,⁴⁶ K. Huang,²⁴ A. K. Ichikawa,²⁴ K. Ieki,²⁴ M. Ikeda,⁴⁷ J. Imber,⁹ J. Insler,²⁷ R. A. Intonti,¹⁷ T. J. Irvine,⁴⁸ T. Ishida,^{13,*} T. Ishii,^{13,*} E. Iwai,¹³ K. Iwamoto,⁴⁰ A. Izmaylov,^{15,21} A. Jacob,³⁵ B. Jamieson,⁵⁷ M. Jiang,²⁴ S. Johnson,⁶ J. H. Jo,³² P. Jonsson,¹⁶ C. K. Jung,^{32,†} M. Kabirnezhad,³¹ A. C. Kaboth,^{41,45} T. Kajita,^{48,†} H. Kakuno,⁴⁹ J. Kameda,⁴⁷ D. Karlen,^{52,51} I. Karpikov,²¹ T. Katori,³⁸ E. Kearns,^{2,22,†} M. Khabibullin,²¹ A. Khotjantsev,²¹ D. Kielczewska,⁵³ T. Kikawa,²⁴ H. Kim,³⁴ J. Kim,³ S. King,³⁸ J. Kisiel,⁴⁴ A. Knight,⁵⁵ A. Knox,²⁵ T. Kobayashi,^{13,*} L. Koch,⁴² T. Koga,⁴⁶ A. Konaka,²⁴ K. Kondo,²⁴ A. Kopylov,²¹ L. L. Kormos,²⁵ A. Korzenev,¹¹ Y. Koshio,^{33,†} W. Kropp,⁴ Y. Kudenko,^{21,‡} R. Kurjata,⁵⁴ T. Kutter,²⁷ J. Lagoda,³¹ I. Lamont,²⁵ E. Larkin,⁵⁵ P. Lasorak,^{38,38} M. Laveder,¹⁹ M. Lawe,²⁵ M. Lazos,²⁶ T. Lindner,⁵¹ Z. J. Liptak,⁶ R. P. Litchfield,⁵⁵ X. Li,³² A. Longhin,¹⁹ J. P. Lopez,⁶ L. Ludovici,²⁰ X. Lu,³⁵ L. Magaletti,¹⁷ K. Mahn,²⁹ M. Malek,⁴³ S. Manly,⁴⁰ A. D. Marino,⁶ J. Marteau,²⁸ J. F. Martin,⁵⁰ P. Martins,³⁸ S. Martynenko,³² T. Maruyama,^{13,*} V. Matveev,²¹ K. Mavrokoridis,²⁶ W. Y. Ma,¹⁶ E. Mazzucato,⁵ M. McCarthy,⁵⁹ N. McCauley,²⁶ K. S. McFarland,⁴⁰ C. McGrew,³² A. Mefodiev,²¹ M. Mezzetto,¹⁹ P. Mijakowski,³¹ A. Minamino,²⁴ O. Mineev,²¹ S. Mine,⁴ A. Missert,⁶ M. Miura,^{47,†} S. Moriyama,^{47,†} Th. A. Mueller,⁹ S. Murphy,¹⁰ J. Myslik,⁵² T. Nakadaira,^{13,*} M. Nakahata,^{47,22} K. G. Nakamura,²⁴ K. Nakamura,^{22,13,*} K. D. Nakamura,²⁴ S. Nakayama,^{47,†} T. Nakaya,^{24,22} K. Nakayoshi,^{13,*} C. Nantais,³ C. Nielsen,³⁴ M. Nirkko,¹ K. Nishikawa,^{13,*} Y. Nishimura,⁴⁸ J. Nowak,²⁵ H. M. O'Keeffe,²⁵ R. Ohta,^{13,*} K. Okumura,^{48,22} T. Okusawa,³² W. Oryszczak,⁵³ S. M. Oser,³ T. Ovsyannikova,²¹ R. A. Owen,³⁸ Y. Oyama,^{13,*} V. Palladino,¹⁸ J. L. Palomino,³² V. Paolone,³⁷ N. D. Patel,²⁴ M. Pavin,³⁶ D. Payne,²⁶ J. D. Perkin,⁴³ Y. Petrov,³ L. Pickard,⁴³ L. Pickering,¹⁶ E. S. Pinzon Guerra,⁵⁹ C. Pistillo,¹ B. Popov,^{36,8} M. Posiadala-Zezula,⁵³ J.-M. Poutissou,⁵¹ R. Poutissou,⁵¹ P. Przewlocki,³¹ B. Quilain,²⁴ E. Radicioni,¹⁷ P. N. Ratoff,²⁵ M. Ravonel,¹¹ M. A. M. Rayner,¹¹ A. Redij,¹ E. Reinherz-Aronis,⁷ C. Riccio,¹⁸ P. Rojas,⁷ E. Rondio,³¹ S. Roth,⁴² A. Rubbia,¹⁰ A. Rychter,⁵⁴ R. Sacco,³⁸ K. Sakashita,^{13,*} F. Sánchez,¹⁴ F. Sato,¹³ E. Scantamburlo,¹¹ K. Scholberg,^{8,†} S. Schoppmann,⁴² J. Schwehr,⁷ M. Scott,⁵¹ Y. Seiya,³⁴ T. Sekiguchi,^{13,*} H. Sekiya,^{47,22,†} D. Sgalaberna,¹⁰ R. Shah,^{45,35} A. Shaikhiev,²¹ F. Shaker,⁵⁷ D. Shaw,²⁵ M. Shiozawa,^{47,22} T. Shirahige,³³ S. Short,³⁸ M. Smy,⁴ J. T. Sobczyk,⁵⁸ M. Sorel,¹⁵ L. Southwell,²⁵ P. Stamoulis,¹⁵ J. Steinmann,⁴² T. Stewart,⁴⁵ Y. Suda,⁴⁶ S. Suvorov,²¹ A. Suzuki,^{23,*} K. Suzuki,²⁴ S. Y. Suzuki,¹³ Y. Suzuki,²² R. Tacik,^{39,51} M. Tada,^{13,*} S. Takahashi,²⁴ A. Takeda,⁴⁷ Y. Takeuchi,^{23,22} H. K. Tanaka,^{47,†} H. A. Tanaka,^{50,51,||} D. Terhorst,⁴² R. Terri,³⁸ T. Thakore,²⁷ L. F. Thompson,⁴³ S. Tobayama,³ W. Toki,⁷ T. Tomura,⁴⁷ C. Touramanis,²⁶ T. Tsukamoto,^{13,*} M. Tzanov,²⁷ Y. Uchida,¹⁶ A. Vacheret,³⁵ M. Vagins,^{22,4} Z. Vallari,³² G. Vasseur,⁵ T. Wachala,¹² K. Wakamatsu,³⁴ C. W. Walter,^{8,†} D. Wark,^{45,35} W. Warzocha,⁵³ M. O. Wascko,¹⁶ A. Weber,^{45,35} R. Wendell,^{24,†} R. J. Wilkes,⁵⁶ M. J. Wilking,³² C. Wilkinson,¹ J. R. Wilson,³⁸ R. J. Wilson,⁷ Y. Yamada,^{13,*} K. Yamamoto,³⁴ M. Yamamoto,²⁴ C. Yanagisawa,^{32,||} T. Yano,²³ S. Yen,⁵¹ N. Yershov,²¹ M. Yokoyama,^{46,†} K. Yoshida,²⁴ T. Yuan,⁶ M. Yu,⁵⁹ A. Zalewska,¹² J. Zalipska,³¹ L. Zambelli,^{13,*} K. Zaremba,⁵⁴ M. Ziembicki,⁵⁴ E. D. Zimmerman,⁶ M. Zito,⁵ and J. Żmuda⁵⁸

(T2K Collaboration)

¹Albert Einstein Center for Fundamental Physics, Laboratory for High Energy Physics (LHEP), University of Bern, Bern, Switzerland

²Department of Physics, Boston University, Boston, Massachusetts, USA

³Department of Physics and Astronomy, University of British Columbia, Vancouver, British Columbia, Canada

⁴Department of Physics and Astronomy, University of California, Irvine, Irvine, California, USA

⁵IRFU, CEA Saclay, Gif-sur-Yvette, France

- ⁶*Department of Physics, University of Colorado at Boulder, Boulder, Colorado, USA*
- ⁷*Department of Physics, Colorado State University, Fort Collins, Colorado, USA*
- ⁸*Department of Physics, Duke University, Durham, North Carolina, USA*
- ⁹*Ecole Polytechnique, IN2P3-CNRS, Laboratoire Leprince-Ringuet, Palaiseau, France*
- ¹⁰*ETH Zurich, Institute for Particle Physics, Zurich, Switzerland*
- ¹¹*University of Geneva, Section de Physique, DPNC, Geneva, Switzerland*
- ¹²*H. Niewodniczanski Institute of Nuclear Physics PAN, Cracow, Poland*
- ¹³*High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki, Japan*
- ¹⁴*Institut de Fisica d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Campus UAB, Bellaterra (Barcelona) Spain*
- ¹⁵*IFIC (CSIC & University of Valencia), Valencia, Spain*
- ¹⁶*Department of Physics, Imperial College London, London, United Kingdom*
- ¹⁷*INFN Sezione di Bari and Università e Politecnico di Bari, Dipartimento Interuniversitario di Fisica, Bari, Italy*
- ¹⁸*Dipartimento di Fisica, INFN Sezione di Napoli and Università di Napoli, Napoli, Italy*
- ¹⁹*Dipartimento di Fisica, INFN Sezione di Padova and Università di Padova, Padova, Italy*
- ²⁰*INFN Sezione di Roma and Università di Roma "La Sapienza", Roma, Italy*
- ²¹*Institute for Nuclear Research of the Russian Academy of Sciences, Moscow, Russia*
- ²²*Kavli Institute for the Physics and Mathematics of the Universe (WPI), The University of Tokyo Institutes for Advanced Study, University of Tokyo, Kashiwa, Chiba, Japan*
- ²³*Kobe University, Kobe, Japan*
- ²⁴*Department of Physics, Kyoto University, Kyoto, Japan*
- ²⁵*Physics Department, Lancaster University, Lancaster, United Kingdom*
- ²⁶*Department of Physics, University of Liverpool, Liverpool, United Kingdom*
- ²⁷*Department of Physics and Astronomy, Louisiana State University, Baton Rouge, Louisiana, USA*
- ²⁸*Université de Lyon, Université Claude Bernard Lyon 1, IPN Lyon (IN2P3), Villeurbanne, France*
- ²⁹*Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan, USA*
- ³⁰*Department of Physics, Miyagi University of Education, Sendai, Japan*
- ³¹*National Centre for Nuclear Research, Warsaw, Poland*
- ³²*Department of Physics and Astronomy, State University of New York at Stony Brook, Stony Brook, New York, USA*
- ³³*Department of Physics, Okayama University, Okayama, Japan*
- ³⁴*Department of Physics, Osaka City University, Osaka, Japan*
- ³⁵*Department of Physics, Oxford University, Oxford, United Kingdom*
- ³⁶*UPMC, Université Paris Diderot, CNRS/IN2P3, Laboratoire de Physique Nucléaire et de Hautes Energies (LPNHE), Paris, France*
- ³⁷*Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, Pennsylvania, USA*
- ³⁸*School of Physics and Astronomy, Queen Mary University of London, London, United Kingdom*
- ³⁹*Department of Physics, University of Regina, Regina, Saskatchewan, Canada*
- ⁴⁰*Department of Physics and Astronomy, University of Rochester, Rochester, New York, USA*
- ⁴¹*Department of Physics, Royal Holloway University of London, Egham, Surrey, United Kingdom*
- ⁴²*RWTH Aachen University, III. Physikalisches Institut, Aachen, Germany*
- ⁴³*Department of Physics and Astronomy, University of Sheffield, Sheffield, United Kingdom*
- ⁴⁴*Institute of Physics, University of Silesia, Katowice, Poland*
- ⁴⁵*STFC, Rutherford Appleton Laboratory, Harwell Oxford, and Daresbury Laboratory, Warrington, United Kingdom*
- ⁴⁶*Department of Physics, University of Tokyo, Tokyo, Japan*
- ⁴⁷*Institute for Cosmic Ray Research, Kamioka Observatory, University of Tokyo, Kamioka, Japan*
- ⁴⁸*Institute for Cosmic Ray Research, Research Center for Cosmic Neutrinos, University of Tokyo, Kashiwa, Japan*
- ⁴⁹*Department of Physics, Tokyo Metropolitan University, Tokyo, Japan*
- ⁵⁰*Department of Physics, University of Toronto, Toronto, Ontario, Canada*
- ⁵¹*TRIUMF, Vancouver, British Columbia, Canada*
- ⁵²*Department of Physics and Astronomy, University of Victoria, Victoria, British Columbia, Canada*
- ⁵³*Faculty of Physics, University of Warsaw, Warsaw, Poland*
- ⁵⁴*Warsaw University of Technology, Institute of Radioelectronics, Warsaw, Poland*
- ⁵⁵*Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵⁶*Department of Physics, University of Washington, Seattle, Washington, USA*
- ⁵⁷*Department of Physics, University of Winnipeg, Winnipeg, Manitoba, Canada*

⁵⁸*Faculty of Physics and Astronomy, Wrocław University, Wrocław, Poland*⁵⁹*Department of Physics and Astronomy, York University, Toronto, Ontario, Canada*

(Received 18 February 2016; published 21 June 2016)

We report the measurement of muon neutrino charged-current interactions on carbon without pions in the final state at the T2K beam energy using 5.734×10^{20} protons on target. For the first time the measurement is reported as a flux-integrated, double-differential cross section in muon kinematic variables $(\cos \theta_\mu, p_\mu)$, without correcting for events where a pion is produced and then absorbed by final state interactions. Two analyses are performed with different selections, background evaluations and cross-section extraction methods to demonstrate the robustness of the results against biases due to model-dependent assumptions. The measurements compare favorably with recent models which include nucleon-nucleon correlations but, given the present precision, the measurement does not distinguish among the available models. The data also agree with Monte Carlo simulations which use effective parameters that are tuned to external data to describe the nuclear effects. The total cross section in the full phase space is $\sigma = (0.417 \pm 0.047(\text{syst}) \pm 0.005(\text{stat})) \times 10^{-38} \text{ cm}^2 \text{ nucleon}^{-1}$ and the cross section integrated in the region of phase space with largest efficiency and best signal-over-background ratio ($\cos \theta_\mu > 0.6$ and $p_\mu > 200 \text{ MeV}$) is $\sigma = (0.202 \pm 0.036(\text{syst}) \pm 0.003(\text{stat})) \times 10^{-38} \text{ cm}^2 \text{ nucleon}^{-1}$.

DOI: 10.1103/PhysRevD.93.112012

I. INTRODUCTION

Accelerator-driven neutrino oscillation measurements [1–3] make use of neutrino beams with energies of a few GeV or lower, at which one of the main interaction processes of neutrinos with nuclei is the charged current quasielastic scattering (CCQE) process. In this process, a muon neutrino (ν_μ) interacts with a neutron n to produce a muon and a proton p through the exchange of a W boson ($\nu_\mu + n \rightarrow \mu^- + p$). This interaction is exploited in long-baseline neutrino oscillation experiments for the signal events with which to measure the neutrino appearance and disappearance probabilities as a function of neutrino energy. It is therefore of primary importance for the CCQE process to be well modeled.

Over the past ten years, a complicated experimental and theoretical picture has emerged regarding CCQE interactions on nuclear targets. The K2K experiment noted that the kinematic distributions of the outgoing muons were not consistent with the prediction of the relativistic Fermi gas (RFG) nuclear model [4,5]. K2K extracted an effective nucleon axial mass $M_A^{\text{QE}} = 1.20 \pm 0.12 \text{ GeV}$ for interactions on a water target [6], in presence of nuclear effects,

compared to the value $1.014 \pm 0.014 \text{ GeV}$ determined from neutrino scattering on deuterium as well as from pion electroproduction [7,8]. MiniBooNE also reported a similar anomaly on mineral oil (CH_2), with a large data set of neutrinos (effective M_A^{QE} of $1.35 \pm 0.17 \text{ GeV}$ [9]) and of antineutrinos [10], as did MINOS, using iron as a target [11]. Both experiments also noted a discrepancy at the lowest values of momentum transfer squared ($Q^2 < 0.2 \text{ GeV}^2$). SciBooNE has also reported similar results [12]. The previous T2K off-axis CCQE measurement [13] was also consistent with a large effective M_A^{QE} . The T2K on-axis measurement [14] has large systematic uncertainties and is compatible with different values of M_A^{QE} , depending on whether only one muon track is reconstructed, or both the muon and proton tracks are reconstructed. Other data sets on nuclear targets from the NOMAD (carbon target) and MINERνA (hydrocarbon target) experiments are in agreement with an M_A^{QE} of 1 GeV. At beam energies of 3–100 GeV, the NOMAD experiment has reported an M_A^{QE} of $1.05 \pm 0.06 \text{ GeV}$ [15]. The MINERνA ($E_\nu \sim 3.5 \text{ GeV}$) experiment has also measured the CCQE cross section with only a muon and a proton in the final state [16] which is consistent with the RFG model and $M_A^{\text{QE}} \sim 1 \text{ GeV}$. Interestingly, previous MINERνA CCQE measurements which use muon information and the calorimetric recoil energy with neutrinos [17] and antineutrinos [18] prefer a transverse enhancement model, suggesting the presence of meson exchange currents [19]. The measurements of the neutrino interaction rate depend on the neutrino cross section and the flux. MiniBooNE and MINERνA are working to improve the flux modeling and thus apply a more precise flux correction to the previously mentioned cross-section measurements.

* Also at J-PARC, Tokai, Japan.

† Also at Kavli IPMU (WPI), the University of Tokyo, Japan.

‡ Also at National Research Nuclear University “MEPhI” and Moscow Institute of Physics and Technology, Moscow, Russia.

§ Also at JINR, Dubna, Russia.

|| Also at Institute of Particle Physics, Canada.

¶ Also at BMCC/CUNY, Science Department, New York, New York, USA.

The anomalies measured in neutrino interactions by modern experiments, using relatively heavy nuclei as targets, may be explained by the contribution of nuclear effects. These were not needed in the models designed to describe bubble-chamber data on deuterium targets. Various different implementations of such nuclear effects have been proposed [20–30]. In experimental measurements the effects of nucleon and hadron initial and final state interactions with the nucleus cannot typically be disentangled from the fundamental neutrino-interaction cross section. The phenomenological interpretation of modern measurements is therefore complicated. Given the discrepancies between the available predictions or their incompleteness in the description of such nuclear effects, it is important to provide experimental measurements which are, as much as possible, model independent, in order to reduce the systematic uncertainties which arise from model dependencies, and to produce results that are useful for comparison with the present and future models.

The Tokai-to-Kamioka (T2K) experiment has a suite of neutrino detectors placed in a neutrino beam with energy peaked at $E_\nu = 0.6$ GeV. This paper describes the measurement of the CCQE-like neutrino interaction cross section, by selecting events without pions in the final sample, with plastic scintillator (C_8H_8) as the target material, using the ND280 off-axis near detector in the T2K beam. Particular care has been taken to avoid model-dependent corrections to the data. Two analyses, which follow different approaches to measure the double-differential cross section as a function of muon momentum and angle, are presented.

The paper is organized as follows. The experimental apparatus is described in Sec. II. The general strategy for the measurement as well as the data samples and the Monte Carlo simulation used are presented in Sec. III, followed by a presentation of the two analyses and their results. The analyses are compared and the results discussed in Sec. IV. Finally, conclusions are reported in Sec. V.

II. EXPERIMENTAL APPARATUS

The T2K long-baseline neutrino oscillation experiment [31] uses a beam of muon neutrinos to study the appearance of electron neutrinos ($\nu_\mu \rightarrow \nu_e$) and to measure or constrain the Pontecorvo-Maki-Nakagawa-Sakata mixing angles θ_{13} and θ_{23} , the mass splitting $|\Delta m_{32}^2|$ and the CP -violating phase δ_{CP} . The neutrinos are produced at the Japan Proton Accelerator Research Complex (J-PARC) in Ibaraki prefecture on the eastern coast of Japan, and travel 295 km through the Earth before reaching the far detector, Super-Kamiokande [32], in Gifu prefecture. A complex of near detectors located 280 meters from the proton beam target is used to characterize the neutrino beam before oscillation, reducing the systematic uncertainties, and to study neutrino interactions as in the measurement reported here.

A. Neutrino beam

T2K uses a conventional neutrino beam, in which the muon neutrinos are produced by the decay of charged pions and kaons. Protons are first accelerated to 30 GeV by a sequence of three accelerators in J-PARC, then extracted to the neutrino beam line where they are directed onto a graphite target. The resulting collisions produce hadrons—predominantly charged pions—which travel inside a 96-m-long decay volume where they decay in flight into muons and muon neutrinos. A set of three magnetic horns is used to focus the positively charged hadrons and defocus the negative particles, enhancing the neutrino component of the beam while reducing its contamination by antineutrinos. At the end of the decay volume, a beam dump stops the muons and remaining hadrons, leaving an almost pure muon neutrino beam with an intrinsic electron neutrino component of the order of a percent, which comes from the decays of kaons and muons, and an intrinsic muon antineutrino component of the order of 6% [33]. The beam stability and direction are monitored by the muon monitor [34,35] which measures the muons of energies higher than about 5 GeV that are able to penetrate the beam dump, and also by INGRID, the on-axis near detector, which samples the neutrino beam 280 m from the proton beam target [36].

B. Off-axis near detector ND280

The ND280 is composed of a series of subdetectors located 280 m from the target in a direction making a 2.5° angle with the average neutrino beam direction and placed within the refurbished UA1/NOMAD magnet which generates a 0.2 T magnetic field. The neutrino beam first passes through the pi-zero detector (P0D) [37] and then the tracker detector, which is used for the present measurement. The tracker is made up of two fine grained detectors (FGD) [38] and three time projection chambers (TPC) [39]. Those detectors are surrounded by electromagnetic calorimeters (ECals) [40] and side muon range detectors (SMRDs) [41], as can be seen in Fig. 1. The ND280 reference system is also shown in Fig. 1: the muon angle is defined as the polar angle (θ) between the muon momentum and the z axis (which corresponds, to a good approximation, to the beam direction and thus to the average neutrino direction). In general, most of the muons are expected to be “forward” ($\theta < 90^\circ$), i.e. to move in the same direction as the incoming neutrino beam, while events with “backward” muons ($\theta > 90^\circ$) correspond typically to interactions with high transferred Q^2 .

In the tracker, the target mass is provided by the FGDs. The first FGD (FGD1) is made only of scintillator bars, while the second FGD (FGD2) is made of alternating layers of scintillator bars and water. To measure cross sections on carbon, neutrino interactions occurring in FGD1 are selected. The FGD1 fiducial volume has an elemental composition of 86.1% carbon and 7.35% hydrogen with remaining contributions from oxygen (3.70%) and small




FIG. 1. Exploded view of the ND280 off-axis detector from Ref. [31]. Only one half of the UA1 magnet yoke is shown in this figure.

quantities of other elements (Ti, Si, N) [38,42]. The number of neutrons (nucleons) in the fiducial volume is 2.75×10^{29} (5.50×10^{29}). By chance, the excess of neutrons present in titanium (which is in the coating of the scintillator bars) is roughly equal to the number of protons from hydrogen, leading to the total number of protons and neutrons being equal (which is not what would be expected for the scintillator itself— C_8H_8). While the neutrino target for the present result is FGD1 with its complicated elemental composition, the measurement can be effectively considered as on carbon; the difference to the results from correcting for the other components is expected to be negligible with respect to the other normalization uncertainties.

III. ANALYSIS STRATEGY

The measurement of the ν_μ double-differential flux-integrated cross section of the charged current process on carbon without pions in the final state ($CC0\pi$) is presented as a function of the muon momentum and angle. Two analyses have been performed which make use of different selections and different cross-section extraction methods. The two analyses labelled I and II are considered to be equally valid, and no primary analysis was chosen. The consistency between their results is an indication of the robustness of the measurement. Particular care has been taken to perform a measurement that is highly model independent:

- (i) the cross section is measured as a function of the kinematics of the outgoing muon, as opposed to reconstructed variables that relate to the neutrino, such as the neutrino energy or the transferred momentum squared, which would depend on assumptions made about the nuclear model;
- (ii) a flux-integrated cross section is extracted, rather than a flux-averaged or flux-unfolded cross section, thus avoiding E_ν -dependent flux corrections. In

particular a flux-averaged cross section, as a function of muon kinematics, would require one to correct each bin in muon momentum and angle by the integral of the neutrino flux in that particular bin. In order to perform this correction, model-dependent assumptions are needed to establish the minimum and maximum neutrino energy allowed for each configuration in muon kinematics.

- (iii) the signal is defined in terms of the particles which exit the nucleus and can be observed in the detector. Compared to a signal defined in terms of interactions at the nucleon level, this removes the dependence on the modeling of the reinteractions of the final state particles in the nuclear medium. The definition used here includes CCQE interactions, but also events where, for example, one pion is produced at the interaction point and then reabsorbed in the nuclear environment;
- (iv) the cross-section measurement is designed to be robust to background-modeling uncertainties through the use of control samples or a reduced phase space, thereby removing regions with a small signal-to-background ratio.

The first analysis (“Analysis I”) uses a dedicated selection for CCQE-like events where a single muon (with a proton above or below detection threshold) is required and no other tracks. The cross section is extracted through a binned likelihood fit. The second analysis (“Analysis II”) follows the T2K oscillation analysis and MiniBooNE selection strategy, where CCQE-like interactions are identified by vetoing the presence of pions in the final state and Bayesian unfolding is used to correct for detector effects.

A. Event samples and simulation

1. Data samples

The analyses use data from the three T2K run periods between November 2010 and May 2013, where T2K was operating with a beam of mostly muon neutrinos. Only data recorded with all detectors correctly working are used, corresponding to 5.734×10^{20} protons on target (POT).

2. Monte Carlo samples

In order to correct for the detector response, acceptance and efficiency, simulations have been produced which correspond to ten times the data POT used, where the specific detector and beam configuration during each data run was modeled. The flux of neutrinos reaching the detectors—assuming the absence of oscillations—is predicted using simulations tuned to external measurements. Details of the simulation can be found in Ref. [43]. Interactions of protons in the graphite target and the resulting hadron production are simulated using the FLUKA 2008 package [44,45], weighted to match measurements of hadron production [46–50]. The propagation and

decay of those hadrons is performed in a GEANT3 [51] simulation which uses the GCALOR package [52] to model hadron reinteractions and decays outside the target. Uncertainties on the proton beam properties, horn current, hadron production model and alignment are taken into account to produce an energy-dependent systematic uncertainty on the neutrino flux. Flux tuning using NA61/SHINE data [46,50,53] reduces the uncertainty on the overall normalization of the integrated flux to 8.5%.

Neutrinos are then propagated through the ND280 detector and interactions are simulated with the NEUT neutrino event generator. NEUT 5.1.4.2 [54,55] uses the Llewellyn-Smith CCQE neutrino-nucleon cross-section formalism [56] with the relativistic Fermi gas model by Smith and Moniz [4,5] as the nuclear model. Dipole forms were used for both the axial-vector and vector form factors. From tuning to Super-Kamiokande atmospheric data and K2K data, the nominal axial mass M_A^{QE} was set to 1.21 GeV. Neutrino-induced pion production is simulated based on the Rein Sehgal model [57] in NEUT with the axial mass $M_A^{\text{RES}} = 1.21$ GeV. The parton distribution function GRV98 [58] with corrections by Bodek and Yang [59] is used for the deep inelastic scattering interactions. Secondary interactions of pions inside the nucleus [so-called final state interactions (FSIs)] are simulated using an intranuclear cascade model based on the method of Oset [60], tuned to external π - ^{12}C data.

The GENIE neutrino generator v2.6.4 [61] has been used as an alternative simulation to test the dependence of the analyses on the assumed signal and background models, with the primary difference to NEUT arising from different values of $M_A^{\text{QE}} = 0.99$ GeV [7] and $M_A^{\text{RES}} = 1.12$ GeV [62].

3. Event preselection

The FGD1 detector is used as the target for the neutrino interactions, and particles are reconstructed in the FGD1 itself and in TPC2, which is situated immediately downstream from FGD1. Initially, a ν_μ charged current selection is performed by looking for a muon candidate. The preselection criteria require good detector and beam conditions, correct timing with respect to the beam bunches, and good track reconstruction quality in TPC. The muon candidate is identified as the highest-momentum negatively charged track which passes the TPC track quality cut and starts in the FGD fiducial volume (FV). Further cuts are applied to remove events where the interaction happens outside the FV.

The further event selection depends on the analysis strategy and will be explained in the sections below. After the full selection cuts are applied, the background comes from events with one, or a number of, true pions which are misidentified or not reconstructed ($\text{CC}1\pi$ and CC_{other}), neutral current interactions (NC) and interactions that occurred outside of the fiducial volume but were reconstructed inside (OOFV).

4. Control samples for detector systematics

The detector systematics are described in detail in [2]. The systematics on track efficiency in the FGD and TPC, particle identification, charge identification, and momentum scale and resolution were evaluated by dedicated data and simulation comparisons using independent control samples (specially selected event samples that are designed to be sensitive to specific sources of uncertainty). In addition to reconstruction-related detector uncertainties, we also estimated the uncertainties on the number of simultaneous events (pileup) and OOFV events. Pions and nucleons from initial neutrino interactions in FGD1 can reinteract and be absorbed further in the detector. While the intranuclear final state interactions are simulated by the NEUT generator, the secondary interactions (SI) which follow these from pions and nucleons are treated by GEANT4.9.4 [63]. This additional uncertainty has also been evaluated from a control sample and is one of the dominant detector systematics.

5. Uncertainties due to neutrino interaction model

Uncertainties on the neutrino interaction model are described in detail in Ref. [2]. A set of systematic parameters characterizes the uncertainties on the predictions of the NEUT generator and are propagated through the analyses to estimate the uncertainty on the background and signal modeling, as well as the effect of the final state interactions. A number of those parameters are normalization uncertainties for the different interaction modes simulated by NEUT (energy dependent for the dominant modes at the T2K neutrino energy spectrum). Other parameters describe uncertainties on the values of the axial mass (using separate parameters for CCQE and resonant interactions), of the binding energy and of the Fermi momentum. An additional systematic parameter covers the difference between the predictions obtained with the default relativistic Fermi gas model used by NEUT and a spectral function describing the momentum and energy of nucleons inside the nucleus [64].

Finally, for the analysis using information from the presence or absence of a reconstructed proton to separate events between different categories (Analysis I, described below), the effects of the reinteractions of produced protons in the nuclear medium are evaluated using the GENIE neutrino interaction generator [61] by varying the parameters of the intranuclear cascade model describing those final state interactions.

B. Analysis I

This analysis uses a binned likelihood fit performed simultaneously in four signal regions and two control regions to constrain the backgrounds caused by resonant pion production and deep inelastic scattering (DIS).

The signal includes events where the muon is reconstructed in the FGD but does not reach the TPC, thus increasing the efficiency for muons that have small momentum or are emitted at high angle. Signal events with and without a reconstructed proton are treated separately, thus allowing the proton FSI parameters to be constrained using data. This is a first step towards a future differential measurement as a function of the proton kinematics.

The normalization and the shape of the background are extracted from data: the various background processes are parametrized in the same way as for the T2K oscillation analysis [2] and the values of such parameters are constrained by simultaneously fitting the control regions.

A likelihood fit is performed to the number of $CC0\pi$ events, in bins of muon momentum and angle. Detector, flux, and model uncertainties are included as nuisance parameters and a penalty term is added to the likelihood to constrain detector and model uncertainties. All the systematics are evaluated with “toy” Monte Carlo experiments sampling over the values of the nuisance parameters, as described in detail in Sec. III B 3.

1. Event selection

The preselection described in Sec. III A 3 is extended to also include muons which do not reach the TPC (i.e., are FGD only or FGD plus the ECAL or the SMRD or both). In this case the proton needs to be reconstructed as a positively charged track in the TPC with a vertex in the FGD FV, where this track has to pass the TPC track quality cut and a protonlike PID is required. The preselected events are then divided into four signal regions:

region 1 single-track events with a muon candidate in the TPC,

region 2 two-track events with a muon and a proton candidates in TPC,

region 3 two-track events with a muon candidate in TPC and a proton candidate in FGD,

region 4 two-track events with a proton candidate in TPC and a muon candidate in FGD (possibly reaching the ECAL or the SMRD or both).

Muon and proton candidates are identified using the $\frac{dE}{dx}$ measurement in the TPC or the energy deposited in the FGD. The kinematics of the muon candidate in each selection region for the $CC0\pi$ signal and the various backgrounds are shown in Fig. 2. The selection is highly dominated by events with one reconstructed muon and no other tracks (region 1). The signal regions where the muon is reconstructed in the TPC (regions 1,2,3) have very similar momentum distributions, although events with a reconstructed proton (regions 2,3) tend to have muons at slightly larger angles, while the region with the muon in the FGD and the proton in the TPC (region 4) have muons with much smaller momenta and larger angles. The overall selection efficiency for the $CC0\pi$ selection is 39%; Fig. 3 shows the efficiency as a function of the muon candidate momentum and angle. The loss in

efficiency at low momentum is due to the detector threshold for muon tracking in FGD, while the loss for backward muons is due to limitations in the reconstruction algorithm. This is the first T2K ND280 analysis with a nonzero efficiency and non-negligible event statistics for backwards-going muons; future analyses will benefit from a new reconstruction algorithm with further improvements in backwards-track reconstruction.

Two additional control regions are selected to constrain charged current event rates with single-pion and multiple-pion production. After preselection, a reconstructed negative track in TPC with muonlike PID and a positive track in TPC with pionlike PID are required. Events with exactly two tracks are included in *region 5* ($CC1\pi$ control region) while events with more than two tracks are included in *region 6* (CC_{other} control region). The background composition of the control regions as a function of muon momentum is shown in Fig. 4. The fraction of $CC0\pi$ signal in the control regions is very low and the $CC1\pi$ (CC_{other}) purity is quite good in region 5 (region 6), thus allowing unbiased constraints to be put on the background shape and normalization.

2. Cross-section extraction

We perform a fit to the number of selected events as a function of the muon kinematic variables (p_μ and $\cos\theta_\mu$), simultaneously in the four signal and two control regions. The detector-related and theoretical systematic uncertainties are parametrized and included in the fit through nuisance parameters. The number of selected events in each signal region and in each bin of reconstructed kinematics j is computed as

$$N_j = \sum_i^{\text{true bins}} \left[c_i \left(N_i^{\text{MC } CC0\pi} \prod_a^{\text{model}} w(a_\alpha)_{ij}^{CC0\pi} \right) + \sum_k^{\text{bkg reactions}} N_i^{\text{MC bkg } k} \prod_a^{\text{model}} w(a_\alpha)_{ij}^k \right] t_{ij}^{\text{det}} r_j^{\text{det}}, \quad (1)$$

where i runs over the bins of the “true” muon kinematics prior to detector smearing effects and k runs over the background reactions ($CC1\pi$, CC_{other} etc.). The parameters of interest c_i adjust the Monte Carlo $CC0\pi$ cross section to match with the observation in data. The t_{ij}^{det} matrix describes the transfer from the true (i) to the reconstructed (j) muon kinematics bins, and the r_j^{det} vector contains the nuisance parameters in the fit describing the detector systematics and which are constrained by a prior covariance matrix. The product \prod_a^{model} runs over the systematics related to the theoretical modeling of signal and background. Each $w(a_\alpha)_{ij}^k$ term is a weighting function describing how the generated and reconstructed muon kinematics change (in bins i , j and for each signal and background process) as a function of the value of a particular theoretical parameter a_α . The vector a_α contains all the nuisance





FIG. 2. Distribution of events in different regions for Analysis I. Each row corresponds to a signal region from 1 (top) to 4 (bottom). Figures in the left column are plotted against the reconstructed muon momentum and the right column against the reconstructed muon $\cos\theta$. Histograms are stacked.

parameters related to modeling systematics in the fit, constrained by a prior covariance matrix.

For simplicity, we use the same binning for the fit to the reconstructed p_μ , $\cos\theta_\mu$ distribution and for the extraction of the data/Monte Carlo cross-section corrections c_i . A nonrectangular p_μ , $\cos\theta_\mu$ binning (different p_μ binning for each $\cos\theta_\mu$ bin) is chosen on the basis of the available event numbers (which are much smaller in the high angle and

backward regions), the signal-over-background ratio (which is much smaller in the high momentum region) and of the detector resolution (to avoid large migrations of events between nearby bins).

As the parameters of interest for the fit rescale the overall number of CC0 π events in the four signal regions and two control regions together, the resulting cross section is extracted inclusively for all the regions simultaneously.

FIG. 3. Efficiency of reconstruction and selection of CC0 π events for Analysis I.

A future analysis will measure separate cross sections, with and without reconstructed protons.

A binned likelihood fit is performed

$$\chi^2 = \chi^2_{\text{stat}} + \chi^2_{\text{syst}} = \sum_j^{\text{reco bins}} 2 \left(N_j - N_j^{\text{obs}} + N_j^{\text{obs}} \ln \frac{N_j^{\text{obs}}}{N_j} \right) + \chi^2_{\text{syst}}, \quad (2)$$

where χ^2_{syst} is a penalty term for the systematics:

$$\chi^2_{\text{syst}} = (r_j^{\text{det}} - r_j^{\text{det,prior}})(V_{j,j'}^{\text{det}})^{-1}(r_j^{\text{det}} - r_j^{\text{det,prior}}) + (a_\alpha^{\text{model}} - a_\alpha^{\text{model,prior}})(V_{\alpha,\alpha'}^{\text{model}})^{-1}(a_\alpha^{\text{model}} - a_\alpha^{\text{model,prior}}) \quad (3)$$

where r_j^{det} and a_α^{model} are the parameters for detector and theory systematics running over the reconstructed bins j , and $r_j^{\text{det,prior}}$, $a_\alpha^{\text{model,prior}}$ indicate our initial knowledge of the detector response and theory parameters and $V_{j,j'}^{\text{det}}$, $V_{\alpha,\alpha'}^{\text{model}}$ are the corresponding covariance matrices.

The number of selected events as a function of the “true” kinematics extracted from the fit

$$N_i^{\text{CC0}\pi} = \sum_t^{\text{regions}} \sum_j^{\text{reco bins}} c_i N_{jt}^{\text{MC CC0}\pi} t_{tij}^{\text{det}} r_{tj}^{\text{det}} \prod_\alpha^{\text{model syst}} w(a_\alpha)_{ij}^{\text{CC0}\pi} \quad (4)$$

is corrected by the selection efficiency in each bin ϵ_i and divided by the overall integrated flux and the number of




FIG. 4. Distribution of events in different control regions for Analysis I. The first row is the CC1 π control region (region 5) and the second row is the DIS control region (region 6). Figures in the left column are plotted against the reconstructed muon momentum and the right column against the reconstructed muon $\cos\theta$. Histograms are stacked.

nucleons in the fiducial volume to extract a flux-integrated cross section:

$$\frac{d\sigma}{dx_i} = \frac{N_i^{\text{CC}0\pi}}{\epsilon_i \Phi N_{\text{nucleons}}^{\text{FV}}} \times \frac{1}{\Delta x_i}. \quad (5)$$

3. Treatment of systematic uncertainties

The detector systematics are stored in a covariance matrix (V_{jk}^{det}) as uncertainties on the total number of reconstructed events in bins of reconstructed muon momentum and angle for each signal and control region.

The flux systematic uncertainties affect the measured cross section in two ways: they affect the fit, by varying the signal differently in each bin and varying the signal over background ratio bin by bin, and thus changing the shape of the measured cross section, and they also affect the overall cross-section normalization. These two contributions are treated separately in the analysis. While the fit to the control regions has the power to constrain the flux systematic uncertainties, the flux is not included as a nuisance parameter in the fit; this is to reduce the impact of model-dependent assumptions when the signal-over-background distribution is extrapolated from the control regions to the signal regions.

The systematics due to signal and background modeling are based on the parametrization discussed in Sec. III A 2. The systematics due to background modeling and pion and proton FSIs are included as nuisance parameters in the fit. The fit to the control regions reduces the background modeling and pion FSI systematics by about a factor of 4. Systematics related to signal cross-section modeling, on the other hand, are not constrained from data because including them in the fit with a specific parametrization (e.g.: M_A^{QE}) would introduce a model-dependent bias to the result. The effect of signal modeling on the estimation of the efficiency in Eq. (5) is therefore described by a large systematic uncertainty, without trying to constrain it from the fit to the data. The small uncertainty on the efficiency that arises from proton and pion FSIs is also included.

In summary, all the systematic uncertainties are included in the fit as nuisance parameters, except for signal modeling and flux uncertainties. Finally the effect from the statistical uncertainty on the Monte Carlo samples is included in the bin-by-bin efficiency in Eq. (5).

To evaluate all the systematic and statistical uncertainties, we produce a large number of toy experiments. To assess the statistical uncertainty, the number of reconstructed events in each bin is fluctuated according to the Poisson distribution in each of the toy experiments. To evaluate the systematics, the values of the parameters governing the various systematic uncertainties are varied in each toy data set according to a Gaussian distribution, following the prior covariance matrices.

A summary of statistical and systematic errors is shown in Fig. 5. Theoretical uncertainties for the background cross

section, pion FSIs and proton FSIs are varied together, while separate toy experiments are made for the signal-modeling systematics, the flux systematics and the detector systematics. With the chosen binning, the statistical uncertainty is dominant. The largest uncertainty is from the flux normalization (8.5%) but, being fully correlated between all bins, it does not affect the cross-section shape. The effect of the flux uncertainty on the cross-section shape is small ($\leq 1\%$) in the region relevant for the signal ($p_\mu \approx 0.3\text{--}1$ GeV) while it reaches 5%–10% at low and high momenta, where the magnitude of the effect is similar to that from the detector systematics. The systematics due to the model of background cross sections and pion FSIs are larger in the forward and high momentum regions where most of the background is located, but even in these regions they remain below 2%, due to the constraint of nuisance parameters from the control regions. The Monte Carlo statistical uncertainty on the efficiency is about 2% or lower in most of the bins, but it is as much as 4% in the lowest and highest momentum bins, where the statistical uncertainties from data are also large. The systematics on the efficiency due to signal modeling are typically about a few percent, except in the high-angle region where the efficiency is lower and therefore we depend more on the simulation to extrapolate to the full phase space. Finally the detector systematics are of the order of a few percent and become larger (up to 10%) in the low and high momentum regions, where the detector resolution and efficiency is less well known. The detector systematics are the dominant shape uncertainties in most of the phase space, except at very high angles where the uncertainty on the signal modeling is larger. For the final results, the systematics uncertainty is evaluated on a separate set of toy experiments by varying all the theory nuisance parameters and the flux parameters at once. The uncertainties due to detector systematics and the systematics on the efficiency due to Monte Carlo statistics and signal modeling are added in quadrature.

4. Results

The total signal cross section per nucleon integrated over all the muon kinematics phase space is

$$\sigma = (0.417 \pm 0.047(\text{syst}) \pm 0.005(\text{stat})) \times 10^{-38} \text{ cm}^2 \text{ nucleon}^{-1} \quad (6)$$

to be compared with the NEUT prediction of $0.444 \times 10^{-38} \text{ cm}^2 \text{ nucleon}^{-1}$. The uncertainty is dominated by the flux normalization systematics (8.5%), while other sources of systematic uncertainty are a few percent or less.

The double-differential flux-integrated cross section is shown in Fig. 6, comparing to prediction of NEUT and GENIE Monte Carlo (see also Fig. 17 in the Appendix where the high momentum bins up to 30 GeV are included). The systematic uncertainties and the data statistical




FIG. 5. Statistical and systematic uncertainties in bins of true muon kinematics for Analysis I. The flux normalization uncertainty is flat (8.5%) and not shown, the uncertainty due to MC statistics is negligible in all bins and not shown.




FIG. 6. Measured cross section with shape uncertainties (error bars: internal systematics, external statistical) and fully correlated normalization uncertainty (gray band). The results from the fit to the data are compared to predictions from NEUT (blue solid line), and from GENIE (green dashed line).

uncertainties are summed in quadrature and shown as error bars. The uncertainty related to the flux normalization is given as a gray band.

In Fig. 7 the results are compared to the model of Martini *et al.* [20,65] and of Nieves *et al.* [21,66] (with a cut on the three-momentum transfer of $q_3 < 1.2$ GeV). These models include corrections to the interaction for collective nuclear effects calculated with random phase approximation, as well as 2p2h contributions, i.e. neutrino interactions with nucleon-nucleon correlated pairs and meson exchange currents. These models do not include the contribution of CC1 π with pion reabsorption due to FSIs, but they do include the production of a Δ resonance followed by pionless decay. Moreover the region of very small transferred Q^2 (most forward muon angles and higher muon momentum) could be sensitive to the shell structure of the nucleus and to the excitations of low-lying giant resonances [30,67]. Therefore the comparison to data has been limited to muon momenta below 3 GeV. Finally, the results are compared to the same models with and without the 2p2h contribution in Figs. 8 and 9. Those results will be discussed in details in Sec. IV B, together with the results of the second analysis presented in the next section.

C. Analysis II

Analysis II makes use of a Bayesian unfolding procedure to extract the CC0 π differential cross section from a single selection which is designed around the vetoing of pions. The selection used is the same as that used for the near detector fits in recent T2K oscillation analyses. In this analysis there is no direct constraint on the background. The background uncertainties are taken from fits to external data from MiniBooNE performed by T2K, which are also used as priors in the fit of Analysis I.

1. Event selection

After the preselection described in Sec. III A 3, the selection is subdivided based on the observed number of pions as done in recent T2K oscillation analyses [2]. Charged pions are tagged by searching for either a pionlike track in the TPC, a pionlike track in the FGD, or a Michel electron from muon decay in the FGD. Neutral pions are tagged by searching for electronlike tracks in the TPC. Events with no evidence for pions, charged or neutral, are kept and placed in the so-called CC0 π signal sample. This event selection is found to be 72% pure according to our NEUT Monte Carlo prediction of background events.

Figure 10 shows the kinematics of the events selected as simulated by our NEUT prediction, and extracted from data. The Monte Carlo is shown divided by true final-state topology. In Fig. 11, the same is shown but the simulation is divided by true reaction type. As can be seen, the selection has no efficiency in the backward region. Regarding the main backgrounds, the contribution from Δ resonance (RES) is larger than the contribution due to CC1 π topology

while the DIS component is smaller than the CC_{Other} topology, clearly showing the importance of pion FSI effects.

2. Cross-section extraction

The Bayesian unfolding procedure as described by D'Agostini [68] is used to convert from reconstructed variables to true variables.

The inputs to the Bayesian unfolding are

- (i) $P(t_i) = N(t_i) / \sum_{\alpha}^{\text{true bins}} N_{t_\alpha}$, the prior probability of finding an event in true bin i .
- (ii) $P(r_j|t_i)$, the probability of an event being reconstructed in bin j , given it originated in true bin i .
- (iii) B_{r_j} , the predicted background in each bin.

Using these inputs, it is possible to define the efficiency of selecting events in true bin i ,

$$\epsilon_i = \sum_{\alpha}^{\text{reco bins}} P(r_\alpha|t_i) \quad (7)$$

and the prior probability of being reconstructed in bin j ,

$$P(r_j) = \sum_{\alpha}^{\text{true bins}} P(r_j|t_\alpha)P(t_\alpha) \quad (8)$$

$$= \frac{N_{r_j}}{\sum_{\alpha}^{\text{true bins}} N_{t_\alpha}}. \quad (9)$$

Applying Bayes' theorem to the probabilities we have, results in the “unsmearing” matrix, which gives the probability for an event originated in true bin, i , given it was reconstructed in bin j ,

$$P(t_i|r_j) = \frac{P(r_j|t_i)P(t_i)}{P(r_j)} \quad (10)$$

and we can then use this unsmearing matrix, along with the efficiency, to obtain the unfolded estimate of the true event distribution.

$$N_{t_i}^{\text{CC0}\pi} = \frac{1}{\epsilon_i} \sum_j P(t_i|r_j)(N_{r_j} - B_{r_j}). \quad (11)$$

The cross section is then calculated in the same manner as Analysis I—by scaling the unfolded true number of events in each bin by the flux, number of targets, efficiency, and bin width, as in Eq. (5).

It is possible to iterate this procedure by feeding the unfolded true distribution in the start as an updated prior. Fake data studies showed that the first iteration is sufficient to correct for detector effects, even when the prior and fake data were generated according to models chosen to have exaggerated differences between each other. This is because the reconstruction resolution is very good, with




FIG. 7. Measured cross section with shape uncertainties (error bars: internal systematics, external statistical) and fully correlated normalization uncertainty (gray band). The results from fit to the data are compared to the predictions from Nieves *et al.* (red dotted line), and from Martini *et al.* (red solid line).




FIG. 8. Measured cross section with shape uncertainties (error bars: internal systematics, external statistical) and fully correlated normalization uncertainty (gray band). The results from the fit to the data are compared to predictions from Martini *et al.* without 2p2h (black line), and with 2p2h (red line).




FIG. 9. Measured cross section with shape uncertainties (error bars: internal systematics, external statistical) and fully correlated normalization uncertainty (gray band). The results from the fit to the data are compared to predictions from Nieves *et al.* without 2p2h (black dashed line), and with 2p2h (red dotted line).




FIG. 10. Distribution of events in selection, separated by true final state topology, as a function of muon momentum on the left and muon angle on the right. In this selection there is negligible coverage at angles above 90 degrees.




FIG. 11. Distribution of events in selection, separated by true reaction type, as a function of muon momentum on the left and muon angle on the right.

over 60% of events being reconstructed in their true bin, and under 5% of events being reconstructed more than one bin away.

There are limitations to this approach. We note that while the technique outlined in Ref. [68] should in principle unfold both background and signal, for this analysis, the unfolding procedure was applied subsequent to background subtraction. Because the background is subtracted before unfolding, this method can yield negative cross sections in some bins that contain large backgrounds.

3. Treatment of systematic uncertainties

All systematic uncertainties were propagated using a sample of toy experiments that were generated assuming different underlying parameters. Each toy experiment was unfolded using the same algorithm, and the results were used to calculate a covariance matrix defined as

$$V_{ij} = \frac{1}{N} \sum_{s_n=1}^N (\sigma_i^{(s_n)} - \sigma_i^{\text{nominal}})(\sigma_j^{(s_n)} - \sigma_j^{\text{nominal}}), \quad (12)$$

where, for each source of uncertainty, labeled by s , N pseudoexperiments are performed, giving a new differential cross section $\sigma^{(s_n)}$ each time, and the nominal cross section in bin i is given by $\sigma_i^{\text{nominal}}$.

4. Region of reported results

The region below 0.2 GeV in muon momentum contains a significant amount of external backgrounds, and suffers from a very low efficiency due to reconstruction difficulties. For this reason, no result is reported below 0.2 GeV. This should not be interpreted as measuring zero cross section in this region, rather that the cross section has not been measured in this region.

5. Results

The total signal cross section per nucleon integrated over the restricted muon kinematics phase space ($p_\mu > 0.2$ GeV and $\cos \theta_\mu > 0.6$) is

$$\sigma = (0.202 \pm 0.036(\text{syst}) \pm 0.003(\text{stat})) \times 10^{-38} \text{ cm}^2 \text{ nucleon}^{-1} \quad (13)$$

to be compared with the NEUT prediction: $0.232 \times 10^{-38} \text{ cm}^2 \text{ nucleon}^{-1}$. The uncertainty is fully dominated by the flux normalization. When considering the full phase space the results agree well with those in Analysis I; however, they suffer from large uncertainties that arise from extrapolating beyond the visible phase space. Figure 12 shows the cross section as a function of momentum for different angular bins. Those results will




FIG. 12. Measured cross section with shape uncertainties (error bars: internal systematics, external statistical) and fully correlated normalization uncertainty (gray band) as a function of muon momentum for different angular bins.

be discussed together with the results of the first analysis in Sec. IV B.

IV. DISCUSSION

The results are only presented in the form of a double-differential cross section, and a total flux integrated cross

section per nucleon. Since Analysis II has no efficiency in the low-momentum and high-angle region, the phase space has been reduced in that analysis.

Cross sections are often calculated as a function of Q^2 , providing one-dimensional distributions which are useful for comparisons between models; however, because of the limited acceptance of ND280, this would make the result

very model dependent. Indeed, each Q^2 bin would contain contributions from events with different muon kinematics thus requiring very different efficiency corrections. For instance the efficiency for forward muons in ND280 is very good, while it is lower for backward tracks (zero in Analysis II). Therefore the efficiency correction in each bin of Q^2 depends strongly on the relative number of events with forward or backward muons, which depends in turn on the particular model assumption. Moreover, in fake data studies for Analysis II, it was shown that changing the assumed (prior) distribution in (p_μ, θ_μ) space could have a drastic effect on the shape and normalization in Q^2 , or p_μ , and θ_μ separately. The efficiency corrections in bins of (p_μ, θ_μ) are instead mostly model independent since these are the actual variables measured in the detector.

The cross section is reported for all events without any pions in the final state. Thus the signal includes contributions from $CC1\pi$ events where the pion has been reabsorbed through FSIs. Such contributions can only be estimated from Monte Carlo, and are accompanied by large uncertainties.

The limitations described before (limited acceptance and presence of irreducible background) and the uncertainties in the background and signal modeling make it particularly relevant to compare the results of analyses which try to address those problems in different ways. We will start by looking more in detail at the differences between the two analyses, before discussing their results.

A. Comparison between the analyses

The two analyses differ, most substantially, in the methods used for the estimation of backgrounds, signal selection, and cross-section extraction. Analysis I uses two control regions to constrain the background parametrization, while Analysis II makes use of Monte Carlo predictions to estimate the background. Given the discrepancy between the present modeling of the $CC1\pi$ process and the available measurements in MiniBooNE [9] and MINERvA [16,17], there are large uncertainties on this process and it is important to constrain them from T2K data. In Fig. 13 the results of the fit to the control regions in Analysis I are shown; the most visible effect is a reduction of the events with very forward muons in the $CC1\pi$ -dominated region. The overall normalization of the DIS-dominated region is also slightly increased. Figure 14 shows a comparison between data and MC for background-enhanced selections in Analysis II. The selections are orthogonal to the signal selection, with one enhanced in $CC1\pi^+$ events, and the other enhanced in multipion and deep inelastic scattering events. Although the data are systematically lower than the prediction, this difference is covered by the assumed uncertainties on the prediction. Analysis II includes, indeed, large uncertainties in the backgrounds that are subtracted, to cover possible Monte Carlo mismodeling of this type, and it has limited the phase space to above 0.2 GeV to avoid the region where the signal-over-background ratio is low.




FIG. 13. Results of the fit to the control regions: the distribution of selected events in the $CC1\pi$ control region (region 5, top) and in the DIS control region (region 6, bottom), as a function of muon momentum (left) and muon $\cos \theta$ (right). The data are shown in black (with statistical errors), Monte Carlo predictions before the fit are shown by the dotted blue line, and those after the fit are in solid red (with systematics errors indicated by the pink band).




FIG. 14. Background comparison for a selection enhanced in $CC1\pi^+$ (top), and a selection enhanced in multipion and deep inelastic scattering (bottom). The data and MC agree within the uncertainties assigned to the MC.

The event selection for Analysis I, based on track counting and explicit proton identification, results in a better $CC0\pi$ purity (87%), at the expense of a smaller efficiency (39%), with respect to the selection used in Analysis II (purity 72%, efficiency 48%). Analysis II is instead based on pion counting, and thus is required to take into account uncertainties due to possible mismodeling of pions in the detector. Because of the detector geometry, pion rejection is worse at lower momenta and higher angles. The track counting-based selection of Analysis I has also a smaller contamination from events with $CC1\pi$ interactions where the pion has been reabsorbed through FSIs (i.e., a larger $CCQE$ purity) with respect to Analysis II; the contribution of $CC1\pi$ production with pion absorption is of the order of a few percent, except in the forward bin (up to 15%) and in the first momentum bin of each angular bin (up to 50%). Analysis I includes events where the muon does not reach the TPC (region 4) thus increasing the coverage of the phase space for high-angle muons, but in this case a proton has to be reconstructed in the TPC to improve the purity. On the other hand, the explicit reconstruction and identification of protons in Analysis I can be affected by cross-section modeling and proton FSI uncertainties. The selection of Analysis II is not subject to uncertainties related to the proton kinematics, as it does not attempt to measure the protons in the event. In particular, events which have two protons, where both are energetic enough to result in a reconstructible track in the detector, are included in Analysis II while they are excluded in Analysis I. These events can be due to nucleon-nucleon correlations or to meson exchange current processes or can

be due to proton production through FSIs. Studies with generators which include these effects suggest that in this data set we would expect to see around 70 events with two visible protons (to be compared with a number of signal selected events of about 10000). This category of events will be studied in dedicated analyses in the future. It should be noted that although the event selections are different, there is an overlap of approximately 80% between the two samples.

For the differential cross-section extraction, Analysis II uses an unfolding method to correct for detector effects, as with previous cross-section analyses by T2K [42,69], MiniBooNE [9] and MINERvA [9,16,17]. Analysis I uses a likelihood fit instead that is similar to the one used for T2K oscillation analyses [2]. The likelihood fit in Analysis I allows the fit parameters describing the theory systematics and the detector systematics to be kept separate, in contrast to the unfolding procedure where the final result is a convolution of the detector and theory parameters. Given the present poor knowledge of the modeling of signal and background, the likelihood fit allows one to check that the systematics theory parameters converge to meaningful values. The unfolding procedure in Analysis II has the feature that the statistical and systematic error estimates depend on the amount of regularization (or number of iterations), which needs to be considered. On the other hand, given the complexity of the fit in Analysis I and the large number of nuisance parameters it includes, it is important to compare the results using an independent and simpler method, as represented by the unfolding used in Analysis II.




FIG. 15. Comparison between the two analyses performed for the regions where the angular binning is the same (the measurement from Analysis II is reported only above $p_\mu > 200$ MeV). A good level of agreement is found. Error bars include all uncertainties except for the flux normalization uncertainty.

Despite the aforementioned differences between the two strategies, the results from the two analyses are in good agreement (see Fig. 15). This is a strong demonstration of the model independence of the results. Moreover, both analyses were tested using different model assumptions, to understand the dependence of the result on the signal and background model. Different generators were used for these tests, such as GENIE [61] and NuWro [70], as well as reweightings of the nominal NEUT model as a function of E_ν , Q^2 , or $(p_\mu, \cos\theta_\mu)$. The results were found in good agreement in all the tests, and to be within the estimated uncertainties.

B. Discussion of the results

In Figs. 6–9 and Fig. 12 the results are compared to various models. The data agree, inside current uncertainties, with the NEUT and GENIE simulations. In NEUT, the value of the M_A^{QE} is higher than it is in GENIE to accommodate cross-section enhancement due to nuclear effects, as suggested by recent quasi-elastic-like measurements on bounded nucleons inside nuclei, as discussed in Sec. I. The data tend to lie in between the NEUT and GENIE predictions, indicating the presence of nuclear effects which cannot be completely reproduced by changing the M_A^{QE} value.

In more recent models from Martini *et al.* [20,65] and Nieves *et al.* [21,66] the CCQE parameters are tuned to deuterium neutrino-scattering data, to low energy pion scattering and electron scattering on nuclei and the nuclear effects are explicitly implemented in the form of long- and short-range nucleon-nucleon correlations. We find that predictions from these new models agree with the data; in particular, the data suggest the presence of 2p2h with respect to pure CCQE predictions with the random phase approximation (RPA). The cross-section enhancement due to 2p2h is broadly distributed in muon kinematics and both the models would provide a poor description of the data without it. The prediction for pure CCQE with RPA are similar between Martini *et al.* and Nieves *et al.* models, while the prediction for multinucleons interactions is about a factor 2 larger in the first. The data can accommodate a large multinucleons component but to extract a precise estimation of the latter, the component due to pion-production events with subsequent pion absorption in nuclear matter need to be included in the models. Quantifying the agreement between the data and the various models is, indeed, not straightforward. The correlations between uncertainties in different bins must be considered, but the experimental measurements are affected by theoretical systematics, which are especially




FIG. 16. Left: the dash-dotted line is the predicted CC0 π cross section from NEUT as a function of the neutrino energy. The shapes of the neutrino fluxes for T2K and MiniBooNE are also shown. Right: A comparison of the measured CC0 π flux integrated cross section from MiniBooNE and the two T2K measurements presented in this paper, using the full (Analysis I) and restricted (Analysis II) phase space. As on the left, the dash-dotted lines represent the prediction from NEUT for each respective measurement.

large in the backward region. On the other hand, the models have known limitations (the lack of FSIs and large uncertainties related to nuclear effects, in particular for the very forward region) which should be considered in the comparison to the data. For all these reasons, we do not attempt a quantitative comparison of the experimental results with the various models. Such phenomenological studies will be pursued further in the T2K collaboration, possibly combining measurements for different channels and targets.

In Fig. 16, the flux-integrated cross sections measured in this paper and the one measured by MiniBooNE [9] are compared with the NEUT prediction. The MINER ν A results [16,17] are not included since comparisons to these measurements would depend on model-dependent assumptions: the analysis presented in [17] includes only pure CCQE events after subtracting CC1 π events where the pion is absorbed by the nuclear medium, and the analysis in Ref. [16] only includes events with both the muon and proton being reconstructed, and thus is dependent on the modeling of nucleon FSIs and nuclear effects. We look forward to new double-differential CC0 π results from MINER ν A. In Fig. 16 the full-phase-space result from Analysis I and the restricted phase space result from Analysis II ($\cos\theta_\mu > 0$ and $p_\mu > 200$ MeV) are reported and compared to the MiniBooNE measurement. The three measurements are compatible with NEUT predictions within their uncertainties, but there is a trend across the measurements as their acceptances vary; MiniBooNE, which has 4π acceptance, tends to measure a larger cross section, compared to the NEUT prediction, than the T2K full-phase-space result, which in turn gives a larger cross section, compared to NEUT, than our restricted-phase-space result. The uncertainties are too large for any

conclusive statement; however, this may indicate that NEUT does not correctly describe the angular shape of the cross section, as can also be seen in Fig. 6 and Fig. 12 where the results from data tend to lie above (below) NEUT at high (small) angle. It is thus crucial for future measurements to significantly increase the high angle acceptance, as well as to report results in the restricted phase space where the experiments have good efficiency.

V. CONCLUSIONS

In modern experiments, which use relatively heavy nuclear targets, CCQE cross-section measurements, previously considered to be well understood, have been found to contain potentially significant contributions from nuclear effects that are not well known and difficult to disentangle experimentally. The narrow-band T2K off-axis beam, which has a peak energy of 0.6 GeV, provides a powerful probe to study these CCQE interactions. In this paper the measurements of interactions on carbon with the production of a muon with no associated pions is presented in the form of a double-differential, flux-integrated cross section.

In this paper, we presented the results of two analyses designed to be robust against the dependence on the signal model assumed in the analysis. The results show the presence of sizeable nuclear effects for all muon angles and momenta. Those effects can only partially be modeled by an effective parametrization using an increased value of M_A^{QE} . More recent models with explicit implementation of 2p2h contributions agree well with the data. We are however not currently able to select between the two 2p2h models considered in this paper, because of the uncertainties in the modeling of the signal and background reactions, in particular CC1 π events followed by pion




FIG. 17. Measured cross section with shape uncertainties (error bars: internal systematics, external statistical) and fully correlated normalization uncertainty (gray band). The results from the fit to the data are compared to predictions from NEUT (blue solid line), and from GENIE (green dashed line). The measurement is presented in a logarithmic scale including the high momentum bins up to 30 GeV.

absorption which cannot be experimentally separated from CCQE interactions.

Finally, the limitations in past and present analyses have been highlighted and discussed in the paper, together with suggestions on ways to address such issues. In particular, future measurements using the T2K off-axis beam will include more data and improved algorithms for backwards-going track and proton reconstruction, which will enable exclusive measurements of the muon and proton final state to further elucidate the nature of nuclear effects in neutrino interactions and possibly solve the present degeneracy between different models.

ACKNOWLEDGMENTS

We thank the J-PARC staff for the superb accelerator performance and the CERN NA61 collaboration for providing valuable particle production data. We acknowledge the support of MEXT, Japan; NSERC (Grant No. SAPPJ-2014-00031), NRC and CFI, Canada; CEA and CNRS/

IN2P3, France; DFG, Germany; INFN, Italy; National Science Centre (NCN), Poland; RSF, RFBR and MES, Russia; MINECO and ERDF funds, Spain; SNSF and SERI, Switzerland; STFC, UK; and DOE, USA. We also thank CERN for the UA1/NOMAD magnet, DESY for the HERA-B magnet mover system, NII for SINET4, the WestGrid and SciNet consortia in Compute Canada, and GridPP, UK. In addition, participation of individual researchers and institutions has been further supported by funds from ERC (FP7), H2020 RISE-GA644294-JENNIFER, EU; JSPS, Japan; Royal Society, UK; DOE Early Career program, USA.

APPENDIX: CROSS-SECTION MEASUREMENT AT HIGH MOMENTUM

The double-differential flux-integrated cross section extracted from Analysis I is compared to NEUT and GENIE Monte Carlo in Fig. 6 including high momentum bins up to 30 GeV.

-
- [1] M. H. Ahn *et al.* (K2K Collaboration), *Phys. Rev. D* **74**, 072003 (2006).
 - [2] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. D* **91**, 072010 (2015).
 - [3] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **110**, 161801 (2013).
 - [4] R. Smith and E. Moniz, *Nucl. Phys.* **B43**, 605 (1972).
 - [5] R. Smith and E. Moniz, *Nucl. Phys.* **B101**, 547 (1975).
 - [6] R. Gran *et al.* (K2K Collaboration), *Phys. Rev. D* **74**, 052002 (2006).
 - [7] K. S. Kuzmin, V. V. Lyubushkin, and V. A. Naumov, *Eur. Phys. J. C* **54**, 517 (2008).
 - [8] A. Bodek, S. Avvakumov, R. Bradford, and H. S. Budd, *Eur. Phys. J. C* **53**, 349 (2008).
 - [9] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **81**, 092005 (2010).
 - [10] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **88**, 032001 (2013).
 - [11] P. Adamson *et al.* (MINOS Collaboration), *Phys. Rev. D* **91**, 012005 (2015).
 - [12] J. L. Alcaraz-Aunión and J. Walding (SciBooNE Collaboration), *AIP Conf. Proc.* **1189**, 145 (2009).
 - [13] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. D* **92**, 112003 (2015).
 - [14] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. D* **91**, 112002 (2015).
 - [15] V. Lyubushkin *et al.* (NOMAD Collaboration), *Eur. Phys. J. C* **63**, 355 (2009).
 - [16] T. Walton *et al.* (MINERvA Collaboration), *Phys. Rev. D* **91**, 071301 (2015).
 - [17] G. Fiorentini *et al.* (MINERvA Collaboration), *Phys. Rev. Lett.* **111**, 022502 (2013).
 - [18] L. Fields *et al.* (MINERvA Collaboration), *Phys. Rev. Lett.* **111**, 022501 (2013).
 - [19] A. Bodek, H. S. Budd, and M. E. Christy, *Eur. Phys. J. C* **71**, 1726 (2011).
 - [20] M. Martini, M. Ericson, G. Chanfray, and J. Marteau, *Phys. Rev. C* **80**, 065501 (2009).
 - [21] J. Nieves, I. R. Simo, and M. V. Vacas, *Phys. Lett. B* **707**, 72 (2012).
 - [22] O. Benhar, A. Lovato, and N. Rocco, *Phys. Rev. C* **92**, 024602 (2015).
 - [23] A. M. Ankowski, *Phys. Rev. C* **86**, 024616 (2012).
 - [24] A. Butkevich, *Phys. Rev. C* **80**, 014610 (2009).
 - [25] A. Bodek, M. Christy, and B. Coopersmith, *Eur. Phys. J. C* **74**, 3091 (2014).
 - [26] T. Leitner, O. Buss, L. Alvarez-Ruso, and U. Mosel, *Phys. Rev. C* **79**, 034601 (2009).
 - [27] C. Maieron, M. Martinez, J. Caballero, and J. Udias, *Phys. Rev. C* **68**, 048501 (2003).
 - [28] A. Meucci, C. Giusti, and F. D. Pacati, *Nucl. Phys.* **A739**, 277 (2004).
 - [29] A. Lovato, S. Gandolfi, R. Butler, J. Carlson, E. Lusk, S. C. Pieper, and R. Schiavilla, *Phys. Rev. Lett.* **111**, 092501 (2013).
 - [30] V. Pandey, N. Jachowicz, T. Van Cuyck, J. Ryckebusch, and M. Martini, *Phys. Rev. C* **92**, 024606 (2015).
 - [31] K. Abe *et al.* (T2K Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **659**, 106 (2011).
 - [32] Y. Fukuda *et al.* (Super-Kamiokande Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **501**, 418 (2003).
 - [33] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. D* **87**, 012001 (2013); **87**, 019902(E) (2013).

- [34] K. Matsuoka *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **624**, 591 (2010).
- [35] K. Suzuki *et al.* (T2K Collaboration), *Prog. Theor. Exp. Phys.* **2015**, 053C01 (2015).
- [36] K. Abe *et al.* (T2K Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **694**, 211 (2012).
- [37] S. Assylbekov *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **686**, 48 (2012).
- [38] P. Amaudruz *et al.* (T2K ND280 FGD Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **696**, 1 (2012).
- [39] N. Abgrall *et al.* (T2K ND280 TPC Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **637**, 25 (2011).
- [40] D. Allan *et al.* (T2K UK Collaboration), *J. Instrum.* **8**, P10019 (2013).
- [41] S. Aoki, G. Barr, M. Batkiewicz, J. Blocki, J. Brinson *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **698**, 135 (2013).
- [42] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. D* **87**, 092003 (2013).
- [43] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. D* **87**, 012001 (2013).
- [44] A. Ferrari, P.R. Sala, A. Fasso, and J. Ranft, Reports No. CERN-2005-010, No. SLAC-R-773, and No. INFN-TC-05-11, 2005.
- [45] G. Battistoni, F. Cerutti, A. Fassò, A. Ferrari, S. Muraro, J. Ranft, S. Roesler, and P.R. Sala, *AIP Conf. Proc.* **896**, 31 (2007), we used FLUKA2008, which was the latest version at the time of this study. A new version, FLUKA2011, has been already released now and the comparison with data would be different.
- [46] N. Abgrall *et al.* (NA61/SHINE Collaboration), *Phys. Rev. C* **85**, 035210 (2012).
- [47] T. Eichten *et al.*, *Nucl. Phys.* **B44**, 333 (1972).
- [48] J. V. Allaby *et al.*, CERN Technical Report. 70-12, 1970.
- [49] I. Chemakin *et al.*, *Phys. Rev. C* **77**, 015209 (2008).
- [50] N. Abgrall *et al.* (NA61/SHINE Collaboration), *Eur. Phys. J. C* **76**, 84 (2016).
- [51] R. Brun, F. Carminati, and S. Giani, Report No. CERN-W5013, 1994.
- [52] C. Zeitnitz and T. A. Gabriel, *Nucl. Instrum. Methods Phys. Res., Sect. A* **349**, 106 (1994).
- [53] N. Abgrall *et al.* (NA61/SHINE Collaboration), *Phys. Rev. C* **84**, 034604 (2011).
- [54] Y. Hayato, *Nucl. Phys. B, Proc. Suppl.* **112**, 171 (2002).
- [55] Y. Hayato, *Acta Phys. Pol. B* **40**, 2477 (2009).
- [56] C. H. Llewellyn Smith, *Phys. Rep.* **3**, 261 (1972).
- [57] D. Rein and L. M. Sehgal, *Ann. Phys. (N.Y.)* **133**, 79 (1981).
- [58] M. Gluck, E. Reya, and A. Vogt, *Eur. Phys. J. C* **5**, 461 (1998).
- [59] A. Bodek and U.K. Yang, *AIP Conf. Proc.* **670**, 110 (2003).
- [60] L. L. Salcedo, E. Oset, M. J. Vicente-Vacas, and C. Garcia-Recio, *Nucl. Phys.* **A484**, 557 (1988).
- [61] C. Andreopoulos, A. Bell, D. Bhattacharya, F. Cavanna, J. Dobson *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **614**, 87 (2010).
- [62] K. S. Kuzmin, V. V. Lyubushkin, and V. A. Naumov, *Acta Phys. Pol. B* **37**, 2337 (2006).
- [63] S. Agostinelli *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [64] O. Benhar, A. Fabrocini, S. Fantoni, and I. Sick, *Nucl. Phys.* **A579**, 493 (1994).
- [65] M. Martini, M. Ericson, G. Chanfray, and J. Marteau, *Phys. Rev. C* **81**, 045502 (2010).
- [66] J. Nieves, F. Sanchez, I. Ruiz Simo, and M. Vicente Vacas, *Phys. Rev. D* **85**, 113008 (2012).
- [67] M. Martini and M. Ericson, *Phys. Rev. C* **90**, 025501 (2014).
- [68] G. D'Agostini, *Nucl. Instrum. Methods Phys. Res., Sect. A* **362**, 487 (1995).
- [69] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. Lett.* **113**, 241803 (2014).
- [70] T. Golan, C. Juszczak, and J. T. Sobczyk, *Phys. Rev. C* **86**, 015505 (2012).