



symmetry

IMPACT
FACTOR
2.2

CITESCORE
5.4

Article

A New Subclass of Analytic Functions Associated with the q -Derivative Operator Related to the Pascal Distribution Series

Ying Yang, Rekha Srivastava and Jin-Lin Liu

Special Issue

Geometric Function Theory and Special Functions II

Edited by

Prof. Dr. Erhan Deniz



<https://doi.org/10.3390/sym16030280>

Article

A New Subclass of Analytic Functions Associated with the q -Derivative Operator Related to the Pascal Distribution Series

Ying Yang ¹, Rekha Srivastava ²  and Jin-Lin Liu ^{3,*}

¹ Department of Mathematics, Maanshan Teacher's College, Maanshan 243000, China; graceyying@massz.edu.cn

² Department of Mathematics and Statistics, University of Victoria, Victoria, BC V8W 3R4, Canada; rekhas@math.uvic.ca

³ Department of Mathematics, Yangzhou University, Yangzhou 225002, China

* Correspondence: jlliu@yzu.edu.cn

Abstract: A new subclass $TX_q[\lambda, A, B]$ of analytic functions is introduced by making use of the q -derivative operator associated with the Pascal distribution. Certain properties of analytic functions in the subclass $TX_q[\lambda, A, B]$ are derived. Some known results are generalized.

Keywords: analytic function; q -derivative operator; differential subordination; Pascal distribution series; Hadamard product

MSC: Primary 30C45; 30C50

1. Introduction

Let H be the class of analytic functions g of the following form

$$g(z) = z + \sum_{k=2}^{\infty} a_k z^k \quad (z \in D = \{z \in \mathbb{C} : |z| < 1\}). \quad (1)$$

Also, let T be the subclass of H consisting of analytic functions which have the following form

$$g(z) = z - \sum_{k=2}^{\infty} |a_k| z^k \quad (z \in D). \quad (2)$$

If $g \in T$ satisfies the condition:

$$\left| \frac{g'(z) - 1}{(\mu - \nu)\tau - \nu[g'(z) - 1]} \right| < 1 \quad (z \in D),$$

then g is said to belong to the class $R^\tau(\mu, \nu)$ ($\tau \in \mathbb{C} \setminus \{0\}$, $-1 \leq \nu < \mu \leq 1$). The class $R^\tau(\mu, \nu)$ was defined by Dixit and Pal [1].

Let P denote the class of Caratéodory functions that are analytic in D and have the following form

$$h(z) = 1 + \sum_{k=1}^{\infty} h_k z^k \quad (3)$$

so that

$$\operatorname{Re}\{h(z)\} > 0 \quad (z \in D).$$

We now recall here the principle of subordination between analytic functions [2]. If g and h are two functions in H , we say that g is subordinate to h , written $g \prec h$ or $g(z) \prec h(z)$, if there is a function v which is analytic in D with $v(0) = 0$ and $|v(z)| < 1$,



Citation: Yang, Y.; Srivastava, R.; Liu, J.-L. A New Subclass of Analytic Functions Associated with the q -Derivative Operator Related to the Pascal Distribution Series. *Symmetry* **2024**, *16*, 280. <https://doi.org/10.3390/sym16030280>

Academic Editor: Erhan Deniz

Received: 25 January 2024

Revised: 15 February 2024

Accepted: 20 February 2024

Published: 28 February 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

such that $g(z) = h(v(z))$. Further, if h is univalent in D , then we have the equivalence: $g(z) \prec h(z)$ ($z \in D$) $\Leftrightarrow g(0) = h(0)$ and $g(D) \subset h(D)$.

Now we define q -derivative D_q ($0 < q < 1$) for $g \in H$:

$$(D_q g)(z) = \begin{cases} \frac{g(qz) - g(z)}{(q-1)z} & (z \neq 0) \\ g'(0) & (z = 0). \end{cases} \quad (4)$$

From (4), one can find that

$$D_q g(z) = 1 + \sum_{k=2}^{\infty} [k]_q a_k z^{k-1} \quad (5)$$

and

$$[k]_q = \frac{1 - q^k}{1 - q}. \quad (6)$$

From (6), we can see that if $q \rightarrow 1-$, then $[k]_q \rightarrow k$.

In [3], Jackson first showed the application of q -calculus. Since then, many scholars studied the applications of q -analysis in physics and mathematics ([4–26]). Very recently, Srivastava [2] investigated systematically the applications and mathematical explanation of the fractional q -calculus as well as the fractional q -derivative in GFT.

By using the q -derivative operator D_q , we now define a new class $TX_q[\lambda, A, B]$.

Definition 1. Let $g \in T$ and satisfy the following differential subordination:

$$\frac{\lambda z^3 (zD_q g(z))''' + (1 + 2\lambda)z^2 (zD_q g(z))'' + z(zD_q g(z))'}{\lambda z^2 (zD_q g(z))'' + z(zD_q g(z))'} \prec \frac{1 + Az}{1 + Bz} \quad (z \in D), \quad (7)$$

where $-1 \leq B < A \leq 1$ and $0 \leq \lambda \leq 1$, then g is said to be in $TX_q[\lambda, A, B]$.

If

$$P(Y = k) = \binom{m-1+k}{m-1} (1-s)^m s^k, \quad k = 0, 1, 2, \dots$$

for parameters s and m , respectively; then, Y is called to be the Pascal distribution. In [27] El-Deeb, Bulboacă, and Dziok defined the following power series whose coefficients are probabilities of the Pascal distribution:

$$\Psi_s^m(z) := z + \sum_{k=2}^{\infty} \binom{m-2+k}{m-1} (1-s)^m s^{k-1} z^k \quad (z \in D),$$

where $0 \leq s \leq 1$ and $m \geq 1$.

Very recently, Frasin and Darus [28] introduced a class $C_q(\lambda, \alpha)$ ($0 \leq \alpha < 1, 0 \leq \lambda \leq 1$) associated with the q -derivative operator. They considered the following series:

$$\Phi_s^m(z) := 2z - \Psi_s^m(z) = z - \sum_{k=2}^{\infty} \binom{m-2+k}{m-1} (1-s)^m s^{k-1} z^k \quad (z \in D) \quad (8)$$

and the linear operator $I_s^m : H \rightarrow H$ by the Hadamard product:

$$I_s^m g(z) := g(z) * \Psi_s^m(z) = z + \sum_{k=2}^{\infty} \binom{m-2+k}{m-1} (1-s)^m s^{k-1} a_k z^k \quad (z \in D), \quad (9)$$

where $0 \leq s \leq 1$ and $m \geq 1$.

Now, we recall the following Lemmas.

Lemma 1 ([1,29]). Let $g \in R^{\tau}(\mu, \nu)$, then

$$|a_k| \leq (\mu - \nu) \frac{|\tau|}{k}, \quad k = 2, 3, \dots.$$

Lemma 2. Let $g(z) = z - \sum_{k=2}^{\infty} |a_k| z^k \in T$. If

$$\sum_{k=2}^{\infty} k[k]_q (\lambda k - \lambda + 1) \left[k - 1 + \frac{A - B}{1 + |B|} \right] |a_k| \leq \frac{A - B}{1 + |B|} \quad (z \in D),$$

where $0 \leq \lambda \leq 1$ and $-1 \leq B < A \leq 1$, then $g \in TX_q[\lambda, A, B]$.

Lemma 2 can be proved by using the same way as in [13].

In this article, we will derive certain properties of analytic functions in $TX_q[\lambda, A, B]$. Some known results are also generalized.

In order to facilitate our calculations and proofs, we derive several identities which hold for $m \geq 1$ and $0 \leq s < 1$ as the following:

$$\sum_{k=0}^{\infty} \binom{m-1+k}{m-1} s^k = (1-s)^{-m},$$

$$\sum_{k=2}^{\infty} \binom{m-2+k}{m-1} s^{k-1} = \sum_{k=0}^{\infty} \binom{m-1+k}{m-1} s^k - 1 = (1-s)^{-m} - 1, \quad (10)$$

$$\sum_{k=2}^{\infty} (k-1) \binom{m-2+k}{m-1} s^{k-1} = sm \sum_{k=0}^{\infty} \binom{m+k}{m} s^k = s \binom{m}{m-1} (1-s)^{-(m+1)}, \quad (11)$$

$$\sum_{k=3}^{\infty} (k-2)(k-1) \binom{m-2+k}{m-1} s^{k-1} = 2s^2 \binom{1+m}{m-1} (1-s)^{-(m+2)}, \quad (12)$$

$$\sum_{k=4}^{\infty} (k-3)(k-2)(k-1) \binom{m-2+k}{m-1} s^{k-1} = 6s^3 \binom{2+m}{m-1} (1-s)^{-(m+3)}, \quad (13)$$

and

$$\sum_{k=5}^{\infty} (k-4)(k-3)(k-2)(k-1) \binom{m-2+k}{m-1} s^{k-1} = 24s^4 \binom{3+m}{m-1} (1-s)^{-(m+4)}. \quad (14)$$

2. Main Results

Theorem 1. Let $m \geq 1$. If

$$\Psi(m, \lambda, s, A, B) \leq \frac{A - B}{1 + |B|}, \quad (15)$$

then the function Φ_s^m belongs to $TX_q[\lambda, A, B]$, where

$$\begin{aligned} \Psi(m, \lambda, s, A, B) := & 24\lambda \frac{\binom{3+m}{m-1} s^4}{(1-s)^{m+4}} + 6 \left[1 + \lambda \left(\frac{A-B}{1+|B|} + 8 \right) \right] \frac{\binom{2+m}{m-1} s^3}{(1-s)^{m+3}} \\ & + 2 \left[5 + \frac{A-B}{1+|B|} + \lambda \left[5 \left(\frac{A-B}{1+|B|} \right) + 14 \right] \right] \frac{\binom{1+m}{m-1} s^2}{(1-s)^{m+2}} \\ & + \left[4 + 3 \left(\frac{A-B}{1+|B|} \right) + 4\lambda \left[1 + \frac{A-B}{1+|B|} \right] \right] \frac{\binom{m}{m-1} s}{(1-s)^{1+m}}. \end{aligned}$$

Proof. From Lemma 2 and (8), we need to show that

$$P_q := \sum_{k=2}^{\infty} [k]_q k \left[\frac{A-B}{1+|B|} + k - 1 \right] (\lambda k + 1 - \lambda) \binom{k+m-2}{m-1} (1-s)^m s^{k-1} \leq \frac{A-B}{1+|B|}.$$

□

Letting $q \rightarrow 1-$, we obtain from (6) that $P_q \leq P_1$ and

$$\begin{aligned} P_1 &= \sum_{k=2}^{\infty} k^2 \left[k - 1 + \frac{A-B}{1+|B|} \right] (\lambda k + 1 - \lambda) \binom{k+m-2}{m-1} (1-s)^m s^{k-1} \\ &= \sum_{k=2}^{\infty} \left[k^3 + \left(\frac{A-B}{1+|B|} - 1 \right) k^2 \right] (\lambda k + 1 - \lambda) \binom{k+m-2}{m-1} (1-s)^m s^{k-1} \\ &= \sum_{k=2}^{\infty} \left\{ \lambda k^4 + \left[1 - \lambda \left(2 - \frac{A-B}{1+|B|} \right) \right] k^3 - (1-\lambda) \left(1 - \frac{A-B}{1+|B|} \right) k^2 \right\} \binom{m-2+k}{m-1} (1-s)^m s^{k-1}. \end{aligned}$$

By using the identities

$$k^2 = 1 + 3(k-1) + (k-2)(k-1), \tag{16}$$

$$k^3 = 1 + 7(k-1) + 6(k-2)(k-1) + (k-3)(k-2)(k-1), \tag{17}$$

$$k^4 = 1 + 15(k-1) + 25(k-2)(k-1) + 10(k-3)(k-2)(k-1) + (k-4)(k-3)(k-2)(k-1) \tag{18}$$

and (10)–(13), we have

$$\begin{aligned} P_1 &= \sum_{k=2}^{\infty} \left\{ \lambda k^4 + \left[1 - \lambda \left(2 - \frac{A-B}{1+|B|} \right) \right] k^3 + (\lambda - 1) \left(1 - \frac{A-B}{1+|B|} \right) k^2 \right\} \binom{m-2+k}{m-1} s^{k-1} (1-s)^m \\ &= \lambda \sum_{k=5}^{\infty} (k-4)(k-3)(k-2)(k-1) \binom{m-2+k}{m-1} s^{k-1} (1-s)^m \\ &\quad + \left\{ 10\lambda + \left[1 - \lambda \left(2 - \frac{A-B}{1+|B|} \right) \right] \right\} \sum_{k=4}^{\infty} (k-3)(k-2)(k-1) \binom{m-2+k}{m-1} s^{k-1} (1-s)^m \\ &\quad + \left\{ 25\lambda + 6 \left[1 - \lambda \left(2 - \frac{A-B}{1+|B|} \right) \right] + (\lambda - 1) \left(1 - \frac{A-B}{1+|B|} \right) \right\} \cdot \\ &\quad \sum_{k=3}^{\infty} (k-2)(k-1) \binom{m-2+k}{m-1} s^{k-1} (1-s)^m \\ &\quad + \left\{ 15\lambda + 7 \left[1 - \lambda \left(2 - \frac{A-B}{1+|B|} \right) \right] + 3(\lambda - 1) \left(1 - \frac{A-B}{1+|B|} \right) \right\} \cdot \\ &\quad \sum_{k=2}^{\infty} (k-1) \binom{m-2+k}{m-1} s^{k-1} (1-s)^m \\ &\quad + \left\{ \lambda + \left[1 - \lambda \left(2 - \frac{A-B}{1+|B|} \right) \right] + (1-\lambda) \left(\frac{A-B}{1+|B|} - 1 \right) \right\} \sum_{k=1}^{\infty} \binom{m-2+k}{m-1} s^{k-1} (1-s)^m \\ &= 24\lambda \frac{\binom{3+m}{m-1} s^4}{(1-s)^4} + 6 \left[1 + \lambda \left(\frac{A-B}{1+|B|} + 8 \right) \right] \frac{\binom{2+m}{m-1} s^3}{(1-s)^3} \\ &\quad + 2 \left[5 + \frac{A-B}{1+|B|} + \lambda \left(5 \left(\frac{A-B}{1+|B|} \right) + 14 \right) \right] \frac{\binom{1+m}{m-1} s^2}{(1-s)^2} \\ &\quad + \left[4 + 3 \left(\frac{A-B}{1+|B|} \right) + 4\lambda \left(\frac{A-B}{|B|+1} + 1 \right) \right] \frac{\binom{m}{m-1} s}{1-s} + \left(\frac{A-B}{|B|+1} \right) (1 - (1-s)^m) \\ &= (1-s)^m \Psi(m, \lambda, s, A, B) + \left(\frac{A-B}{|B|+1} \right) (1 - (1-s)^m). \end{aligned}$$

Now, we can find that $P_1 \leq \frac{A-B}{1+|B|}$ if (15) holds true. This proves the Theorem. According to Theorem 1, the following Corollaries are derived.

Corollary 1. Let $m \geq 1$. If

$$6 \frac{\binom{2+m}{m-1} s^3}{(1-s)^{m+3}} + 2 \left[\frac{A-B}{1+|B|} + 5 \right] \frac{\binom{1+m}{m-1} s^2}{(1-s)^{m+2}} + \left[3 \left(\frac{A-B}{1+|B|} \right) + 4 \right] \frac{\binom{m}{m-1} s}{(1-s)^{m+1}} \leq \frac{A-B}{1+|B|},$$

then the function Φ_s^m belongs to $TX_q[0, A, B]$.

Corollary 2. Let $m \geq 1$. If

$$24\lambda \frac{\binom{3+m}{m-1} s^4}{(1-s)^{4+m}} + 6(1+\lambda(9-\alpha)) \frac{\binom{2+m}{m-1} s^3}{(1-s)^{3+m}} + 2(6-\alpha+\lambda(19-5\alpha)) \frac{\binom{1+m}{m-1} s^2}{(1-s)^{2+m}} + (7-3\alpha+4\lambda(2-\alpha)) \frac{\binom{m}{m-1} s}{(1-s)^{1+m}} \leq 1-\alpha,$$

then the function Φ_s^m belongs to $TX_q[\lambda, 1-2\alpha, -1]$, where $\alpha \in [0, 1)$.

Remark 1. Making $\lambda = 0$ in Corollary 2, we have a result given by Frasin and Darus in [28].

Theorem 2. Suppose that $m \geq 1$ and $f \in R^\tau(\mu, \nu)$. If

$$\Phi(\lambda, \mu, \nu, \tau, m, s, A, B) \leq \frac{A-B}{|B|+1}, \quad (19)$$

then $I_s^m f \in TX_q[\lambda, A, B]$, where the operator I_s^m is given by (9) and

$$\begin{aligned} \Phi(\lambda, \mu, \nu, \tau, m, s, A, B) := & (\mu - \nu) |\tau| \left[6\lambda \frac{\binom{2+m}{m-1} s^3}{(1-s)^3} + 2 \left[1 + \lambda \left(\frac{A-B}{1+|B|} + 4 \right) \right] \frac{\binom{1+m}{m-1} s^2}{(1-s)^2} \right. \\ & \left. + \left[1 + \left(\frac{A-B}{1+|B|} + 1 \right) (2\lambda + 1) \right] \frac{\binom{m}{m-1} s}{1-s} + (1 - (1-s)^m) \frac{A-B}{1+|B|} \right]. \end{aligned}$$

Proof. By using Lemma 2, we need only to prove that

$$Q_q := \sum_{k=2}^{\infty} [k]_q k \left[\frac{A-B}{|B|+1} + k-1 \right] (\lambda k - \lambda + 1) \binom{k+m-2}{m-1} (1-s)^m s^{k-1} |a_k| \leq \frac{A-B}{|B|+1}.$$

□

Since $f \in R^\tau(\mu, \nu)$, according to Lemma 1, we know that $|a_n| \leq (\mu - \nu) \frac{|\tau|}{n}$, $n \in N \setminus \{1\}$. Letting $q \rightarrow 1-$, we find from (6) that $Q_q \leq Q_1$ and

$$\begin{aligned} Q_1 & \leq (\mu - \nu) |\tau| \sum_{k=2}^{\infty} k \left[k - 1 + \frac{A-B}{|B|+1} \right] (\lambda k - \lambda + 1) \binom{k+m-2}{m-1} (1-s)^m s^{k-1} \\ & = (\mu - \nu) |\tau| \sum_{k=2}^{\infty} \left[k^2 + k \left(\frac{A-B}{|B|+1} - 1 \right) \right] (\lambda k + 1 - \lambda) \binom{m+k-2}{m-1} (1-s)^m s^{k-1} \\ & = (\mu - \nu) |\tau| \sum_{k=2}^{\infty} \left\{ \lambda k^3 + k^2 \left[1 - \lambda \left(2 - \frac{A-B}{|B|+1} \right) \right] \right. \\ & \quad \left. + (1-\lambda) k \left(\frac{A-B}{|B|+1} - 1 \right) \right\} \binom{m+k-2}{m-1} (1-s)^m s^{k-1}. \end{aligned}$$

By considering $k = 1 + (k - 1)$, (16) and (17), we get

$$\begin{aligned}
 Q_1 &\leq (\mu - \nu)|\tau| \sum_{k=2}^{\infty} \left[\lambda k^3 + k^2 \left[1 - \lambda \left(2 - \frac{A-B}{|B|+1} \right) \right] \right. \\
 &\quad \left. + k(\lambda - 1) \left(1 - \frac{A-B}{|B|+1} \right) \right] \binom{m-2+k}{m-1} (1-s)^m s^{k-1} \\
 &= (\mu - \nu)|\tau| \left\{ \sum_{k=4}^{\infty} (k-3)(k-2)(k-1)\lambda \binom{m-2+k}{m-1} (1-s)^m s^{k-1} \right. \\
 &\quad + \left[6\lambda + \left(1 - \lambda \left(2 - \frac{A-B}{|B|+1} \right) \right) \right] \sum_{k=3}^{\infty} (k-2)(k-1) \binom{m-2+k}{m-1} (1-s)^m s^{k-1} \\
 &\quad + \left[7\lambda + 3 \left(1 - \lambda \left(2 - \frac{A-B}{|B|+1} \right) \right) + \left(1 - \frac{A-B}{|B|+1} \right) (\lambda - 1) \right] \cdot \\
 &\quad \sum_{k=2}^{\infty} (k-1) \binom{m-2+k}{m-1} s^{k-1} (1-s)^m \\
 &\quad + \left[\lambda + \left(1 - \lambda \left(2 - \frac{A-B}{|B|+1} \right) \right) + \left(1 - \frac{A-B}{|B|+1} \right) (\lambda - 1) \right] \cdot \\
 &\quad \left. \sum_{k=2}^{\infty} (k-1) \binom{m-2+k}{m-1} s^{k-1} (1-s)^m \right\} \\
 &= (\mu - \nu)|\tau| \left\{ 6\lambda \frac{\binom{2+m}{m-1} s^3}{(1-s)^3} + 2 \left(\lambda \left(\frac{A-B}{|B|+1} + 4 \right) + 1 \right) \frac{\binom{1+m}{m-1} s^2}{(1-s)^2} \right. \\
 &\quad \left. + \left((2\lambda + 1) \left(\frac{A-B}{|B|+1} + 1 \right) + 1 \right) \frac{\binom{m}{m-1} s}{1-s} + (1 - (1-s)^m) \frac{A-B}{|B|+1} \right\}.
 \end{aligned}$$

Thus we obtain that $Q_1 \leq \frac{A-B}{1+|B|}$ if (19) holds true. This proves the Theorem. According Theorem 2, we have the following corollary.

Corollary 3. Suppose that $m \geq 1$ and $f \in R^\tau(\mu, \nu)$. If the inequality

$$(\mu - \nu)|\tau| \left[\frac{2 \binom{1+m}{m-1} s^2}{(1-s)^2} + \left(2 + \frac{A-B}{|B|+1} \right) \frac{\binom{m}{m-1} s}{1-s} + (1 - (1-s)^m) \frac{A-B}{|B|+1} \right] \leq \frac{A-B}{|B|+1}$$

holds true, then $I_s^m f \in TX_q[0, A, B]$, where I_s^m is given by (9).

Theorem 3. Let $m \geq 1$ and the function Γ_s^m be given by

$$\Gamma_s^m(z) = \int_0^z \frac{\Phi_s^m(w)}{w} dw \quad (z \in D).$$

If

$$\Theta(m, \lambda, s, A, B) \leq \frac{A-B}{1+|B|}, \tag{20}$$

then $\Gamma_s^m \in TX_q[\lambda, A, B]$, where

$$\Theta(m, \lambda, s, A, B) := 6\lambda \frac{\binom{2+m}{m-1} s^3}{(1-s)^{m+3}} + 2 \left[1 + \lambda \left(\frac{A-B}{1+|B|} + 4 \right) \right] \frac{\binom{1+m}{m-1} s^2}{(1-s)^{m+2}} \\ + \left(1 + (2\lambda + 1) \left(\frac{A-B}{1+|B|} + 1 \right) \right) \frac{\binom{m}{m-1} s}{(1-s)^{m+1}}.$$

Proof. From (8), we get

$$\Gamma_s^m(z) = \int_0^z \frac{\Phi_s^m(t)}{t} dt = z - \sum_{k=2}^{\infty} k^{-1} \binom{m-2+k}{m-1} s^{k-1} (1-s)^m z^k \quad (z \in D).$$

□

According to Lemma 2, we find that $\Gamma_s^m \in TX_q[\lambda, A, B]$ if

$$R_q := \sum_{k=2}^{\infty} [k]_q k (\lambda k - \lambda + 1) \left[\frac{A-B}{|B|+1} + k - 1 \right] \binom{m-2+k}{m-1} s^{k-1} (1-s)^m \leq \frac{A-B}{|B|+1}.$$

Now letting $q \rightarrow 1-$, we have from (6) that $R_q \leq R_1$ and

$$R_1 = \sum_{k=2}^{\infty} k \left[k - 1 + \frac{A-B}{|B|+1} \right] (\lambda k - \lambda + 1) \binom{m-2+k}{m-1} (1-s)^m s^{k-1} \\ = \sum_{k=2}^{\infty} (\lambda k - \lambda + 1) \left[k^2 + \left(\frac{A-B}{|B|+1} - 1 \right) k \right] \binom{m-2+k}{m-1} s^{k-1} (1-s)^m \\ = \sum_{k=2}^{\infty} \left\{ \lambda k^3 + \left[1 - \lambda \left(2 - \frac{A-B}{|B|+1} \right) \right] k^2 + k(\lambda - 1) \left(1 - \frac{A-B}{|B|+1} \right) \right\} \binom{m-2+k}{m-1} (1-s)^m s^{k-1}.$$

By using the same method as in Theorem 2, we find that $\Gamma_s^m \in TX_q[\lambda, A, B]$ if (20) holds true. Thus, the Theorem is proved.

Corollary 4. Let $m \geq 1$. If

$$2 \binom{1+m}{m-1} s^2 (1-s)^{-(m+2)} + \left(2 + \frac{A-B}{1+|B|} \right) \binom{m}{m-1} s (1-s)^{-(m+1)} \leq \frac{A-B}{1+|B|},$$

then $\Gamma_s^m \in TX_q[0, A, B]$.

3. Conclusions

In recent years, many scholars (see, e.g., [30–32]) have been devoted to applications of q -analysis in physics and mathematics. In particular, Srivastava [2] systematically investigated the applications and mathematical explanation of the fractional q -calculus as well as the fractional q -derivative in GFT. In this paper, a new analytic function class $TX_q[\lambda, A, B]$ associated with the q -derivative operator and the Pascal distribution series is introduced and studied. Certain properties of functions in $TX_q[\lambda, A, B]$ are derived.

Scholars may consider some new q -analogous derivative operators and utilize these new operators to introduce new subclasses of analytic functions as potential avenues for future investigation. Also, the concepts given in this paper offer potential for extending to other operators such as the symmetric q -derivative operator. In particular, scholars could consider the symmetric q -derivative operator with differential subordination to define new subclasses of analytic functions.

Author Contributions: Writing—original draft, Y.Y., Writing—review & editing, J.-L.L., Supervision, R.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Data are contained within the article.

Acknowledgments: The authors would like to express their sincere thanks to the referees for their careful reading and suggestions which helped us to improve the paper.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Dixit, K.K.; Pal, S.K. On a class of univalent functions related to complex order. *Indian J. Pure Appl. Math.* **1995**, *26*, 889–896.
2. Srivastava, H.M. Operators of basic (or q -) calculus and fractional q -calculus and their applications in geometric function theory of complex analysis. *Iran. J. Sci. Technol. Trans. A Sci.* **2020**, *44*, 327–344. [[CrossRef](#)]
3. Jackson, F.H. On q -functions and a certain difference operator. *Earth Environ. Sci. Trans. R. Soc.* **1909**, *46*, 253–281. [[CrossRef](#)]
4. Ahuja, O.P.; Çetinkaya, A.; Polatoğlu, Y. Bieberbach-de Branges and Fekete-Szegő inequalities for certain families of q -convex and q -close-to-convex functions. *J. Comput. Anal. Appl.* **2019**, *26*, 639–649.
5. Aldweby, H.; Darus, M. On Fekete-Szegő problems for certain subclasses defined by q -derivative. *J. Funct. Spaces* **2017**, *2017*, 7156738. [[CrossRef](#)]
6. Aldweby, H.; Darus, M. A note on q -integral operators. *Electron. Notes Discret. Math.* **2018**, *67*, 25–30. [[CrossRef](#)]
7. Al-Hawary, T.; Yousef, F.; Frasin, B.A. Subclasses of analytic functions of complex order involving Jackson (p, q)-derivative. In Proceedings of International Conference on Fractional Differentiation and Its Applications (ICFDA), Ajman, United Arab Emirates, 14–16 March 2018.
8. Aouf, M.K.; Mostafa, A.O.; Al-Quhali, F.Y. Properties for class of bi-uniformly univalent functions defined by Salagean type q -difference operator. *Int. J. Open Probl. Complex Anal.* **2019**, *11*, 2.
9. Araci, S.; Duran, U.; Acikgoz, M.; Srivastava, H.M. A certain (p, q)-derivative operator and associated divided differences. *J. Inequal. Appl.* **2016**, *2016*, 301. [[CrossRef](#)]
10. Frasin, B.A.; Murugusundaramoorthy, G. A subordination results for a class of analytic functions defined by q -differential operator. *Ann. Univ. Paedagog. Cracov. Stud. Math.* **2020**, *19*, 53–64. [[CrossRef](#)]
11. Ibrahim, R.W.; Darus, M. On a class of analytic functions associated with a complex domain concerning q -differential-difference operator. *Adv. Differ. Equ.* **2019**, *2019*, 515. [[CrossRef](#)]
12. Ismail, M.E.H.; Merkes, E.; Steyr, D. A generalization of starlike functions. *Complex Var. Theory* **1990**, *14*, 77–84. [[CrossRef](#)]
13. Kamali, M.; Akbulut, S. On a subclass of certain convex functions with negative coefficients. *J. Appl. Math. Comput.* **2002**, *145*, 341–350. [[CrossRef](#)]
14. Khan, B.; Srivastava, H.M.; Khan, N.; Darus, M.; Tahir, M.; Ahmad, Q.Z. Coefficient estimates for a subclass of analytic functions associated with a certain leaf-like domain. *Mathematics* **2020**, *8*, 1334. [[CrossRef](#)]
15. Mahmood, S.; Sokół, J. New subclass of analytic functions in conical domain associated with Ruscheweyh q -differential operator. *Res. Math.* **2017**, *71*, 1345–1357. [[CrossRef](#)]
16. Murugusundaramoorthy, G.; Selvaraj, C.; Babu, O.S. Subclasses of starlike functions associated with fractional q -calculus operators. *J. Complex Anal.* **2013**, *2013*, 572718. [[CrossRef](#)]
17. Ramachandran, C.; Soupramanien, T.; Frasin, B.A. New subclasses of analytic function associated with q -difference operator. *Eur. J. Pure Appl.* **2017**, *10*, 348–362.
18. Rehman, M.S.U.; Ahmad, Q.Z.; Srivastava, H.M.; Khan, N.; Darus, M.; Khan, B. Applications of higher-order q -derivatives to the subclass of q -starlike functions associated with the Janowski functions. *AIMS Math.* **2021**, *6*, 1110–1125. [[CrossRef](#)]
19. Srivastava, H.M. Some generalizations and basic (or q -) extensions of the Bernoulli, Euler and Genocchi polynomials. *Appl. Math. Inf. Sci.* **2011**, *5*, 390–444.
20. Srivastava, H.M.; Bansal, D. Close-to-convexity of a certain family of q -Mittag-Leffer functions. *J. Nonlinear Var. Anal.* **2017**, *1*, 61–69.
21. Srivastava, H.M.; Zahoor, Q.; Khan, N.; Khan, N.; Khan, B. Hankel and Toeplitz determinants for a subclass of q -starlike functions associated with a general conic domain. *Mathematics* **2019**, *7*, 181. [[CrossRef](#)]
22. Srivastava, H.M.; Raza, N.; Jarad, E.S.A.A.; Srivastava, G.; AbuJarad, M.H. Fekete-Szegő inequality for classes of (p, q)-starlike and (p, q)-convex functions. *Rev. Real Acad. Cienc. Exactas Fís. Nat. Ser. A Mat.* **2019**, *113*, 3563–3584. [[CrossRef](#)]
23. Srivastava, H.M.; Tahir, M.; Khan, B.; Ahmad, Q.Z.; Khan, N. Some general classes of q -starlike functions associated with the Janowski functions. *Symmetry* **2019**, *11*, 292. [[CrossRef](#)]
24. Srivastava, H.M.; Tahir, M.; Khan, B.; Ahmad, Q.Z.; Khan, N. Some general families of q -starlike functions associated with the Janowski functions. *Filomat* **2019**, *33*, 2613–2626. [[CrossRef](#)]
25. Srivastava, H.M. An introductory overview of Bessel polynomials, the generalized Bessel polynomials and the q -Bessel polynomials. *Symmetry* **2023**, *15*, 822. [[CrossRef](#)]
26. Srivastava, R.; Zayed, H.M. Subclasses of analytic functions of complex order defined by q -derivative operator. *Stud. Univ. Babeş-Bolyai Math.* **2019**, *64*, 69–78. [[CrossRef](#)]
27. El-Deeb, S.M.; Bulboacă, T.; Dziok, J. Pascal distribution series connected with certain subclasses of univalent functions. *Kyungpook Math. J.* **2019**, *59*, 301–314.
28. Frasin, B.A.; Darus, M. Subclass of analytic functions defined by q -derivative operator associated with Pascal distribution series. *AIMS Math.* **2021**, *6*, 5008–5019. [[CrossRef](#)]

29. Silverman, H. Univalent functions with negative coefficients. *Proc. Am. Math. Soc.* **1975**, *51*, 109–116. [[CrossRef](#)]
30. Khan, N.; Khan, S.; Xin, Q.; Tchier, F.; Malik, S.N.; Javed, U. Some applications of analytic functions associated with q -fractional operator. *Mathematics* **2023**, *11*, 930. [[CrossRef](#)]
31. Srivastava, H.M.; Khan, N.; Darus, M.; Khan, S.; Ahmad, Q.Z.; Hussain, S. Fekete-Szegö type problems and their applications for a subclass of q -starlike functions with respect to symmetrical points. *Mathematics* **2020**, *8*, 842. [[CrossRef](#)]
32. Khan, M.F.; Goswami, A.; Khan, S. Certain new subclass of multivalent q -starlike functions associated with q -symmetric calculus. *Fractal Fract.* **2022**, *6*, 367. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.