

---

Faculty of Science

Faculty Publications

---

Coral Oxygen Isotopic Records Capture the 2015/2016 El Niño Event in the Central Equatorial Pacific

Gemma K. O'Connor, Kim M. Cobb, Hussein R. Sayani, Alyssa R. Atwood, Pamela R. Grothe, Samantha Stevenson, Julia K. Baum, Tianran Chen, Danielle C. Claar, Nicholas T. Hitt, Jean Lynch-Stieglitz, Richard A. Mortlock, Gavin A. Schmidt, Rachel Walter

December 2021

© 2021 O'Connor et al. This is an open access article distributed under the terms of the Creative Commons Attribution License. <https://creativecommons.org/licenses/by-nc-nd/4.0/>

This article was originally published at:

<https://doi.org/10.1029/2021GL094036>

---

Citation for this paper:

O'Connor, G. K., Cobb, K. M., Sayani, H. R., Atwood, A. R., Grothe, P. R., Stevenson, S., et al. (2021). Coral oxygen isotopic records capture the 2015/2016 El Niño event in the central equatorial Pacific. *Geophysical Research Letters*, 48(24). <https://doi.org/10.1029/2021GL094036>

# Geophysical Research Letters<sup>®</sup>



## RESEARCH LETTER

10.1029/2021GL094036

## Coral Oxygen Isotopic Records Capture the 2015/2016 El Niño Event in the Central Equatorial Pacific

### Key Points:

- An ensemble of coral oxygen isotope timeseries from the central equatorial Pacific tracks the 2015/16 El Niño event
- Coral oxygen isotope records reflect ~70% contribution from warming and ~30% from freshening during the 2015/16 El Niño event
- *In situ* seawater oxygen isotope data provide quantitative constraints on temperature versus hydrological contributions to coral records

### Supporting Information:

Supporting Information may be found in the online version of this article.














### Correspondence to:

G. K. O'Connor,  
[goconnor@uw.edu](mailto:goconnor@uw.edu)

### Citation:

O'Connor, G. K., Cobb, K. M., Sayani, H. R., Atwood, A. R., Grothe, P. R., Stevenson, S., et al. (2021). Coral oxygen isotopic records capture the 2015/2016 El Niño event in the central equatorial Pacific. *Geophysical Research Letters*, 48, e2021GL094036. <https://doi.org/10.1029/2021GL094036>

Received 29 APR 2021  
Accepted 26 AUG 2021

Gemma K. O'Connor<sup>1,2</sup> , Kim M. Cobb<sup>1</sup> , Hussein R. Sayani<sup>1</sup> , Alyssa R. Atwood<sup>3</sup> , Pamela R. Grothe<sup>4</sup> , Samantha Stevenson<sup>5</sup> , Julia K. Baum<sup>6</sup>, Tianran Chen<sup>7</sup> , Danielle C. Claar<sup>6,8</sup> , Nicholas T. Hitt<sup>9</sup> , Jean Lynch-Stieglitz<sup>1</sup> , Richard A. Mortlock<sup>10</sup> , Gavin A. Schmidt<sup>11</sup> , and Rachel Walter<sup>1</sup> 

<sup>1</sup>School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA, <sup>2</sup>Department of Earth and Space Sciences, University of Washington, Seattle, WA, USA, <sup>3</sup>Department of Earth, Ocean, and Atmospheric Science, Florida State University, Tallahassee, FL, USA, <sup>4</sup>Earth & Environmental Sciences, University of Mary Washington, Fredericksburg, VA, USA, <sup>5</sup>Bren School of Environmental Science and Management, University of California-Santa Barbara, Santa Barbara, CA, USA, <sup>6</sup>Department of Biology, University of Victoria, Victoria, BC, Canada, <sup>7</sup>South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, China, <sup>8</sup>School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA, USA, <sup>9</sup>Victoria University of Wellington, Wellington, New Zealand, <sup>10</sup>Department of Earth and Planetary Sciences, Rutgers, State University of New Jersey, Piscataway, NJ, USA, <sup>11</sup>NASA Goddard Institute for Space Studies, New York, NY, USA

**Abstract** Coral oxygen isotopes ( $\delta^{18}\text{O}$ ) from the central equatorial Pacific provide monthly resolved records of El Niño-Southern Oscillation activity over past centuries to millennia. However, calibration studies using *in situ* data to assess the relative contributions of warming and freshening to coral  $\delta^{18}\text{O}$  records are exceedingly rare. Furthermore, the fidelity of coral  $\delta^{18}\text{O}$  records under the most severe thermal stress events is difficult to assess. Here, we present six coral  $\delta^{18}\text{O}$  records and *in situ* temperature, salinity, and seawater  $\delta^{18}\text{O}$  data from Kiritimati Island (2°N, 157°W) spanning the very strong 2015/16 El Niño event. Local sea surface temperature (SST) anomalies of  $+2.4 \pm 0.4^\circ\text{C}$  and seawater  $\delta^{18}\text{O}$  anomalies of  $-0.19 \pm 0.02\text{‰}$  contribute to the observed coral  $\delta^{18}\text{O}$  anomalies of  $-0.58 \pm 0.05\text{‰}$ , consistent with a ~70% contribution from SST and ~30% from seawater  $\delta^{18}\text{O}$ . Our results demonstrate that Kiritimati coral  $\delta^{18}\text{O}$  records can provide reliable reconstructions even during the largest class of El Niño events.

**Plain Language Summary** Oxygen isotope anomalies in coral skeletons are a well-established proxy for changes in tropical Pacific Ocean temperature variations, which have a profound impact on weather extremes around the planet. However, only a handful of calibrations exist that quantify the relationship between ocean temperature and coral oxygen isotopic composition at a given site, especially across extreme events where this relationship may vary most strongly. Here we compare ocean temperature data from loggers installed on the reef at Kiritimati Island (2°N, 157°W) to coral oxygen isotopic records spanning the record-breaking 2015/16 El Niño event. We find that oxygen isotopes in corals provide accurate reconstructions of ocean temperature extremes during this very strong El Niño event, with ~70% of the signal originating from ocean temperature and the remainder from increased rainfall.

## 1. Introduction

The El Niño-Southern Oscillation (ENSO) is the dominant mode of interannual climate variability, but its response to greenhouse warming remains highly uncertain (Bellenger et al., 2014; Ng et al., 2021; Stevenson, 2012). Although projections of ENSO-related sea surface temperature (SST) variability differ across climate models, simulations forced with projected anthropogenic greenhouse gas emissions generally agree on an increase in the hydrological extremes associated with ENSO (Bonfils et al., 2015; Brown et al., 2020; Cai et al., 2014; Power et al., 2013). This tendency translates to an increase in the occurrence of ‘extreme’ El Niño events (as defined based on equatorial Pacific precipitation) under continued greenhouse warming (Cai et al., 2015), although models reflect a range of responses (Stevenson et al., 2021). Another question is whether greenhouse gas-driven changes to ENSO have already taken place; observational studies of 20th century ENSO extremes suggest a shift toward more Central Pacific warming events (Wang et al., 2019), consistent with results from analysis of a large network of coral records spanning the last several centuries (Freund et al., 2019). A number of paleoclimate

© 2021. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs License](https://creativecommons.org/licenses/by/4.0/), which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

ENSO reconstructions document enhanced late 20th-century ENSO variability relative to the preindustrial era (Cobb et al., 2013; Grothe et al., 2020; Li et al., 2013; Liu et al., 2017; McGregor et al., 2013), supporting the idea that greenhouse warming has already led to a significant shift in ENSO properties. However, it remains unclear how much of the observed shift in ENSO variance is due to a strengthening of SST anomalies or hydrological extremes during ENSO events. This is especially true for the largest trove of monthly resolved ENSO reconstructions based on modern and fossil corals from the central tropical Pacific, which documents a 25% increase in interannual variability from the last millennia to the most recent decades (Grothe et al., 2020).


Oxygen isotope ratios ( $\delta^{18}\text{O}$ ) in corals from the central tropical Pacific track ENSO-related changes in SST and hydrology, with warmer and wetter conditions during El Niño events driving negative coral  $\delta^{18}\text{O}$  anomalies, while cool and dry conditions during La Niña events drive positive coral  $\delta^{18}\text{O}$  anomalies (Cobb et al., 2001; Evans et al., 1999; Nurhati et al., 2009). Cores collected from living and fossil coral colonies have provided high-fidelity records of past ENSO activity over recent decades (Cobb et al., 2001; Nurhati et al., 2009, 2011), centuries (Cobb et al., 2003), and millennia (Cobb et al., 2013; Grothe et al., 2020; McGregor et al., 2013). The largest such data set comes from Kiritimati Island (2°N, 157°W), where ENSO dominates coral  $\delta^{18}\text{O}$  variability as evidenced by correlation coefficients of  $\sim 0.80$  between modern coral  $\delta^{18}\text{O}$  records and the NIÑO3.4 index (Grothe et al., 2020) - a key index of large-scale ENSO variability.

Despite their high-fidelity representation of instrumental SST variability over the satellite era, coral  $\delta^{18}\text{O}$ -based ENSO reconstructions are associated with a number of uncertainties. For one, extremely high SSTs during strong El Niño events can induce thermal stress, which slows or halts coral calcification, limiting the ability of coral proxies to capture the full extent of temperature SST change during such events (e.g., Carilli et al., 2017). Indeed, previous coral studies from the Eastern Pacific and Western Atlantic have found skeletal growth hiatuses during very strong El Niño events (Dunbar et al., 1994), as well as systematic geochemical biases initiated by extreme El Niño events (Hetzinger et al., 2016). Due to limited coral ensemble sizes and *in situ* climate data, it remains unknown how well corals from the central tropical Pacific record ENSO activity during extreme events. Second, coral  $\delta^{18}\text{O}$  values reflect the combined influence of SST and seawater  $\delta^{18}\text{O}$ , but the relative contribution of SST versus seawater  $\delta^{18}\text{O}$  to Kiritimati coral  $\delta^{18}\text{O}$  records during ENSO extremes is difficult to assess without *in situ* data. Previous work using paired Sr/Ca and coral  $\delta^{18}\text{O}$  measurements showed a stronger than normal seawater  $\delta^{18}\text{O}$  contribution in Kiritimati coral  $\delta^{18}\text{O}$  during the very strong 1997/98 El Niño event (McGregor et al., 2013), but coverage among multiple strong events paired with *in situ* data is needed to better understand these contributions. In particular, there are no systematic surveys of seawater  $\delta^{18}\text{O}$  variations across El Niño events, even though such variability has been inferred from both models and observational data (Conroy et al., 2014; Fairbanks et al., 1997; Russon et al., 2013).

In this study, we present six new coral  $\delta^{18}\text{O}$  records as well as *in situ* records of SST, salinity, and seawater  $\delta^{18}\text{O}$  from Kiritimati Island spanning the 2015/16 El Niño event, one of the strongest El Niño events on record. These data allow us to quantify the oceanographic changes that occurred across this event, and their contribution to the observed coral  $\delta^{18}\text{O}$  anomalies. By comparing our results from the 2015/16 El Niño event to available coral  $\delta^{18}\text{O}$  records and *in situ* SST, salinity, and seawater  $\delta^{18}\text{O}$  data from Kiritimati across past strong El Niño events of the 20th century, we assess the stability of the relationship between SST, seawater  $\delta^{18}\text{O}$ , and coral  $\delta^{18}\text{O}$  across different El Niño events. This assessment is key to improving interpretation of coral  $\delta^{18}\text{O}$  records from the region.

## 2. Methods

Six modern coral cores from *Porites* spp. were recovered from a leeward open ocean reef flat ranging from 6–9 m depth (labeled “drill site” on Figure S1 in the Supporting Information S1) at Kiritimati Island during expeditions in April and November 2016. The cores were prepared and analyzed for coral  $\delta^{18}\text{O}$  composition using standard procedures (Sayani et al., 2019, see Figure S6 in the Supporting Information S1 for X-rays) on either a Thermo-Finnigan Delta V or MAT253, both equipped with a Kiel IV Carbonate Device, with analytical precisions of  $\pm 0.05\text{‰}$  ( $1\sigma$ ) and  $\pm 0.06\text{‰}$  ( $1\sigma$ ), respectively. Age models for each record were reconstructed by peak matching the coral  $\delta^{18}\text{O}$  data with  $1^\circ \times 1^\circ$  monthly NOAA OISSTv2 SST data (Reynolds et al., 2002) from the grid box containing Kiritimati Island, following procedures outlined in Cobb (2002). Each monthly resolved record covers the 2015/16 El Niño event and extends back 3–9 years prior to the event, depending on core length (Table S1 in



**Figure 1.** Coral  $\delta^{18}\text{O}$  records (colored lines) from Kiritimati Island plotted with monthly resolved ERSSTv5 (gray dashed line) at Kiritimati Island (note inverted y-axis for coral  $\delta^{18}\text{O}$ ). (a) New modern coral  $\delta^{18}\text{O}$  records spanning the 2015/16 El Niño event as presented in this study. (b) Same as in (a), but with the ensemble mean of the corals shown. Coral  $\delta^{18}\text{O}$  offsets have been applied (Text S3 in the Supporting Information S1) and are denoted in the legend in panel (a) in units of per mil.

the Supporting Information S1). Following Sayani et al. (2019), we apply offsets of up to 0.19‰ to each record to align them to a common ensemble mean (see Text S1 and S3 in the Supporting Information S1).


We also present *in situ* SST, salinity, and seawater  $\delta^{18}\text{O}$  data spanning the 2015/16 El Niño event from Kiritimati Island. We present continuous time series of SST from a Seabird SBE56 temperature logger (Site 5 logger), and weekly SST and salinity from a Seabird SBE37 conductivity-temperature-depth (CTD) sensor (see Figure S1 in the Supporting Information S1 for locations). We present 97 paired salinity and seawater  $\delta^{18}\text{O}$  data from seawater samples collected during field expeditions to Kiritimati Island from 2014 to 2016, and ~30 paired salinity and seawater  $\delta^{18}\text{O}$  samples collected on the RV Moana Wave from October to November 1997. We include two *in situ* SST logger datasets from Kiritimati Island presented by Claar et al., 2019 in our analyses: the first from the drill site, where corals in this study were collected, and the second from a southward-facing reef flat several kilometers away (Figure S1 in the Supporting Information S1). We also use data from three gridded SST products, using the grid point closest to Kiritimati Island:  $1^\circ \times 1^\circ$  weekly OISSTv2,  $1^\circ \times 1^\circ$  monthly HadISSTv1.1 (Rayner et al., 2003), and  $2^\circ \times 2^\circ$  monthly ERSSTv5 (Huang et al., 2017).

### 3. Results

#### 3.1. Coral $\delta^{18}\text{O}$ Records

All six coral  $\delta^{18}\text{O}$  records closely follow instrumental SST and are highly reproducible on monthly to annual timescales (Figure 1). Most of the variability in coral  $\delta^{18}\text{O}$  is driven by ENSO, as seasonal SST variability at this site is relatively small. The records are well correlated with ERSSTv5, HadISST, and OISSTv2 ( $R$  between  $-0.81$  and  $-0.91$ ,  $p < 0.05$ , Table S8 in the Supporting Information S1). A composite coral  $\delta^{18}\text{O}$  record, formed by averaging overlapping records, has a greater correlation with these SST products than any individual record ( $R \sim -0.93$ ; Figure 1b, Table S8 in the Supporting Information S1). All coral  $\delta^{18}\text{O}$  records demonstrate consistent monthly variability across the study period, with correlations among records ranging between 0.64 and 0.90 ( $p < 0.05$ ). The  $1\sigma$  spread in coral  $\delta^{18}\text{O}$  values during any given month of overlapping coral  $\delta^{18}\text{O}$  records ranges from 0.04 to 0.24‰, with the maximum variance occurring in November 2015, during the peak of the 2015/16 El Niño event. We note that there is greater consistency during the 3-month peak of the mild El Niño event in 2014/15 ( $1\sigma = 0.10\%$ ).

During the 3-month peak of the event (October 2015 to December 2015), all six coral records exhibit significant coral  $\delta^{18}\text{O}$  depletion (mean of  $-5.59 \pm 0.05\%$  (1SE, Text S4 in the Supporting Information S1)), coinciding with warmer and fresher conditions (Figure 2). For the two-year baseline preceding the El Niño event (from January



**Figure 2.** *In situ* sea surface temperature (SST), salinity, and seawater (sw)  $\delta^{18}\text{O}$  data spanning the 2015/16 El Niño event at Kiritimati Island. (a) Weekly SST records from four *in situ* temperature loggers (see Figure S1 in the Supporting Information S1 for locations). Also plotted are weekly OISSTv2, monthly HadISSTv1.1, and monthly ERSSTv5. (b) Salinity records from the conductivity-temperature-depth (CTD; red) and seawater bottle samples (green circles). (c) Seawater  $\delta^{18}\text{O}$  measurements from seawater bottle samples (blue diamonds). Orange lines represent the median of the seawater  $\delta^{18}\text{O}$  data. The boxes show the 25%–75% interquartile range.

2013 to January 2015, ending 9 months prior to the 3-month peak), the coral  $\delta^{18}\text{O}$  records exhibit a mean isotopic value of  $-5.01 \pm 0.02\text{‰}$  (1 standard error (1SE)), reflecting a change in coral  $\delta^{18}\text{O}$  of  $-0.58 \pm 0.05\text{‰}$  (1SE).

### 3.2. *In Situ* SST, Salinity, and Seawater $\delta^{18}\text{O}$ Measurements

We compare *in situ* SST, salinity, and seawater  $\delta^{18}\text{O}$  from Kiritimati Island to constrain the individual contributions of SST and seawater  $\delta^{18}\text{O}$  to coral  $\delta^{18}\text{O}$  during the 2015/16 El Niño event. The four *in situ* SST datasets all capture the 2015/16 El Niño event with varying coverage (Figure 2a). The *in situ* SST timeseries exhibit strong correlations with all three gridded SST data products, on monthly timescales (R values of 0.90–0.99, Table S8 in the Supporting Information S1). We calculate change in SST across the 2015/16 El Niño by subtracting the 2-year-averaged baseline SST (2013–2015) from the 3-month peak of the event. Using just the Site 5 logger (which has the longest coverage across this event), we calculate an increase in SST of  $2.4 \pm 0.4^\circ\text{C}$  (1SE) at Kiritimati Island. In comparison, ERSSTv5 and HadISST show similar warming of  $2.4 \pm 0.4^\circ\text{C}$  (1SE) and  $2.4 \pm 0.3^\circ\text{C}$  (1SE), respectively, and OISSTv2 shows a slightly larger mean change of  $2.8 \pm 0.5^\circ\text{C}$  (1SE), still within the uncertainty ranges of the other products.


To constrain the hydrological contribution to the interannual variability in the coral records, we analyze *in situ* CTD salinity and salinity/seawater  $\delta^{18}\text{O}$  from seawater bottle samples (Figures 2b and 2c). Due to the shorter duration of these timeseries, we use a 1-year baseline from mid-2014 to mid-2015 (and the same 3-month peak) to calculate the change in salinity during the event. The CTD tracks a decrease in salinity of  $0.91 \pm 0.05$  psu (1SE), reaching a value of 34.14 psu during the peak of the El Niño event. Salinity data from 148 seawater bottle samples show a mean decrease of  $0.96 \pm 0.08$  psu (1SE), reaching a mean value of 34.41 psu during the peak of the event. Although the absolute salinity of the two data sources is offset by approximately 0.3 psu

(which may be attributed to differences in instrumental calibration), they capture the same relative change across the study interval. Using the same time periods as the salinity calculation, the seawater  $\delta^{18}\text{O}$  data capture a decrease of  $0.19 \pm 0.02\text{‰}$  (1SE), reaching a mean of 0.19‰ during the peak of the event. We note that the use of a 1-year baseline here may underestimate the magnitude of the El Niño anomalies, as there is less warming (and likely less freshening) from this 2014/15 baseline (Figure 1). We account for this in conclusions based on this calculation.

### 3.3. Comparison to Past Strong El Niño Events

Coral  $\delta^{18}\text{O}$  timeseries from Kiritimati Island show progressive depletion in mean coral  $\delta^{18}\text{O}$  during the peaks of the 1982/83, 1997/98, and 2015/16 strong El Niño events. Mean coral  $\delta^{18}\text{O}$  anomalies (3-month average centered around the peak of each event) were  $-5.23 \pm 0.07\text{‰}$  during the peak of the 1982/83 El Niño event, decreased to  $-5.49 \pm 0.10\text{‰}$  during the 1997/98 event, and then decreased to  $-5.59 \pm 0.05\text{‰}$  during the 2015/16 event (Figure 3). However, these El Niño are superimposed on background warming and freshening trends in the tropical Pacific (Nurhati et al., 2009), which produce more depleted coral  $\delta^{18}\text{O}$  values over this time interval. When the amplitude of each event is isolated from the mean state of the Pacific (by taking the difference between the 3-month average during the peak and the 2-year baseline prior to the event; Figure 3), a student *t*-test ( $p < 0.05$ ) shows no statistically significant difference in coral  $\delta^{18}\text{O}$  magnitude of these three events (Figure S3 in the Supporting Information S1).

The amplitude of change during the 1997/98 event, both in terms of coral  $\delta^{18}\text{O}$  and observed SST, is similar to that of the 2015/16 event. Five out of six corals that span the 1997/98 event show similar depletion in coral  $\delta^{18}\text{O}$  during the peak of the 1997/98 event. The sixth coral (Nurhati-09) does not fully capture SST and SSS changes across this event (Figure 3), possibly due to thermal stress or sublethal bleaching (Nurhati et al., 2009). Excluding this record, the amplitude of the 1997/98 event is  $-0.56 \pm 0.06\text{‰}$  (1SE), which is indistinguishable from the



**Figure 3.** Coral  $\delta^{18}\text{O}$  records (solid lines) and monthly sea surface temperature (SST; dashed lines) during the (a) 1982/83, (b) 1997/98, and (c) 2015/16 El Niño events. Horizontal bars at the top of each panel show the 2-year baseline (gray) and 3-month peak (red) periods used to calculate SST and coral  $\delta^{18}\text{O}$  anomalies for each El Niño event. Coral  $\delta^{18}\text{O}$  records shown are the “X16” corals (this study), “X12-3” and “X12-6” (Grothe et al., 2020), “X12-FS” fossil corals and “X12-D6-1” fossil coral (Hitt et al., 2021), “Nurhati-09” (Nurhati et al., 2009), and “Evans-99” (Evans et al., 1999). Coral records are plotted with offsets applied, as shown in Tables S1 and S2 in the Supporting Information S1. Gridded SST data are from ERSSTv5 and OISSTv2. Seawater  $\delta^{18}\text{O}$  (open circles) and sea surface salinity (open diamonds) measurements are shown where available.

$-0.58 \pm 0.05\text{‰}$  (1SE) amplitude of the 2015/2016 event. Similarly, ERSSTv5 shows statistically indistinguishable SST anomalies during the 1997/98 El Niño and 2015/16 El Niño ( $+2.1 \pm 0.8^\circ\text{C}$  vs.  $2.4 \pm 0.4^\circ\text{C}$ , respectively; Figure 3). Similar results are found when this calculation is performed with OISSTv2 and HadISST (Figure S4 and Table S8 in the Supporting Information S1).

As limited salinity and seawater  $\delta^{18}\text{O}$  data is available from the 1997/98 event, we only compare bottle seawater  $\delta^{18}\text{O}$  measurements from the peaks of the 1997/98 and 2015/16 events. Salinity during November 1997 reached a mean of 34.17 psu, slightly lower than that of the 2015/16 event mean in seawater bottles (34.41 psu). Seawater  $\delta^{18}\text{O}$  reached a minimum of  $0.11\text{‰}$  during October 1997, significantly less than the  $0.19\text{‰}$  observed during the peak of the 2015/16 event. Thus, available data show that the 1997/98 El Niño event was characterized by similar changes in coral  $\delta^{18}\text{O}$  and SST but slightly lower minimum salinity and seawater  $\delta^{18}\text{O}$  values compared to the 2015/16 event.

For the 1982/83 event, we use two of the three coral  $\delta^{18}\text{O}$  records that span this interval to calculate anomalies associated with this strong El Niño, excluding the Nurhati et al., 2009 record as the authors note that it does not fully capture SST and SSS changes across the peak of the event. We calculate a coral  $\delta^{18}\text{O}$  amplitude of  $-0.52 \pm 0.05\text{‰}$  (1SE) for the 1982/83 event, which is statistically indistinguishable from the 1997/98 and 2015/16 events (Figure S3 in the Supporting Information S1). ERSSTv5 shows an increase of  $+1.7 \pm 0.3^\circ\text{C}$  (1SE) at Kiritimati across the 1982/83 El Niño, again statistically indistinguishable from SST during the 1997/98 and 2015/16 events (Figure S3 in the Supporting Information S1) and consistent with previous work (Huang et al., 2016).

### 3.4. SST and Hydrological Contributions to Coral $\delta^{18}\text{O}$

To quantify the relative contributions of SST and seawater  $\delta^{18}\text{O}$  to coral  $\delta^{18}\text{O}$  during the 2015/16 El Niño event, we calculate the expected temperature contribution to coral  $\delta^{18}\text{O}$  using the empirical relationship of  $-0.2\text{‰ }^\circ\text{C}^{-1}$  (Epstein et al., 1951), and then subtract the temperature contribution from coral  $\delta^{18}\text{O}$  to isolate seawater  $\delta^{18}\text{O}$

**Table 1**

Comparison of Sea Surface Temperature (From ERSSTv5) and Estimated Seawater (sw)  $\delta^{18}\text{O}$  Contributions to Observed Coral  $\delta^{18}\text{O}$  Anomalies Associated With Strong El Niño Events at Kiritimati Island

		1982/83 event	1997/98 event	2015/16 event
A	Observed $\Delta$ coral $\delta^{18}\text{O}$	$-0.52 \pm 0.05\text{‰}$	$-0.56 \pm 0.06\text{‰}$	$-0.58 \pm 0.05\text{‰}$
B	Observed $\Delta$ SST (ERSSTv5)	$+1.7 \pm 0.3^\circ\text{C}$	$+2.1 \pm 0.8^\circ\text{C}$	$+2.4 \pm 0.4^\circ\text{C}$
C	Estimated SST-driven $\Delta$ coral $\delta^{18}\text{O}$ ( $C \times 0.2\text{‰}^\circ\text{C}^{-1}$ )	$-0.35 \pm 0.07\text{‰}$	$-0.42 \pm 0.16\text{‰}$	$-0.48 \pm 0.08\text{‰}$
D	Estimated $\Delta$ sw $\delta^{18}\text{O}$ (A–C)	$-0.18 \pm 0.07\text{‰}$	$-0.14 \pm 0.15\text{‰}$	$-0.10 \pm 0.08\text{‰}$
E	Observed $\Delta$ sw $\delta^{18}\text{O}$ (SW bottle samples)	N/A	N/A	$-0.19 \pm 0.02\text{‰}$
F	Estimated sw $\delta^{18}\text{O}$ contribution (D/A $\times$ 100)	$34 \pm 13\%$	$26 \pm 28\%$	$18 \pm 13\%$
G	Observed sw $\delta^{18}\text{O}$ contribution (E/A $\times$ 100)	N/A	N/A	$33 \pm 4\%$

*Note.* Changes are calculated as the 3-month peak of the event minus a 2-year baseline, reported with 1SE uncertainties (Text S4, S5, and Table S4 in the Supporting Information S1). For the 2015/16 event, observed changes in seawater  $\delta^{18}\text{O}$  and associated contributions are also shown, with 1SE uncertainty, calculated using the available 1-year baseline.

changes (Cahyarini et al., 2008; Ren et al., 2003, Text S5 in the Supporting Information S1). The SST change of  $+2.4 \pm 0.4^\circ\text{C}$  (1SE) calculated from ERSSTv5 during the 2015/16 event corresponds to an expected coral  $\delta^{18}\text{O}$  change of  $-0.48 \pm 0.08\text{‰}$  (1SE). As the observed coral  $\delta^{18}\text{O}$  change during this event is  $-0.58 \pm 0.05\text{‰}$  (1SE), we estimate a seawater  $\delta^{18}\text{O}$  contribution of  $-0.10 \pm 0.08\text{‰}$  (1SE), which implies that seawater  $\delta^{18}\text{O}$  accounts for  $18 \pm 13\%$  (1SE) of the observed coral  $\delta^{18}\text{O}$  change during the 2015/16 event, with SST contributing to the remaining  $82 \pm 13\%$  (1SE; Table 1 and S4 in the Supporting Information S1). We find similar results when repeating this calculation with OISSTv2 and HadISST (Tables S5–S8 in the Supporting Information S1). We note that SST uncertainty dominates this calculation, and a full propagation of the 2SE uncertainties through SST, coral  $\delta^{18}\text{O}$ , and seawater  $\delta^{18}\text{O}$  would likely double the range of permissible percentages.

The *in situ* seawater  $\delta^{18}\text{O}$  measurements provide a powerful additional constraint on the coral  $\delta^{18}\text{O}$  budget, given the weak constraints afforded by SST and coral  $\delta^{18}\text{O}$  observations outlined above. Given the observed change in  $\delta^{18}\text{O}_{\text{sw}}$  of  $-0.19 \pm 0.02\text{‰}$  (1SE) during the peak of the 2015/16 El Niño, the seawater  $\delta^{18}\text{O}$  contribution is constrained to  $\sim 30\text{--}35\%$  (including all values that fall within the 1 SE uncertainty range; Tables 1, S4, and Figure S5 in the Supporting Information S1). These values do not change if different SST products are used (Tables S5 and S6 in the Supporting Information S1), and are relatively insensitive to the choice of 1-year versus 2-year baselines for the calculations of anomalies (Table S7 in the Supporting Information S1). We note that the mild 2014/15 El Niño event is included in the 2-year baseline, which could bias the baseline toward slightly more warming than normal years (Figure 3).

Using the same approach, we calculate the SST and seawater  $\delta^{18}\text{O}$  contributions to coral  $\delta^{18}\text{O}$  changes across the 1997/98 and 1982/83 El Niño events (Tables 1 and S5 in the Supporting Information S1), although the lack of observed seawater  $\delta^{18}\text{O}$  data translates to large uncertainties in these estimates. For the 1997/98 event, we find a hydrological contribution of approximately  $26 \pm 28\%$  (1SE), which is similar in mean but associated with larger uncertainty. For the 1982/83 event, we find a seawater  $\delta^{18}\text{O}$  contribution of  $34 \pm 13\%$  (1SE). In short, we find consistent relative contributions from SST and seawater  $\delta^{18}\text{O}$  anomalies (roughly 70% and 30%, respectively) to coral  $\delta^{18}\text{O}$  anomalies across the three strong El Niño events in question.

#### 4. Discussion and Conclusions

Our analysis of Kiritimati coral records demonstrates that coral  $\delta^{18}\text{O}$  at this site reliably records SST and hydrological anomalies associated with the strongest El Niño events. Paired *in situ* temperature, seawater  $\delta^{18}\text{O}$ , and coral  $\delta^{18}\text{O}$  confirm that all six cores collected in 2016 accurately capture the SST and seawater  $\delta^{18}\text{O}$  changes observed during the 2015/2016 El Niño event, within uncertainties (Figure S5 in the Supporting Information S1). We observe no evidence of hiatuses in x-rays of our cores (Figure S6 in the Supporting Information S1), such as those found in Galapagos coral during strong El Niños (Dunbar et al., 1994; Jimenez et al., 2018), nor do we observe any signs of thermal stress in our coral  $\delta^{18}\text{O}$  data (e.g., Hetzinger et al., 2016). However, we observe the largest variance in coral  $\delta^{18}\text{O}$  during peak El Niño warming, which could bias studies of past El Niño amplitudes

based on single coral  $\delta^{18}\text{O}$  records. Nurhati et al., 2009 hypothesized that the attenuated signal of the 1997/98 El Niño event in their Kiritimati coral  $\delta^{18}\text{O}$  records may have resulted from a reduced precipitation or a growth hiatus. Alternatively, high coral  $\delta^{18}\text{O}$  variance during peak El Niño conditions may reflect more meter-scale variance in temperature and seawater  $\delta^{18}\text{O}$  during El Niño extremes, given the strong gradients that occur at this time between the surface and depth, and between the lagoon and the open ocean, during a prolonged period of reduced wind-driven mixing.

Our investigation of the relative contributions of SST and seawater  $\delta^{18}\text{O}$  anomalies to Kiritimati coral  $\delta^{18}\text{O}$  anomalies across the 2015/16 El Niño event shows that SST conditions are responsible for  $\sim 70\%$  of the coral  $\delta^{18}\text{O}$  signal. Similar estimates for SST contributions to Kiritimati coral  $\delta^{18}\text{O}$  records spanning the 1982/83 and 1997/98 El Niño events (mean values of  $66 \pm 13$  and  $74 \pm 28\%$ , respectively) bolsters our confidence in our findings from the 2015/16 El Niño event, although these estimates are associated with much larger uncertainties given the lack of *in situ* seawater  $\delta^{18}\text{O}$  observations. Our findings are consistent with isotope-enabled climate model studies (Russon et al., 2013) which found an upper bound of 75% for the SST contribution to Kiritimati coral  $\delta^{18}\text{O}$ , as well as with previous empirically derived estimates (e.g., McGregor et al., 2011).

When placed within the context of centuries worth of coral  $\delta^{18}\text{O}$  data from Kiritimati Island and nearby sites, our results provide key constraints on the physical drivers of the recent intensification of interannual coral  $\delta^{18}\text{O}$  variability in central tropical Pacific corals (Grothe et al., 2020). Given that our results show a  $\sim 30\%$  contribution from seawater  $\delta^{18}\text{O}$  to the coral  $\delta^{18}\text{O}$  anomalies during the three largest El Niño events in recent decades, it is unlikely that the observed  $\sim 25\%$  increase in interannual coral  $\delta^{18}\text{O}$  variability in recent decades relative to the past millennia (Grothe et al., 2020) is caused exclusively by an amplification of the hydrologic response to ENSO-related SST anomalies. Furthermore, instrumental climate data indicate that regional freshening is dynamically linked to warm SST anomalies during El Niño events in this location (Ropelewski & Halpert, 1987), suggesting that the observed increase in coral  $\delta^{18}\text{O}$  variability is driven by an increase in both temperature and hydrological variability.

Our data support the fidelity of Kiritimati coral  $\delta^{18}\text{O}$  records for long-term ENSO reconstruction, even under extreme temperature stress associated with a very strong El Niño event. Taken together, the coral  $\delta^{18}\text{O}$  records, *in situ* SST, and seawater  $\delta^{18}\text{O}$  data analyzed here show that SST dominates the ENSO-related coral  $\delta^{18}\text{O}$  signal ( $\sim 70\%$ ), with a smaller influence from seawater  $\delta^{18}\text{O}$  ( $\sim 30\%$ ). Such quantitative constraints are made possible by *in situ* seawater  $\delta^{18}\text{O}$  observations collected across the event, demonstrating the value of prioritizing *in situ* seawater  $\delta^{18}\text{O}$  observations in the design of regional to global-scale ocean observing systems, including TPOS2020 (Kessler et al., 2019). Calibration studies such as these are necessary to better understand the ENSO signals captured in coral  $\delta^{18}\text{O}$  records and provide insight on signals captured by coral records over longer timescales. Our results suggest that a documented increase in interannual coral  $\delta^{18}\text{O}$  variability at the site from the preindustrial to the present (Grothe et al., 2020) likely reflects an increase in ENSO-related SST.

## Data Availability Statement

All data and metadata presented here are archived at NCDC (<https://www.ncdc.noaa.gov/paleo-search/study/28291>) and have registered International Geo Sample Numbers IECXI000B, IECXI000C, IECXI000D, IECXI000E, IECXI000F, and IECXI000G (i.e., [igsn.org/IECXI000B](https://doi.org/10.7927/H4TJ)). We use additional coral data from Hitt et al., 2021, Grothe et al., 2020, Nurhati et al., 2009, and Evans et al., 1999, and additional *in situ* logger data from Claar et al., 2019. The ERSSTv5 and OISSTv2 datasets can be found at NOAA (<https://psl.noaa.gov/data/gridded/>), and the HadISSTv1.1 data set can be found at <https://www.metoffice.gov.uk/hadobs/hadisst/>.

## References

- Bellenger, H., Guilyardi, E., Leloup, J., Lengaigne, M., & Vialard, J. (2014). ENSO representation in climate models: From CMIP3 to CMIP5. *Climate Dynamics*, 42(7–8), 1999–2018. <https://doi.org/10.1007/s00382-013-1783-z>
- Bonfils, C. J. W., Santer, B. D., Phillips, T. J., Marvel, K., Leung, L. R., Doutriaux, C., & Capotondi, A. (2015). Relative contributions of mean-state shifts and ENSO-driven variability to precipitation changes in a warming climate. *Journal of Climate*, 28(24), 9997–10013. <https://doi.org/10.1175/JCLI-D-15-0341.1>
- Brown, J. R., Brierley, C. M., An, S.-I., Guarino, M., Stevenson, S., Williams, C. J. R., et al. (2020). Comparison of past and future simulations of ENSO in CMIP5/PMIP3 and CMIP6/PMIP4 models. *Climate of the Past*, 16, 1777–1805. <https://doi.org/10.5194/cp-16-1777-2020>

## Acknowledgments

We acknowledge the Republic of Kiribati for granting us permits to collect samples and conduct research on Kiritimati Island (number 005/13), provided by the Environment and Conservation Division. We especially thank Tiito Teabi for his support and expertise in the field on numerous expeditions to Kiritimati from 2012 to 2016. This research was funded by National Science Foundation Awards 1502832, 1658182, 1635068, 1836645, to K. M. Cobb, 1446343 to S. Stevenson, 1446274 to J. K. Baum, and 9802056 (Moana Wave Cruise). Additional funding from a Georgia Tech Presidential Undergraduate Research Award and a National Science Foundation Graduate Research Fellowship was used to support G. K. O'Connor.

- Cahyarini, S. Y., Pfeiffer, M., Timm, O., Dullo, W., & Schonberg, D. G. (2008). Reconstructing seawater  $\delta^{18}\text{O}$  and Sr/Ca ratios: Methods, error analysis and problems, with examples from Tahiti (French Polynesia) and Timor (Indonesia). *Geochimica et Cosmochimica Acta*, 72(12), 2841–2853. <https://doi.org/10.1016/j.gca.2008.04.005>
- Cai, W., Borlace, S., Lengaigne, M., van Rensch, P., Collins, M., Vecchi, G., et al. (2014). Increasing frequency of extreme El Niño events due to greenhouse warming. *Nature Climate Change*, 4, 111–116. <https://doi.org/10.1038/nclimate2100>
- Cai, W., Santoso, A., Wang, G., Yeh, S.-W., An, S.-I., Cobb, K. M., et al. (2015). ENSO and greenhouse warming. *Nature Climate Change*, 5, 849–859. <https://doi.org/10.1038/nclimate2743>
- Carilli, J. E., Hartmann, A. C., Heron, S. F., Pandolfi, J. M., Cobb, K. M., Sayani, H. S., et al. (2017). *Porites* coral response to an oceanographic and human impact gradient in the Line Islands. *Limnology and Oceanography*, 62(6), 2850–2863. <https://doi.org/10.1002/lno.10670>
- Claar, D. C., Cobb, K. M., & Baum, J. K. (2019). In situ and remotely sensed temperature comparison on a Central Pacific atoll. *Coral Reefs*, 38(6), 1343–1349. <https://doi.org/10.1007/s00338-019-01850-4>
- Cobb, K. M. (2002). *Coral records of the El Niño-Southern Oscillation and tropical Pacific climate over the last millennium*. University of California.
- Cobb, K. M., Charles, C. D., Cheng, H., & Edwards, R. L. (2003). El Niño/Southern Oscillation and tropical Pacific climate during the last millennium. *Nature*, 424, 271–276. <https://doi.org/10.1038/nature01779>
- Cobb, K. M., Charles, C. D., & Hunter, D. E. (2001). A central tropical Pacific coral demonstrates Pacific, Indian, and Atlantic decadal climate connections. *Geophysical Research Letters*, 28(11), 2209–2212. <https://doi.org/10.1029/2001GL012919>
- Cobb, K. M., Westphal, N., Sayani, H. R., Watson, J. T., Di Lorenzo, E., Cheng, H., et al. (2013). Highly variable El Niño–Southern Oscillation throughout the Holocene. *Science*, 339(6115), 67–70. <https://doi.org/10.1126/science.1228246>
- Conroy, J. L., Cobb, K. M., Lynch-Stieglitz, J., & Polissar, P. J. (2014). Constraints on the salinity–oxygen isotope relationship in the central tropical Pacific Ocean. *Marine Chemistry*, 161, 26–33. <https://doi.org/10.1016/j.marchem.2014.02.001>
- Dunbar, R. B., Wellington, G. W., Colgan, M. W., & Glynn, P. W. (1994). Eastern Pacific sea surface temperature since 1600 AD: The  $\delta^{18}\text{O}$  record of climate variability in Galapagos corals. *Paleoceanography*, 9(2), 291–315. <https://doi.org/10.1029/93PA03501>
- Epstein, S., Buchsbaum, R., Lowenstam, H., & Urey, H. (1951). Carbonate-water isotopic temperature scale. *Geological Society of America Bulletin*, 62(4), 417–426. [https://doi.org/10.1130/0016-7606\(1951\)62\[417:CITS\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1951)62[417:CITS]2.0.CO;2)
- Evans, M. N., Fairbanks, R. G., & Rubenstone, J. L. (1999). The thermal oceanographic signal of El Niño reconstructed from a Kiritimati Island coral. *Journal of Geophysical Research*, 104(C6), 13409–13421. <https://doi.org/10.1029/1999JC900001>
- Fairbanks, R. G., Evans, M. N., Rubenstone, J. L., Mortlock, R. A., Broad, K., Moore, M. D., & Charles, C. D. (1997). Evaluating climate indices and their geochemical proxies measured in corals. *Coral Reefs*, 16, S93–S100. <https://doi.org/10.1007/s003380050245>
- Freund, M. B., Henley, B. J., Karoly, D. J., McGregor, H. V., Abram, N. J., & Dommenges, D. (2019). Higher frequency of Central Pacific El Niño events in recent decades relative to past centuries. *Nature Geoscience*, 12, 450–455. <https://doi.org/10.1038/s41561-019-0353-3>
- Grothe, P. R., Cobb, K. M., Liguori, G., Di Lorenzo, E., Capotondi, A., Lu, Y., et al. (2020). Enhanced El Niño–Southern Oscillation variability in recent decades. *Geophysical Research Letters*, 47(7), e2019GL083906. <https://doi.org/10.1029/2019GL083906>
- Hetzinger, S., Pfeiffer, M., Dullo, W., Zinke, J., & Garbe-Schonberg, D. (2016). A change in coral extension rates and stable isotopes after El Niño-induced coral bleaching and regional stress events. *Scientific Reports*, 6, 32879. <https://doi.org/10.1038/srep32879>
- Hitt, N. T., Sayani, H. R., Atwood, A. R., Grothe, P. R., Maupin, C., O’Connor, G. K., et al. (2021). Central equatorial Pacific warming and freshening in the 20th century: Insights from a coral ensemble approach. *Geophysical Research Letters*. <https://doi.org/10.1029/2021GL094051>
- Huang, B., L’Heureux, M., Hu, Z., Zhang, H. (2016). Ranking the strongest ENSO events while incorporating SST uncertainty. *Geophysical Research Letters*, 43, 9165–9172. <https://doi.org/10.1002/2016gl070888>
- Huang, B., Thorne, P. W., Banzon, V. F., Boyer, T., Chepurin, G., Lawrimore, J. H., et al. (2017). Extended reconstructed sea surface temperature version 5 (ERSSTv5), upgrades, validations, and intercomparisons. *Journal of Climate*, 30(20), 8179–8205. <https://doi.org/10.1175/JCLI-D-16-0836.1>
- Jimenez, G., Cole, J. E., Thompson, D. M., & Tudhope, A. W. (2018). Northern Galapagos corals reveal twentieth century warming in the eastern tropical Pacific. *Geophysical Research Letters*, 45(4), 1981–1988. <https://doi.org/10.1002/2017GL075323>
- Kessler, W. S., Wijffels, S. E., Cravatte, S., Smith, N., Yosuke, F., Large, W., et al. (2019). Second report of TPOS 2020. GOOS-234.
- Li, J., Xie, S.-P., Cook, E. R., Morales, M. S., Christie, D. A., Johnson, N. C., et al. (2013). El Niño modulations over the past seven centuries. *Nature Climate Change*, 3, 822–826. <https://doi.org/10.1038/nclimate1936>
- Liu, Y., Cobb, K. M., Song, H., Li, Q., Li, C.-Y., Nakatsuka, T., et al. (2017). Recent enhancement of central Pacific El Niño variability relative to last eight centuries. *Nature Communications*, 8, 15386. <https://doi.org/10.1038/ncomms15386>
- McGregor, H. V., Fischer, M. J., Gagan, M. K., Fink, D., & Woodroffe, C. D. (2011). Environmental control of the oxygen isotope composition of *Porites* coral microatolls. *Geochimica et Cosmochimica Acta*, 75, 3930–3944. <https://doi.org/10.1016/j.gca.2011.04.017>
- McGregor, S., Timmermann, A., England, M. H., Elison Timm, O., & Wittenberg, A. T. (2013). Inferred changes in El Niño–Southern Oscillation variance over the past six centuries. *Climate of the Past*, 9(5), 2269–2284. <https://doi.org/10.5194/cp-9-2269-2013>
- Ng, B., Cai, W., Cowan, T., & Bi, D. (2021). Impacts of low-frequency internal climate variability and greenhouse warming on El Niño–Southern Oscillation. *Journal of Climate*, 34(6), 2205–2218. <https://doi.org/10.1175/JCLI-D-20-0232.1>
- Nurhati, I. S., Cobb, K. M., Charles, C. D., & Dunbar, R. B. (2009). Late 20th century warming and freshening in the central tropical Pacific. *Geophysical Research Letters*, 36(21), L21606. <https://doi.org/10.1029/2009GL040270>
- Nurhati, I. S., Cobb, K. M., & Di Lorenzo, E. (2011). Decadal-scale SST and salinity variations in the central tropical Pacific: Signatures of natural and anthropogenic climate change. *Journal of Climate*, 24(13), 3294–3308. <https://doi.org/10.1175/2011JCLI3852.1>
- Power, S., Delage, F., Chung, C., Kociuba, G., & Keay, K. (2013). Robust twenty-first-century projections of El Niño and related precipitation variability. *Nature*, 502, 541–545. <https://doi.org/10.1038/nature12580>
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., et al. (2003). Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *Journal of Geophysical Research*, 108(D14), 4407. <https://doi.org/10.1029/2002JD002670>
- Ren, L., Linsley, B. K., Wellington, G. M., Schrag, D. P., & Hugh-Guldberg, O. (2003). Deconvolving the  $\delta^{18}\text{O}$  seawater component from sub-seasonal coral  $\delta^{18}\text{O}$  and Sr/Ca at Rarotonga in the southwestern subtropical Pacific for the period 1726 to 1997. *Geochimica et Cosmochimica Acta*, 67, 1609–1621. [https://doi.org/10.1016/S0016-7037\(02\)00917-1](https://doi.org/10.1016/S0016-7037(02)00917-1)
- Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., & Wang, W. (2002). An improved in situ and satellite SST analysis for climate. *Journal of Climate*, 15(13), 1609–1625. [https://doi.org/10.1175/1520-0442\(2002\)015<1609:AIISAS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1609:AIISAS>2.0.CO;2)
- Ropelewski, C. F., & Halpert, M. S. (1987). Global and regional scale precipitation patterns associated with the El Niño/Southern Oscillation. *Monthly Weather Review*, 115(7), 1606–1626. [https://doi.org/10.1175/1520-0493\(1987\)115<1606:GARSPP>2.0.CO;2](https://doi.org/10.1175/1520-0493(1987)115<1606:GARSPP>2.0.CO;2)

- Russon, T., Tudhope, A. W., Hegerl, G. C., Collins, M., & Tindall, J. (2013). Inter-annual tropical Pacific climate variability in isotope-enabled CGCM: Implications for interpreting coral stable oxygen isotope records of ENSO. *Climate of the Past*, 9(4), 1542–1557. <https://doi.org/10.5194/cp-9-1543-2013>
- Sayani, H. R., Cobb, K. M., DeLong, K., Hitt, N. T., & Druffel, E. R. M. (2019). Intercolony  $\delta^{18}\text{O}$  and Sr/Ca variability among *Porites* spp. corals at Palmyra Atoll: Toward more robust coral-based estimates of climate. *Geochemistry, Geophysics, Geosystems*, 20(11), 5270–5284. <https://doi.org/10.1029/2019GC008420>
- Stevenson, S., Wittenberg, A. T., Fasullo, J., Coats, S., & Otto-Bliesner, B. (2021). Understanding diverse model projections of future extreme El Niño. *Journal of Climate*, 34(2), 449–464. <https://doi.org/10.1175/JCLI-D-19-0969.1>
- Stevenson, S. L. (2012). Significant changes to ENSO strength and impacts in the twenty-first century: Results from CMIP5. *Geophysical Research Letters*, 39(17), L17703. <https://doi.org/10.1029/2012GL052759>
- Wang, B., Luo, X., Yang, Y., Sun, W., Cane, M. A., Cai, W., et al. (2019). Historical change of El Niño properties sheds light on future changes of extreme El Niño. *Proceeding of the National Academy of Sciences*, 116(45), 22512–22517. <https://doi.org/10.1073/pnas.1911130116>