



symmetry



Article

Non-Standard Finite Difference and Vieta-Lucas Orthogonal Polynomials for the Multi-Space Fractional-Order Coupled Korteweg-de Vries Equation

Khaled M. Saad and Rekha Srivastava

Special Issue

Theory and Applications of Special Functions II

Edited by



Dr. Diego Caratelli



<https://doi.org/10.3390/sym16020242>

Article

Non-Standard Finite Difference and Vieta-Lucas Orthogonal Polynomials for the Multi-Space Fractional-Order Coupled Korteweg-de Vries Equation

Khaled M. Saad ¹  and Rekha Srivastava ^{2,*} 

¹ Department of Mathematics, College of Sciences and Arts, Najran University, Najran 1988, Saudi Arabia; khaledma_sd@hotmail.com or kmalhamam@nu.edu.sa

² Department of Mathematics and Statistics, University of Victoria, Victoria, BC V8W 3R4, Canada

* Correspondence: rekhasrivastava@uvic.ca

Abstract: This paper focuses on examining numerical solutions for fractional-order models within the context of the coupled multi-space Korteweg-de Vries problem (CMSKDV). Different types of kernels, including Liouville-Caputo fractional derivative, as well as Caputo-Fabrizio and Atangana-Baleanu fractional derivatives, are utilized in the examination. For this purpose, the nonstandard finite difference method and spectral collocation method with the properties of the Shifted Vieta-Lucas orthogonal polynomials are employed for converting these models into a system of algebraic equations. The Newton-Raphson technique is then applied to solve these algebraic equations. Since there is no exact solution for non-integer order, we use the absolute two-step error to verify the accuracy of the proposed numerical results.

Keywords: multi-space fractional-order coupled Korteweg-de Vries equation; Shifted Vieta-Lucas orthogonal polynomials; Vieta-Lucas spectral collocation method; operators of fractional calculus



Citation: Saad, K.M.; Srivastava, R. Non-Standard Finite Difference and Vieta-Lucas Orthogonal Polynomials for the Multi-Space Fractional-Order Coupled Korteweg-de Vries Equation. *Symmetry* **2024**, *16*, 242. <https://doi.org/10.3390/sym16020242>

Academic Editors: Diego Caratelli and Junesang Choi

Received: 28 November 2023

Revised: 2 February 2024

Accepted: 6 February 2024

Published: 16 February 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Mathematicians who delve into fractional calculus broaden the principles of differentiation and integration to include orders that are non integer. Fractional calculus allows for differentiation and integration to be performed using non-integer , rather than being confined to whole numbers or integers [1–4]. In the 17th century, pioneers like Isaac Newton and Gottfried Leibniz made early strides in advancing fractional calculus. However, it was not until the 19th century that mathematicians such as Karl Weierstrass and Augustin-Louis Cauchy embarked on the development of a structured theory for fractional calculus [3].

There are alternative approaches to defining fractional derivatives and integrals, which encompass the Riemann-Liouville, Liouville-Caputo, and Grünwald-Letnikov definitions. These definitions entail expanding the classical methods of integration and differentiation to include orders that are not integer [5].

Fractional calculus has extensive use across various industries like engineering, physics, finance, and signal processing. It offers an advanced approach to comprehend phenomena like as anomalous diffusion, fractal time series, and, viscoelasticity, all of which exhibit fractional or fractal characteristics [6–14]. In addition, fractional calculus provides a potent mathematical tool for modeling and examining complex systems that cannot be entirely elucidated by employing the standard methods of integer-order calculus [15].

When studying nonlinear dispersive waves, researchers often turn to the Korteweg-de Vries (KdV) equations. For more information [16].

In 1895, Korteweg and de Vries formulated these equations with the aim of simulating shallow water waves in a canal [17].

A nonlinear partial differential equation known as the coupled Korteweg-de Vries (cKdV) equation describes the evolution of many broad, weakly nonlinear waves in a

dispersed medium. It is a generalization of the well-known Korteweg-de Vries (KdV) equation, which describes the behavior of an isolated single wave in particular. Hirota and Satsuma first introduced the development of the linked KdV equation in a study identified as [18].

The KdV equations elucidate the interaction between two lengthy waves characterized by distinct dispersion relations [18,19].

It is important to highlight that the Korteweg-de Vries equations appear in diverse fields within the physical sciences, ranging from plasmas and fluids to vibrations in low-temperature crystal lattices. Although these applications may appear unrelated, they all originate from a comprehensive physical model. When a particular aspect of the problem is examined closely, they all lead to the Korteweg-de Vries equation. In this sense, the Korteweg-de Vries equation is considered to have broad and universal applicability.

Vieta-Lucas polynomials are widely used in spectral methods used to solve partial and ordinary differential equations [20]. These methods show fast exponential convergence or spectral precision, particularly for problems in simple geometric contexts where the solutions are smooth. These methods differ from finite difference techniques in that the determination of the approximation coefficient entirely defines the solution at any specified point within the interval. Therefore, utilizing operational matrix spectral methods that rely on Vieta-Lucas polynomials for solving the system outlined in Equation (1) below is of considerable significance [21–23]. For the application of spectral collocation method and finite difference method, some powerful mathematical techniques have been used in recent years by researchers [24–28].

The objective of this study is to offer numerical solutions for the multi-space fractional-order coupled Korteweg-de Vries problem. Various kernels, Liouville-Caputo fractional derivatives, Caputo-Fabrizio, and Atangana-Baleanu fractional derivatives are employed in this work. The solution method integrates the Vieta-Lucas collocation method with the nonstandard finite difference method to discretize the spatial fractional-order diffusion equation. This results in a set of ordinary differential equations, simplifying the conditions significantly. The nonstandard finite difference method (FDM) is then applied to solve this set of ordinary differential equations. For more detailed information, refer to the provided references [29–31].

The paper is organized as follows: In Section 2, we delve into function approximation and the application of Vieta-Lucas polynomials. Section 3 demonstrates the use of the spectral method to solve the multi-space fractional-order coupled Korteweg-de Vries problem, employing Liouville-Caputo, Caputo-Fabrizio, and Atangana-Baleanu fractional derivatives. Section 4 presents an explanation and numerical results for the multi-space fractional-order coupled Korteweg-de Vries, incorporating the three kernels utilized in this study. Finally, concluding remarks are provided in Section 5.

2. Vieta-Lucas Polynomials and Function Approximations

In this section, we furnish the definitions of the shifted Vieta-Lucas polynomials (VLPs), along with their corresponding notations and key properties. This serves as a foundational understanding for their application in the subsequent analyses [32]. Our research has primarily focused on a specific category of orthogonal polynomials known as the Vieta-Lucas polynomials. This family of orthogonal polynomials can be generated through the utilization of recurrence relations and analytical equations associated with these polynomials. Also we will introduce the convergence analysis and the nonstandard finite difference scheme notations.

2.1. Shifted Vieta-Lucas Polynomials

The Vieta-Lucas polynomials $\Psi_j(z)$, referred to as Lucas polynomials (VLPs), are discussed along with their respective notations and properties as outlined by Vieta. Specif-

ically, for polynomials of degree $j \in \mathbb{N}_0$, $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\}$, the definition is provided as follows [32]:

$$\Psi_j(t) = 2 \cos(j \psi), \quad \psi = \arccos(0.5z), \quad \psi \in [0, \pi], \quad -2 \leq z \leq 2.$$

The following recurrence relation for $\Psi_{j(z)}$ is easily demonstrated:

$$\Psi_j(z) = z \Psi_{j-1}(z) - \Psi_{j-2}(z), \quad j = 2, 3, \dots, \quad \Psi_0(z) = 2, \quad \Psi_1(z) = z.$$

A set of orthogonal polynomials defined on the interval $[0, 1]$ is established by incorporating VLPs and $z = 4\alpha - 2$. These polynomials will be represented by the symbol $\Phi_m(\beta)$, as illustrated in: $\Phi_j(\alpha) = \Psi(4\alpha - 2) = \Psi(2\sqrt{\alpha})$.

$\Phi_j(\alpha)$ have the following recurrence relation:

$$\Phi_{j+1}(\alpha) = (4\alpha - 2)\Phi_{j-1}(\alpha) - \Phi_{j-2}(\alpha), \quad j = 2, 3, \dots,$$

where, $\Phi_0(\alpha) = 2$, $\Phi_1(\alpha) = 4\alpha - 2$. Also, we find $\Phi_j(0) = 2(-1)^j$ and $\Phi_j(1) = 2$, $j = 0, 1, 2, \dots$.

The analytical formula for $\Phi_j(\alpha)$ is:

$$\Phi_k(\alpha) = 2j \sum_{j=0}^k (-1)^j \frac{4^{k-j} \Gamma(2k-j)}{\Gamma(j+1) \Gamma(2k-2j+1)} \alpha^{k-j}, \quad k = 2, 3, \dots$$

The polynomials $\Phi_k(\alpha)$ are orthogonal polynomials on $[0, 1]$ w.r.t. $\frac{1}{\sqrt{\alpha-\alpha^2}}$, and so we have:

$$\langle \Phi_k(\alpha), \Phi_j(\alpha) \rangle = \int_0^1 \frac{\Phi_k(\alpha) \Phi_j(\alpha)}{\sqrt{\alpha-\alpha^2}} d\alpha = \begin{cases} 0, & k \neq j \neq 0, \\ 4\pi, & k = j = 0, \\ 2\pi, & k = j \neq 0. \end{cases}$$

Let $\Omega(\alpha) \in L^2[0, 1]$, then using $\Phi_k(\alpha)$, we have:

$$\Omega(\alpha) = \sum_{k=0}^{\infty} c_k \Phi_k(\alpha). \quad (1)$$

To convert $\Omega(\alpha)$ into terms of $\Phi_m(\alpha)$, it is necessary to evaluate c_k . This can be expressed by considering only the first $m+1$ terms of (1), we obtain that

$$\Omega_m(\alpha) = \sum_{k=0}^m c_k \Phi_k(\alpha), \quad (2)$$

The values of c_k for $k = 0, 1, 2, \dots, m$ can be computed from:

$$c_0 = \frac{1}{4\pi} \int_0^1 \frac{\phi(\alpha) \Phi_0(\alpha)}{\sqrt{\alpha-\alpha^2}} d\alpha, \quad c_k = \frac{1}{2\pi} \int_0^1 \frac{\phi(\alpha) \Phi_k(\alpha)}{\sqrt{\alpha-\alpha^2}} d\alpha, \quad k = 1, 2, 3, \dots \quad (3)$$

2.2. Convergence Analysis

Lemma 1. Assuming that the function $\Omega_m(\alpha)$ belongs to $L^2[0, 1]$ with the weight function $\frac{1}{\sqrt{\alpha-\alpha^2}}$ and $|\Omega''(\alpha)| \leq \varepsilon$, for some constant ε , the series given by (2) uniformly converges to $\Omega(\alpha)$ as $m \rightarrow \infty$. Furthermore, the estimates listed below are accomplished:

1. Equation (2) has a bounded series of coefficients, meaning

$$|c_k| \leq \frac{\varepsilon}{4k(k^2-1)}, \quad k > 2.$$

2. The norm for the error estimate ($L^2[0, 1]$ -norm) can be articulated as follows:

$$\|\Omega(\alpha) - \Omega_m(\alpha)\| < \frac{L}{12\sqrt{m^3}}.$$

3. If $\Omega^{(m)}(\alpha) \in C[0,1]$, then the absolute error bound is satisfied

$$\|\Omega(\alpha) - \Omega_m(\alpha)\| \leq \frac{\Delta \Pi^{m+1}}{(m+1)!} \sqrt{\pi}, \quad \Delta = \max_{\alpha \in [0,1]} \Omega^{(m+1)}(\alpha), \quad \text{and } \Pi = \max\{1 - \alpha_0, \alpha_0\}.$$

For a more in-depth exploration of the convergence analysis and the use of polynomials to approximate (2), readers may refer [33].

2.3. Nonstandard Finite Difference Scheme Notations

We provide the following elucidation of the discrete first derivative:

$$\frac{d\Omega}{d\beta} \rightarrow \frac{\Omega_{s+1} - \psi_1(h)\Omega_s}{\psi_2(h)}, \quad (4)$$

where $\psi_1(h)$ and $\psi_2(h)$ are functions in the step-size discretization $h = \Delta(\beta)$ and

$$\psi_1(h) = 1 + O(h^2) \quad \text{and} \quad \psi_2(h) = h + O(h^2).$$

The term used to denote the first derivative in the nonstandard finite difference technique is referred to as the presentation formula. Moreover, it is required that the denominator function adheres to the condition that $0 < \psi_1(h) < 1$, $h \rightarrow 0$. While the exact base function $\psi_2(h)$ is not specified, commonly used functions in the non-standard finite difference method can be introduced as follows: $\psi_2(h) = \exp(h) - 1$, $\psi_2(h) = h$, $\psi_2(h) = \sinh(h)$, $\psi_2(h) = \frac{1 - \exp(-\lambda h)}{\lambda}$, etc. (see [34]).

3. Spectral Method for Solving CMSKDV

This section is subdivided into three sections, each delineating the fractional derivatives associated with distinct kernels. Recent works have investigated a fractional derivative based on the exponential decay rule, which can be described as a generalized power law function. A more generalized version of the exponential function called the Mittag-Leffler function served as the foundation for the fractional derivative with a non-local kernel that Dumitru Baleanu and Abdou Atangana presented. The depiction of intricate physical issues that follow the rules of power and exponential decay is made possible by this derivative [35,36].

The coupled KdV equations in the classical case are given as follows:

$$\phi_\beta - c \phi_{\alpha\alpha\alpha} - \mu_2 \phi \phi_\alpha - \mu_1 \phi \varphi_\alpha = 0, \quad (5)$$

and

$$\varphi_\beta - c \varphi_{\alpha\alpha\alpha} + \mu_1 \phi \varphi_\alpha = 0, \quad (6)$$

where c , μ_1 and μ_2 are constant parameters.

3.1. CMSKDV via the Liouville-Caputo Fractional Derivative

When replacing the classical derivative of the coupled KdV equations with a fractional derivative employing a power-law kernel or Liouville-Caputo fractional derivative (LC), the resulting system of equations for the coupled KdV reads as follows:

$$\phi_\beta - c {}^{\text{LC}}D_\alpha^{\nu_1} \phi - \mu_2 \phi {}^{\text{LC}}D_\alpha^{\nu_2} \phi - \mu_1 \varphi {}^{\text{LC}}D_\alpha^{\nu_2} \varphi = 0 \quad (7)$$

and

$$\varphi_\beta - c {}^{\text{LC}}D_\alpha^{\nu_1} \varphi + \mu_1 \varphi {}^{\text{LC}}D_\alpha^{\nu_2} \varphi = 0, \quad (8)$$

$$2 < \nu_1 \leq 3, \quad 0 < \nu_2 \leq 1,$$

where ${}^{\text{LC}}D_\alpha^{\nu_1}$ and ${}^{\text{LC}}D_\alpha^{\nu_2}$ are the LC fractional derivatives of order ν_1 and ν_2 , respectively for the functions $\phi(\alpha)$ and $\varphi(\alpha)$ that belong to $H^1(0, b)$ defined in the following form [37,38].

$${}^{\text{LC}}D_\alpha^{\nu_1} \varphi(\alpha) = \frac{1}{\Gamma(3 - \nu_1)} \int_0^\alpha \frac{\varphi^{(3)}(\tau)}{(\alpha - \tau)^{\nu_1 - 2}} d\tau \quad (\alpha > 0), \quad (9)$$

$${}^{\text{LC}}D_\alpha^{\nu_2} \varphi(\alpha) = \frac{1}{\Gamma(1 - \nu_2)} \int_0^\alpha \frac{\varphi^{(1)}(\tau)}{(\alpha - \tau)^{\nu_2}} d\tau \quad (\alpha > 0). \quad (10)$$

The following theorem provides an approximate formula for the fractional derivative ${}^{\text{LC}}D_\alpha^\nu \Omega(\alpha)$, $n - 1 < \nu \leq n$, with the power-law kernel.

Theorem 1 (see [39]). *The LC-fractional-order derivative of the function $\Omega_m(\alpha)$, as defined in Equation (2), can be determined using the following approximate formula:*

$${}^{\text{LC}}D_\alpha^\nu (\Omega_m(\alpha)) = \sum_{k=\lceil \nu \rceil}^m \sum_{\ell=0}^{k-\mu} \alpha_k \vartheta_{k,\ell} \Pi_{\nu,\ell}^{\text{LC}}(\alpha), \quad (11)$$

where

$$\vartheta_{k,\ell} = \frac{(-1)^\ell 4^{k-\ell} 2k\Gamma(2k - \ell)}{\Gamma(\ell + 1)\Gamma(2k - 2\ell + 1)}, \quad (12)$$

and

$$\Pi_{\nu,\ell}^{\text{LC}}(\alpha) = \frac{\Gamma(k - \ell + 1)}{\Gamma(k - \ell - \nu + 1)} \alpha^{k-\ell-\nu}. \quad (13)$$

Subsequently, we move forward with solving Equations (7) to (8) using the Vieta-Lucas spectral collocation technique, which is explained as follows:

$$\phi_m(\alpha, \beta) = \sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha) \quad (14)$$

and

$$\varphi_m(\alpha, \beta) = \sum_{k=0}^m \varphi_k(\beta) \Phi_k(\alpha). \quad (15)$$

The two Equations (7) through (8) can be represented as follows in accordance with Theorem 1 and the expansions (14)–(15)

$$\begin{aligned} \sum_{k=0}^m \frac{d\phi_k(\beta)}{d\beta} \Phi_k(\alpha) &= c \sum_{k=\lceil v_1 \rceil}^m \sum_{\ell=\lceil v_1 \rceil}^{k-\lceil v_1 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{v,\ell}^{\text{LC}}(\alpha) + \mu_2 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{\theta,\ell}^{\text{LC}}(\alpha) \right) + \mu_1 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil v_1 \rceil}^m \sum_{\ell=\lceil v_1 \rceil}^{k-\lceil v_1 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{\theta,\ell}^{\text{LC}}(\alpha) \right) \end{aligned} \quad (16)$$

and

$$\begin{aligned} \sum_{k=0}^m \frac{d\varphi_k(\beta)}{d\beta} \Phi_k(\alpha) &= c \sum_{k=\lceil v_1 \rceil}^m \sum_{\ell=\lceil v_1 \rceil}^{k-\lceil v_1 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{v,\ell}^{\text{LC}}(\alpha) - \mu_1 \left(\sum_{k=0}^m \varphi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{\theta,\ell}^{\text{LC}}(\alpha) \right). \end{aligned} \quad (17)$$

For the Equations (16) and (17), we will evaluate them at m points α_p ($p = 0, 1, \dots, m-1$) as outlined below:

$$\begin{aligned} \sum_{k=0}^m \frac{d\phi_k(\beta)}{d\beta} \Phi_k(\alpha_p) &= c \sum_{k=\lceil v_1 \rceil}^m \sum_{\ell=\lceil v_1 \rceil}^{k-\lceil v_1 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{v,\ell}^{\text{LC}}(\alpha_p) + \mu_2 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha_p) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{\theta,\ell}^{\text{LC}}(\alpha_p) \right) + \mu_1 \left(\sum_{k=0}^m \varphi_k(\beta) \Phi_k(\alpha_p) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{\theta,\ell}^{\text{LC}}(\alpha_p) \right), \end{aligned} \quad (18)$$

$$\begin{aligned} \sum_{k=0}^m \frac{d\varphi_k(\beta)}{d\beta} \Phi_k(\alpha_p) &= c \sum_{k=\lceil v_1 \rceil}^m \sum_{\ell=\lceil v_1 \rceil}^{k-\lceil v_1 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{v,\ell}^{\text{LC}}(\alpha_p) - \mu_1 \left(\sum_{k=0}^m \varphi_k(\beta) \Phi_k(\alpha_p) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{\theta,\ell}^{\text{LC}}(\alpha_p) \right). \end{aligned} \quad (19)$$

Moreover, through substitution from Equation (2), we can articulate the corresponding boundary conditions, leading to the ensuing set of equations:

$$\sum_{k=0}^m (-1)^k \phi_k(\beta) = g_1(\beta), \quad \sum_{k=0}^m (-1)^k \varphi_k(\beta) = h_1(\beta), \quad (20)$$

$$\sum_{k=0}^m \phi_k(\beta) = g_2(\beta), \quad \sum_{k=0}^m \varphi_k(\beta) = h_2(\beta) \quad (21)$$

and

$$\sum_{k=0}^m 2k^2 \phi_k(\beta) = g_3(\beta) \quad \text{and} \quad \sum_{k=0}^m 2k^2 \varphi_k(\beta) = h_3(\beta). \quad (22)$$

Ultimately, applying the nonstandard finite difference approach as described in [39] leads us to a set of nonlinear algebraic equations, we obtain

$$\begin{aligned} \sum_{k=0}^m \frac{\phi_k^p - \phi_k^{p-1}}{0.5(\exp(2h) - 1)} \Phi_k(\alpha_p) &= c \sum_{k=\lceil \nu_1 \rceil}^m \sum_{\ell=\lceil \nu_1 \rceil}^{k-\lceil \nu_1 \rceil} \phi_k^p \vartheta_{k,\ell} \Pi_{\theta,\ell}^{\text{LC}}(\alpha_p) + \mu_2 \left(\sum_{k=0}^m \phi_k^p \Phi_k(\alpha_p) \right) \\ &\cdot \left(\sum_{k=\lceil \nu_2 \rceil}^m \sum_{\ell=\lceil \nu_2 \rceil}^{k-\lceil \nu_2 \rceil} \phi_k^p \vartheta_{k,\ell} \Pi_{\theta,\ell}^{\text{LC}}(\alpha_p) \right) + \mu_1 \left(\sum_{k=0}^m \phi_k^p \Phi_k(\alpha_p) \right) \\ &\cdot \left(\sum_{k=\lceil \nu_2 \rceil}^m \sum_{\ell=\lceil \nu_2 \rceil}^{k-\lceil \nu_2 \rceil} \phi_k^p \vartheta_{k,\ell} \Pi_{\theta,\ell}^{\text{LC}}(\alpha_p) \right) = 0, \end{aligned} \tag{23}$$

$$\begin{aligned} \sum_{k=0}^m \frac{\phi_k^p - \phi_k^{p-1}}{0.5(\exp(2h) - 1)} \Phi_k(\alpha_p) &= c \sum_{k=\lceil \nu_1 \rceil}^m \sum_{\ell=\lceil \nu_1 \rceil}^{k-\lceil \nu_1 \rceil} \phi_k^p \vartheta_{k,\ell} \Pi_{\nu,\ell}^{\text{LC}}(\alpha_p) - \mu_1 \left(\sum_{k=0}^m \phi_k^p \Phi_k(\alpha_p) \right) \\ &\cdot \left(\sum_{k=\lceil \nu_2 \rceil}^m \sum_{\ell=\lceil \nu_2 \rceil}^{k-\lceil \nu_2 \rceil} \phi_k^p \vartheta_{k,\ell} \Pi_{\theta,\ell}^{\text{LC}}(\alpha_p) \right). \end{aligned} \tag{24}$$

These can be solved using well-known methods such as the Newton-Raphson technique, enabling us to ascertain the values of ϕ_k^p and φ_k^p for $k = 0, 1, \dots, m$.

3.2. CMSKDV via the Caputo-Fabrizio Fractional Derivative

By replacing the classical derivative with the fractional derivative incorporating either the exponential-decay kernel or the Caputo-Fabrizio (CF), the coupled (KdV) equations can be represented in the following manner:

$$\phi_\beta - c \text{CF}D_\alpha^{\nu_1} \phi - \mu_2 \phi \text{CF}D_\alpha^{\nu_2} \phi - \mu_1 \phi \text{CF}D_\alpha^{\nu_2} \varphi = 0 \tag{25}$$

and

$$\varphi_\beta - c \text{CF}D_\alpha^{\nu_1} \varphi + \mu_1 \phi \text{CF}D_\alpha^{\nu_2} \varphi = 0. \tag{26}$$

The fractional derivative operator $\text{CF}D_\alpha^\nu$ (ν might be ν_1 or ν_2) of order $n < \nu < n + 1$, which incorporates the exponential-decay kernel, is defined as follows [40]:

$$\text{CF}D_{a+}^\nu \varphi(\alpha) = \text{CF}D_{a+}^\kappa (D^n \varphi(\alpha)) = \frac{M(\kappa)}{1 - \kappa} \int_a^\alpha \varphi^{(n+1)}(\tau) e^{-\frac{\kappa(\alpha-\tau)}{1-\kappa}} d\tau, \tag{27}$$

where $n = \lfloor \nu \rfloor$ is the floor of ν (that is, the integer part) and $\kappa = \lceil \nu \rceil$ is the ceiling of ν (that is, the decimal part) and $M(\kappa)$ is the normalization function.

Theorem 2 (see [30,40]). *The left-hand side features a fractional derivative employing an exponential-decay kernel with an order $\nu \in (n, n + 1)$*

$$\varphi(\alpha) = \alpha^p \quad (p \geq \lceil \nu \rceil)$$

can be represented in the following manner:

$$\text{CF}D_{0+}^\nu \alpha^p = \frac{M(\kappa)\Gamma(p + 1)}{1 - \kappa} \left(\sum_{i=0}^{p-n-1} \frac{(-1)^i \alpha^{p-n-1-i}}{\Gamma(p-n-i)\left(\frac{\kappa}{1-\kappa}\right)^{i+1}} + \frac{(-1)^{p-n}}{\left(\frac{\kappa}{1-\kappa}\right)^{p-n}} e^{-\frac{\kappa\alpha}{1-\kappa}} \right). \tag{28}$$

The formula in Theorem 1 of [40] when $a = 0$ corresponds to a specific case of the Formula (28). For additional details regarding the definitions and characteristics of fractional CF-derivatives, refer to [40].

The following theorem provides the approximation formula of ${}^{\text{CF}}D_{0+}^{\nu} \Omega_m(x)$ by means of (1).

Theorem 3 (see [30]). *The ${}^{\text{CF}}D_{\alpha}^{\nu}(\Omega_m(\alpha))$ can be written as follows:*

$${}^{\text{CF}}D_{0+}^{\nu}(\Omega_m(\alpha)) = \sum_{k=\lceil \nu \rceil}^m \sum_{\ell=\lceil \nu \rceil}^k a_k \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha), \quad (29)$$

where

$$\Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha) = \left(\frac{M(\kappa)}{1-\kappa} \right) \left(\frac{(-1)^{\ell-n}}{\left(\frac{\kappa}{1-\kappa}\right)^{\ell-n}} e^{-\frac{\kappa\alpha}{1-\kappa}} + \sum_{p=0}^{\ell-n-1} \frac{(-1)^p \alpha^{\ell-n-1-p}}{\Gamma(\ell-n-p) \left(\frac{\kappa}{1-\kappa}\right)^{p+1}} \right). \quad (30)$$

By using Equations (25) and (26), (29) and the Formula (2) with CF, we obtain

$$\begin{aligned} \sum_{k=0}^m \frac{d\phi_k(\beta)}{d\beta} \Phi_k(\alpha) &= c \sum_{k=\lceil \nu_1 \rceil}^m \sum_{\ell=\lceil \nu_1 \rceil}^{k-\lceil \nu_1 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha) + \mu_2 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil \nu_2 \rceil}^m \sum_{\ell=\lceil \nu_2 \rceil}^{k-\lceil \nu_2 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha) \right) + \mu_1 \left(\sum_{k=0}^m \varphi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil \nu_2 \rceil}^m \sum_{\ell=\lceil \nu_2 \rceil}^{k-\lceil \nu_2 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha) \alpha^{k-\ell-\theta} \right), \end{aligned} \quad (31)$$

$$\begin{aligned} \sum_{k=0}^m \frac{d\varphi_k(\beta)}{d\beta} \Phi_j(\alpha) &= c \sum_{k=\lceil \nu_1 \rceil}^m \sum_{\ell=\lceil \nu_1 \rceil}^{k-\lceil \nu_1 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha) - \mu_1 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil \nu_2 \rceil}^m \sum_{\ell=\lceil \nu_2 \rceil}^{k-\lceil \nu_2 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha) \right). \end{aligned} \quad (32)$$

To obtain the ultimate expressions denoted as (31) and (32), we will assess them at m specific points α_p ; ($p = 0, 1, \dots, m-1$) according to the following procedure:

$$\begin{aligned} \sum_{k=0}^m \frac{\phi_k^p - \phi_k^{p-1}}{0.5(\exp(2h) - 1)} \Phi_k(\alpha_p) &= c \sum_{k=\lceil \nu_1 \rceil}^m \sum_{\ell=\lceil \nu_1 \rceil}^{k-\lceil \nu_1 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha_p) + \mu_2 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil \nu_2 \rceil}^m \sum_{\ell=\lceil \nu_2 \rceil}^{k-\lceil \nu_2 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha_p) \right) + \mu_1 \left(\sum_{k=0}^m \varphi_k(\beta) \Phi_k(\alpha_p) \right) \\ &\cdot \left(\sum_{k=\lceil \nu_2 \rceil}^m \sum_{\ell=\lceil \nu_2 \rceil}^{k-\lceil \nu_2 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha_p) \right), \end{aligned} \quad (33)$$

and

$$\begin{aligned} \sum_{k=0}^m \frac{\varphi_k^p - \varphi_k^{p-1}}{0.5(\exp(2h) - 1)} \Phi_j(\alpha_p) &= c \sum_{k=\lceil \nu_1 \rceil}^m \sum_{\ell=\lceil \nu_1 \rceil}^{k-\lceil \nu_1 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha_p) - \mu_1 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha_p) \right) \\ &\cdot \left(\sum_{k=\lceil \nu_2 \rceil}^m \sum_{\ell=\lceil \nu_2 \rceil}^{k-\lceil \nu_2 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{CF}}(\alpha_p) \right). \end{aligned} \quad (34)$$

In Equations (33) to (34), the nonstandard finite difference approach as described in [39] was utilized to compute the derivative concerning with β . Consequently, the equations were transformed into a set of algebraic equations.

One can solve (33) to (34) using established methods like the Newton-Raphson technique, allowing us to determine the values of ϕ_k^p and ϕ_k^q for $k = 0, 1, \dots, m$.

3.3. CMSKDV via the Atangana-Baleanu-Caputo Fractional Derivative

When we replace the classical derivative with the fractional derivative incorporating either the Mittag-Leffler kernel or the Atangana-Baleanu-Caputo fractional-order derivative (ABC), the resultant coupled Korteweg-de Vries system of equations is represented as follows:

$$\phi_\beta - c \text{ABC}D_\alpha^{\nu_1} \phi - \mu_2 \phi \text{ABC}D_\alpha^{\nu_2} \phi - \mu_1 \phi \text{ABC}D_\alpha^{\nu_2} \phi = 0 \quad (35)$$

and

$$\phi_\beta - c \text{ABC}D_\alpha^{\nu_1} \phi + \mu_1 \phi \text{ABC}D_\alpha^{\nu_2} \phi = 0. \quad (36)$$

The fractional derivative $\text{ABC}_a D^\nu$ (ν can be ν_1 or ν_2 .) of order $n < \nu \leq n + 1$ is defined by [36,41].

$$\begin{aligned} \text{ABC}_0 D^\nu \varphi(\alpha) &= \text{ABC}_0 D^\kappa (D^n \varphi(\alpha)) \\ &= \frac{M(\kappa)}{1 - \kappa} \int_0^\alpha \varphi^{(n+1)}(\tau) E_\kappa \left(-\frac{\kappa}{1 - \kappa} (\alpha - \tau)^\kappa \right) d\tau, \end{aligned}$$

where $n = \lfloor \nu \rfloor$ = the floor of ν (that is, the integer part) and $\kappa = \lceil \nu \rceil$ = the ceiling of ν (that is, the decimal part).

$M(\kappa)$ is the normalization function such that $M(0) = M(1) = 1$ and

$$E_\nu(\alpha) = \sum_{k=0}^{\infty} \frac{\alpha^k}{\Gamma(k\nu + 1)}$$

is the Mittag-Leffler function [42] Section 4.

Theorem 4 (see [31]). *The ABC-derivative of order $\nu \in (n, n + 1]$ of the following function:*

$$\varphi(\alpha) = \alpha^p \quad (p \geq \lceil \nu \rceil)$$

is given by

$$\text{ABC}_0 D^\nu \alpha^p = \frac{M(\kappa)\Gamma(p+1)}{1 - \kappa} \sum_{i=0}^{\infty} \left(-\frac{\kappa}{1 - \kappa} \right)^i \frac{\alpha^{p+i\kappa-n}}{\Gamma(p+i\kappa-n+1)}. \quad (37)$$

In the following theorem, we present the fundamental approximation formula for $\text{ABC}_0 D^\nu \Omega_m(\alpha)$ by utilizing the approximation (2).

Theorem 5 (see [31]). *The ABC-derivative $\text{ABC}_0 D^\nu (\Omega_m(\alpha))$ can be expressed as follows:*

$$\text{ABC}_0 D^\nu (\Omega_m(\alpha)) = \sum_{k=\lceil \nu \rceil}^m \sum_{\ell=\lceil \nu \rceil}^{k-\lceil \nu \rceil} \alpha_k \vartheta_{k,\ell} \Pi_{k,\ell,\kappa} \text{ABC}(\alpha), \quad (38)$$

where

$$\Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha) = \frac{M(\kappa)\Gamma(\ell+1)}{1-\kappa} \sum_{s=0}^{\infty} \frac{\left(-\frac{\kappa}{1-\kappa}\right)^s \alpha^{\ell+p\kappa-n}}{\Gamma(\ell+s\kappa-n+1)}. \quad (39)$$

It can be readily noticed that

$$\begin{aligned} \sum_{k=0}^m \frac{d\phi_k(\beta)}{d\beta} \Phi_k(\alpha) &= c \sum_{k=\lceil v_1 \rceil}^m \sum_{\ell=\lceil v_1 \rceil}^{k-\lceil v_1 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha) + \mu_2 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha) \right) + \mu_1 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha) \alpha^{k-\ell-\theta} \right), \end{aligned} \quad (40)$$

$$\begin{aligned} \sum_{k=0}^m \frac{d\varphi_k(\beta)}{d\beta} \Phi_k(\alpha) &= c \sum_{k=\lceil v_1 \rceil}^m \sum_{\ell=\lceil v_1 \rceil}^{k-\lceil v_1 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha) - \mu_1 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha) \right). \end{aligned} \quad (41)$$

To derive the final formulations indicated as (40) and (41), we will evaluate them at m distinct points α_p ; ($p = 0, 1, \dots, m-1$) following the outlined procedure:

$$\begin{aligned} \sum_{k=0}^m \frac{\phi_k^p - \phi_k^{p-1}}{0.5(\exp(2h) - 1)} \Phi_k(\alpha_p) &= c \sum_{k=\lceil v_1 \rceil}^m \sum_{\ell=\lceil v_1 \rceil}^{k-\lceil v_1 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha_p) + \mu_2 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \phi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha_p) \right) + \mu_1 \left(\sum_{k=0}^m \varphi_k(\beta) \Phi_k(\alpha_p) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha_p) \right), \end{aligned} \quad (42)$$

and

$$\begin{aligned} \sum_{k=0}^m \frac{\phi_k^p - \phi_k^{p-1}}{0.5(\exp(2h) - 1)} \Phi_k(\alpha_p) &= c \sum_{k=\lceil v_1 \rceil}^m \sum_{\ell=\lceil v_1 \rceil}^{k-\lceil v_1 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha_p) - \mu_1 \left(\sum_{k=0}^m \phi_k(\beta) \Phi_k(\alpha_p) \right) \\ &\cdot \left(\sum_{k=\lceil v_2 \rceil}^m \sum_{\ell=\lceil v_2 \rceil}^{k-\lceil v_2 \rceil} \varphi_k(\beta) \vartheta_{k,\ell} \Pi_{k,\ell,\kappa}^{\text{ABC}}(\alpha_p) \right). \end{aligned} \quad (43)$$

These equations can be resolved utilizing well-known approaches such as the Newton-Raphson technique, enabling us to calculate the values of ϕ_k^p and φ_k^p for $k = 0, 1, \dots, m$.

4. Numerical Results and Discussion

In this section, we utilize the three previously mentioned approaches to perform a numerical evaluation of the multi-space fractional-order coupled Korteweg-de Vries equation, incorporating three different kernels. Figures 1–9 are included to visually represent the numerical results. The numerical results provided in this section will be computed using the designated values: $c = -1$, $\mu_1 = -6$, $\mu_2 = 7$, and $a = 1$.

In Figure 1, a comparison is made between the exact solution and the approximate solution of (7) and (8) in the case of the LC-derivative, as specified above with $\nu_1 = 2.8$, and $\nu_2 = 0.8$.

The exact solutions of the coupled Korteweg-de Vries equation under classical conditions are given as follows:

$$\phi(\alpha, \beta) = \varphi(\alpha, \beta) = \frac{4c^2 \exp(c(\alpha - \beta c^2))}{(\exp(c(\alpha - \beta c^2)) + 1)^2}. \quad (44)$$

In order to apply the initial condition in the exact solution for the coupled Korteweg-de Vries equation, we set $\beta = 0$. Subsequently, we acquire the initial values as follows:

$$\phi(\alpha) = \varphi(\alpha) = \frac{4c^2 \exp(c(\alpha))}{(\exp(c(\alpha)) + 1)^2} \quad (45)$$

Additionally, the boundary conditions, once $\alpha = 0$ and $\alpha = 1$ are set, are expressed as follows:

$$\phi(0, \beta) = \varphi(0, \beta) = \frac{4c^2 \exp(c(-\beta c^2))}{(\exp(c(-\beta c^2)) + 1)^2}, \quad (46)$$

$$\phi(1, \beta) = \varphi(1, \beta) = \frac{4c^2 \exp(c(1 - \beta c^2))}{(\exp(c(1 - \beta c^2)) + 1)^2}, \quad (47)$$

$$\phi_\alpha(0, \beta) = \varphi_\alpha(0, \beta) = \frac{4c^3 e^{c(\beta(-c^2))}}{(e^{c(\beta(-c^2))} + 1)^2} - \frac{8c^3 e^{2c(\beta(-c^2))}}{(e^{c(\beta(-c^2))} + 1)^3}. \quad (48)$$

In all calculations conducted in this study, the initial and boundary conditions mentioned earlier will be utilized.

In Figure 2, we display the absolute discrepancy between the precise solution (44) and the numerical solution (7) and (8) for the identical set of parameters as depicted in Figure 1.

Given the absence of an exact solution for the fractional-order equation in this scenario, we will assess the absolute error between two consecutive steps.

Figure 3 illustrates the plot of the discrepancy between two consecutive stages.

Based on the data presented in the preceding three figures, it is evident that the numerical solutions we have presented exhibit high accuracy with a remarkably low error rate. Our emphasis on fractional-order equations stems from their effectiveness in gauging the error between two consecutive phases, aligning with the primary objective of our study.

For the LC-derivative, we have carried out the necessary calculations and generated the corresponding data representations.

We conducted a comparable investigation to that of LC, as depicted in Figures 4–6, but this time incorporating the CF-derivative operator.

We have also examined the numerical solutions of Equations (42)–(43) in Figures 7–9, this time employing the ABC-derivative operator.

Based on this analysis, we have noted a relatively minimal numerical calculation error. As a result, this study's findings can be extended and applied to other fractional-order equations and systems.




Figure 1. In the case of the LC-derivative, a graph comparing the exact solution to the approximation of (7) in (a) and (8) in (b) for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6,$ and $a = 1.$




Figure 2. In the case of the LC-derivative, a graph of the absolute error of (7) in (a) and (8) in (b) for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6,$ and $a = 1.$




Figure 3. In the case of the LC-derivative, a graph of the absolute error of two step of (7) in (a) and (8) in (b) for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6,$ and $a = 1.$




Figure 4. In the case of the CF-derivative, a graph comparing the exact solution to the approximation of (25) in (a) and (26) in (b) for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6,$ and $a = 1.$




Figure 5. In the case of the CF-derivative, a graph, of the absolute error of (25) in (a) and (26) in (b) for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6,$ and $a = 1.$




Figure 6. In the case of the CF-derivative, a graph of the absolute error of two step of (25) in (a) and (26) in (b) for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6,$ and $a = 1.$




Figure 7. In the case of the ABC-derivative, a graph comparing the exact solution to the approximation of (40) in (a) and (41) in (b) for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6,$ and $a = 1.$




Figure 8. In the case of the ABC-derivative, a graph of the absolute error of (40) in (a) and (41) in (b) for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6,$ and $a = 1.$




Figure 9. In the case of the ABC-derivative, a graph of the absolute error of two step of (40) in (a) and (41) in (b) for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6$, and $a = 1$.

In Tables 1–3, the tables illustrate the absolute error between the approximate solutions of two equations under the LC-derivative, CF-derivative, and ABC-derivative. Additionally, the tables provide insights into the extent of absolute error between consecutive steps when encountering a non-integer order for the LC-derivative, CF-derivative, and ABC-derivative. The results from these tables indicate that the order of the error is notably minimal. This observation serves as a strong indication of the precision and efficacy of the approach presented in this study.

Table 1. The absolute error between Equations (7) and (8) and the exact solution (44) is represented by the first column and the second column while the absolute error of two steps of (7) and (8) is represented by the third column and the fourth column for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6$, and $a = 1$.

α	$ \phi(\alpha, \beta) - \phi_{LC}(\alpha_n, \beta_n) $	$ \varphi(\alpha, \beta) - \varphi_{LC}(\alpha_n, \beta_n) $	$ \phi_{LC,n} - \phi_{LC,n+1} $	$ \varphi_{LC,n} - \varphi_{LC,n+1} $
0	1.11022×10^{-16}	1.11022×10^{-16}	4.97499×10^{-11}	4.97495×10^{-11}
0.2	8.30951×10^{-5}	2.35526×10^{-5}	1.12725×10^{-6}	4.48712×10^{-7}
0.4	1.35067×10^{-4}	4.38737×10^{-5}	1.69840×10^{-6}	7.35767×10^{-7}
0.6	2.28422×10^{-4}	9.50490×10^{-5}	2.41060×10^{-6}	1.11918×10^{-6}
0.8	3.85878×10^{-4}	1.56810×10^{-4}	4.12192×10^{-6}	1.81449×10^{-6}
1	1.11022×10^{-16}	1.11022×10^{-16}	3.63416×10^{-7}	3.63416×10^{-7}

Table 2. In the case of the CF-derivative, the absolute error between Equations (25) and (26) and the exact solution (44) is represented by the first column and the second column while the absolute error of two steps of (25) and (26) is represented by the third column and the fourth column for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6$, and $a = 1$.

α	$ \phi(\alpha, \beta) - \phi_{CF}(\alpha_n, \beta_n) $	$ \varphi(\alpha, \beta) - \varphi_{CF}(\alpha_n, \beta_n) $	$ \phi_{CF,n} - \phi_{CF,n+1} $	$ \varphi_{CF,n} - \varphi_{CF,n+1} $
0	1.92593×10^{-34}	0	4.97498×10^{-11}	4.97501×10^{-11}
0.2	2.75570×10^{-5}	1.59500×10^{-4}	3.27900×10^{-7}	1.07397×10^{-6}
0.4	1.89708×10^{-4}	4.38737×10^{-5}	2.55455×10^{-7}	1.27453×10^{-6}
0.6	2.49232×10^{-6}	1.04730×10^{-4}	1.94268×10^{-7}	8.70804×10^{-7}
0.8	8.67747×10^{-6}	1.73810×10^{-4}	2.06507×10^{-7}	1.54168×10^{-6}
1	3.851860×10^{-34}	1.11022×10^{-16}	3.63417×10^{-7}	3.63417×10^{-7}

Table 3. In the case of the ABC-derivative, the absolute error between Equations (40) and (41) and the exact solution (44) is represented by the first column and the second column while the absolute error of two steps of (40) and (41) is represented by the third column and the fourth column for $\nu_1 = 2.8, \nu_2 = 0.8, c = 1, \mu_1 = 7, \mu_2 = -6$, and $a = 1$.

α	$ \phi(\alpha, \beta) - \phi_{ABC}(\alpha_n, \beta_n) $	$ \varphi(\alpha, \beta) - \varphi_{ABC}(\alpha_n, \beta_n) $	$ \phi_{CF,n} - \phi_{ABC,n+1} $	$ \varphi_{CF,n} - \varphi_{ABC,n+1} $
0	1.11022×10^{-16}	0	4.97496×10^{-11}	4.97498×10^{-11}
0.2	1.75119×10^{-5}	1.50089×10^{-4}	1.29273×10^{-7}	1.11919×10^{-6}
0.4	$2.283092244170781 \times 10^{-5}$	1.54883×10^{-4}	5.88663×10^{-8}	1.09897×10^{-6}
0.6	1.41835×10^{-6}	3.42845×10^{-6}	1.25380×10^{-7}	1.04458×10^{-7}
0.8	$3.190648624473713 \times 10^{-5}$	1.29585×10^{-5}	3.09597×10^{-8}	1.75898×10^{-7}
1	0	0	3.63415×10^{-7}	3.63417×10^{-7}

5. Conclusions

The problem of multi-space fractional-order coupled Korteweg-de Vries, which involves various kernels, was successfully converted into a system of differential equations. Subsequently, these equations were converted into algebraic forms employing the non-standard finite difference approach.

We employed the commonly used numerical technique, often referred to as the Newton-Raphson method, to address the resultant set of nonlinear algebraic equations. The precision of the examined approximations was confirmed by computing the absolute error between the exact solutions and the approximations. Additionally, we determined the absolute error between successive approximations in the scenario of non-integer orders.

We verified the accuracy of the analyzed approximations by calculating the absolute discrepancy between the exact solutions and the approximations. Furthermore, we assessed the absolute error between consecutive approximations in cases involving non-integer orders.

In future work, the current study's findings can be extended to analyze the behavior of positive solutions [43,44]. The scope of the investigation may also be broadened to encompass space-time. Furthermore, the study discussed in this paper can be extended to cover other special functions, such as Legendre and Chebyshev polynomials. In this context, recent research has been conducted on the subject of fractional derivatives and special functions, as in [45–48].

Author Contributions: K.M.S. performed the formal analysis of the investigation, the methodology, the software, and wrote the first draft of the paper. R.S. suggested and initiated this work, performed its validation, as well as reviewed and edited the paper. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Deanship of Scientific Research at Najran University under grant number (NU/DRP/SERC/12/16).

Data Availability Statement: The authors declare that data supporting the findings of this study are available within the article, the figures are concrete expression.

Acknowledgments: The authors are thankful to the Deanship of Scientific Research at Najran University for funding this work under the Distinguished Research Funding Program Grant Code (NU/DRP/SERC/12/16).

Conflicts of Interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. The authors declare that there are no conflicts of interests with publication of this work.

References

1. Kilbas, A.A.; Srivastava, H.M.; Trujillo, J.J. Theory and Applications of Fractional Differential Equations. In *North-Holland Mathematical Studies*; Elsevier: Amsterdam, The Netherlands; London, UK; New York, NY, USA, 2006; Volume 204.
2. Podlubny, I. *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications, Mathematics in Science and Engineering*; Academic Press: New York, NY, USA; London, UK; Sydney, Australia; Tokyo, Japan; Toronto, ON, Canada, 1999; Volume 198.
3. Srivastava, H.M. Fractional-order derivatives and integrals: Introductory overview and recent developments. *Kyungpook Math. J.* **2020**, *60*, 73–116.
4. Samko, S.G.; Kilbas, A.A.; Marichev, O.I. *Fractional Integrals and Derivatives: Theory and Applications*; Gordon and Breach: Yverdon, Switzerland, 1993.
5. Miller, K.S.; Ross, B. *An Introduction to the Fractional Calculus and Fractional Differential Equations*; John Wiley and Sons.: New York, NY, USA; Chichester, UK; Brisbane, Australia; Toronto, ON, Canada, 1993.
6. Sabir, Z.; Said, S.B.; Al-Mdallal, Q. A fractional order numerical study for the influenza disease mathematical model. *Alexandria Engng. J.* **2023**, *65*, 615–626. [[CrossRef](#)]
7. Alqhtani, M.; Owolabi, K.M.; Saad, K.M. Spatiotemporal (target) patterns in sub-diffusive predator-prey system with the Caputo operator. *Chaos Solitons Fractals* **2022**, *160*, 112267. [[CrossRef](#)]
8. Abdoon, M.A.; Hasan, F.L. Advantages of the differential equations for solving problems in mathematical physics with symbolic computation. *Math. Model. Engng. Probl.* **2022**, *9*, 268–276. [[CrossRef](#)]
9. Srivastava, H.M.; Saad, K.M.; Hamanah, W.M. Certain new models of the multi-space fractal-fractional Kuramoto-Sivashinsky and Korteweg-de Vries equations. *Mathematics* **2022**, *10*, 1089. [[CrossRef](#)]
10. Waheed, W.; Deng, G.; Liu, B. Discrete Laplacian operator and its Applications in signal processing. *IEEE Access* **2020**, *8*, 89692–89707. [[CrossRef](#)]
11. Lin, Z.; Wang, H. Modeling and application of fractional-order economic growth model with time delay. *Fractal Fract.* **2021**, *5*, 74. [[CrossRef](#)]
12. Yang, X.; Zhang, Z. On conservative, positivity preserving, nonlinear FV scheme on distorted meshes for the multi-term nonlocal Nagumo-type equations. *Appl. Math. Lett.* **2024**, *150*, 108972. [[CrossRef](#)]
13. Wang, J.; Jiang, X.; Yang, X.; Zhang, H. A nonlinear compact method based on double reduction order scheme for the nonlocal fourth-order PDEs with Burgers' type nonlinearity. *J. Appl. Math. Comput.* **2024**. [[CrossRef](#)]
14. Xiao, W.; Yang, X.; Zhou, Z. Pointwise-in-time α -robust error estimate of the ADI difference scheme for three-dimensional fractional subdiffusion equations with variable coefficients. *Commun. Anal. Mech.* **2024**, *16*, 53–70. [[CrossRef](#)]
15. Pakhira, R.; Ghosh, U.; Garg, H. An inventory model for partial backlogging items with memory effect. *Soft Comput.* **2023**, *27*, 9533–9550. [[CrossRef](#)]
16. Korteweg, D.J.; de Vries, G. On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves. *Philos. Mag.* **1895**, *39*, 422–443. [[CrossRef](#)]
17. Wu, Y.; Geng, X.; Hu, X.; Zhu, S. A generalized Hirota-Satsuma coupled Korteweg-de Vries equation and Miura transformations. *Phys. Lett. A* **1999**, *255*, 259–264. [[CrossRef](#)]
18. Hirota, R.; Satsuma, J. Soliton solutions of a coupled Korteweg-de Vries equation. *Phys. Lett. A* **1981**, *85*, 407–418. [[CrossRef](#)]
19. Srivastava, H.M.; Mandal, H.; Bira, B. Lie symmetry and exact solution of the time-fractional Hirota-Satsuma Korteweg-de Vries system. *Russian J. Math. Phys.* **2021**, *28*, 284–292. [[CrossRef](#)]
20. Dabiri, A.; Butcher, E.A. Efficient modified Chebyshev differentiation matrices for fractional differential equations. *Commun. Nonlinear Sci. Numer. Simul.* **2017**, *50*, 284–310. [[CrossRef](#)]
21. Yang, X.-J.; Hristov, J.; Srivastava, H.M.; Ahmad, B. Modelling fractal waves on shallow water surfaces via local fractional Korteweg-de Vries equation. *Abstr. Appl. Anal.* **2014**, *2014*, 278672. [[CrossRef](#)]
22. Baleanu, D.; Shiri, B.; Srivastava, H.M.; Qurashi, M.A. A Chebyshev spectral method based on operational matrix for fractional differential equations involving non-singular Mittag-Leffler kernel. *Adv. Differ. Equ.* **2018**, *2018*, 353. [[CrossRef](#)]
23. Hadhoud, A.R.; Srivastava, H.M.; Rageh, A.A.M. Non-polynomial B-spline and shifted Jacobi spectral collocation techniques to solve time-fractional nonlinear coupled Burgers' equations numerically. *Adv. Differ. Equ.* **2021**, *2021*, 439. [[CrossRef](#)]
24. Wang, W.; Zhang, H.; Jiang, X.; Yang, X. A high-order and efficient numerical technique for the nonlocal neutron diffusion equation representing neutron transport in a nuclear reactor. *Ann. Nucl. Energy* **2024**, *195*, 110163. [[CrossRef](#)]
25. Wang, W.; Wang, Y.; Zhu, B.; Chen, H. Robust H^1 -norm convergence analysis of L^1 FEM-ADI scheme for 2D/3D subdiffusion equation with initial singularity. *Math. Methods Appl. Sci.* **2023**, *46*, 16144–16155. [[CrossRef](#)]
26. Zhang, H.; Yang, X.; Tang, Q.; Xu, D. A robust error analysis of the OSC method for a multi-term fourth-order sub-diffusion equation. *Comput. Math. Appl.* **2022**, *109*, 180–190. [[CrossRef](#)]
27. Zhang, H.; Liu, Y.; Yang, X. An efficient ADI difference scheme for the nonlocal evolution problem in three-dimensional space. *J. Appl. Math. Comput.* **2023**, *69*, 651–674. [[CrossRef](#)]
28. Tian, Q.; Yang, X.; Zhang, H.; Xu, D. An implicit robust numerical scheme with graded meshes for the modified Burgers model with nonlocal dynamic properties. *Comp. Appl. Math.* **2023**, *42*, 246. [[CrossRef](#)]
29. Khader, M.M.; Saad, K.M. A numerical approach for solving the fractional Fisher equation using Chebyshev spectral collocation method. *Chaos Solitons Fractals* **2018**, *110*, 169–177. [[CrossRef](#)]

30. Khader, M.M.; Saad, K.M.; Hammouch, Z.; Baleanu, D. A spectral collocation method for solving fractional KdV and KdV-Burgers equations with non-singular kernel derivatives. *Appl. Numer. Math.* **2021**, *161*, 137–146. [[CrossRef](#)]
31. Saad, K.M.; Khader, M.M.; Gómez-Aguilar, J.F.; Baleanu, D. Numerical solutions of the fractional Fisher's type equations with Atangana-Baleanu fractional derivative by using spectral collocation methods. *Chaos* **2019**, *29*, 023116. [[CrossRef](#)] [[PubMed](#)]
32. Horadam, A.F. *Vieta Polynomials*; The University of New England: Armidaie, Australia, 2000.
33. Zakaria, M.; Khader, M.M.; Al-Dayel, I.; Al-Tayeb, W. Solving fractional generalized Fisher-Kolmogorov-Petrovsky-Piskunov's equation using compact finite different method together with spectral collocation algorithms. *J. Math.* **2022**, *2022*, 1901131.
34. Mickens, R.E. *Applications of Nonstandard Finite Difference Schemes*; World Scientific: Singapore, 2000.
35. Caputo, M.; Fabrizio, M. A new definition of fractional derivative without singular kernel. *Progr. Fract. Differ. Appl.* **2015**, *1*, 1–13.
36. Atangana, A.; Baleanu, D. New fractional derivative with non-local and non-singular kernel. *Thermal Sci.* **2016**, *20*, 757–763. [[CrossRef](#)]
37. Srivastava, H.M. Some parametric and argument variations of the operators of fractional calculus and related special functions and integral transformations. *J. Nonlinear Convex Anal.* **2021**, *22*, 1501–1520.
38. Srivastava, H.M. An introductory overview of fractional-calculus operators based upon the Fox-Wright and related higher transcendental functions. *J. Adv. Engrg. Comput.* **2021**, *5*, 135–166. [[CrossRef](#)]
39. Agarwal, P.; El-Sayed, A.A. Vieta-Lucas polynomials for solving a fractional-order mathematical physics model. *Adv. Differ. Equ.* **2020**, *2020*, 626. [[CrossRef](#)]
40. Loh, J.R.; Isah, A.; Phang, C.; Toh, Y.T. On the new properties of Caputo-Fabrizio operator and its application in deriving shifted Legendre operational matrix. *Appl. Numer. Math.* **2018**, *132*, 138–153. [[CrossRef](#)]
41. Abdeljawad, T.; Baleanu, D. Integration by parts and its applications of a new nonlocal fractional derivative with Mittag-Leffler nonsingular kernel. *J. Nonlinear Sci. Appl.* **2017**, *9*, 1098–1107. [[CrossRef](#)]
42. Srivastava, H.M. A survey of some recent developments on higher transcendental functions of analytic number theory and applied mathematics. *Symmetry* **2021**, *13*, 2294. [[CrossRef](#)]
43. Yang, X.; Zhang, H.; Zhang, Q.; Yuan, G. Simple positivity-preserving nonlinear finite volume scheme for subdiffusion equations on general non-conforming distorted meshes. *Nonlinear Dyn.* **2022**, *108*, 3859–3886. [[CrossRef](#)]
44. Yang, X.; Zhang, H. The uniform l1 long-time behavior of time discretization for time-fractional partial differential equations with nonsmooth data. *Appl. Math. Lett.* **2022**, *124*, 107644. [[CrossRef](#)]
45. Ibrahim, R.W.; Obaiys, S.J.; Karaca, Y.; Secer, A. Complex Mathematical Modeling for Advanced Fractal-Fractional Differential Operators within Symmetry. *Fractals* **2023**, *31*, 2340194. [[CrossRef](#)]
46. Ibrahim, R.W.; Obaiys, S.J.; Karaca, Y.; Binti Jamaludin, N.A. Symmetrically Conformable Fractional Differential Operators by Computational Numerical Modeling with Special Function. *Fractals* **2023**, *31*, 2340157. [[CrossRef](#)]
47. Srivastava, H.M. An introductory overview of Bessel polynomials, the generalized Bessel polynomials and the q -Bessel polynomials. *Symmetry* **2023**, *15*, 822. [[CrossRef](#)]
48. Srivastava, H.M.; Saad, K.M.; Khader, M.M.; Singh, H. Spectral collocation method based upon special functions for fractional partial differential equations. In *Chapman and Hall/CRC*; Elsevier: Amsterdam, The Netherlands, 2022; Volume 1, Number 1; pp. 79–101. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.