
Faculty of Science

Faculty Publications

Coefficients of a Comprehensive Subclass of Meromorphic Bi-Univalent Functions
Associated with the Faber Polynomial Expansion

Hari Mohan Srivastava, Ahmad Motamednezhad, & Safa Salehian

March 2021

© 2021 H. M. Srivastava et al. This is an open access article distributed under the terms of
the Creative Commons Attribution License <https://creativecommons.org/licenses/by/4.0/>

This article was originally published at:

<https://doi.org/10.3390/axioms10010027>

Citation for this paper:

Srivastava, H. M., Motamednezhad, A., & Salehian, S. (2021). Coefficients of a
Comprehensive Subclass of Meromorphic Bi-Univalent Functions Associated with the Faber
Polynomial Expansion. *Axioms*, 10(1), 1-13. <https://doi.org/10.3390/axioms10010027>.

Article

Coefficients of a Comprehensive Subclass of Meromorphic Bi-Univalent Functions Associated with the Faber Polynomial Expansion

Hari Mohan Srivastava ^{1,2,3,4,*} , Ahmad Motamednezhad ⁵  and Safa Salehian ⁶ 

¹ Department of Mathematics and Statistics, University of Victoria, Victoria, BC V8W 3R4, Canada

² Department of Medical Research, China Medical University Hospital, China Medical University, Taichung 40402, Taiwan

³ Department of Mathematics and Informatics, Azerbaijan University, 71 Jeyhun Hajibeyli Street, Baku AZ1007, Azerbaijan

⁴ Section of Mathematics, International Telematic University Uninettuno, I-00186 Rome, Italy

⁵ Faculty of Mathematical Sciences, Shahrood University of Technology, Shahrood P.O. Box 316-36155, Iran; a.motamedne@gmail.com

⁶ Department of Mathematics, Gorgan Branch, Islamic Azad University, Gorgan P.O. Box 717, Iran; s.salehian84@gmail.com

* Correspondence: harimsri@math.uvic.ca

Abstract: In this paper, we introduce a new comprehensive subclass $\Sigma_B(\lambda, \mu, \beta)$ of meromorphic bi-univalent functions in the open unit disk \mathbb{U} . We also find the upper bounds for the initial Taylor-Maclaurin coefficients $|b_0|$, $|b_1|$ and $|b_2|$ for functions in this comprehensive subclass. Moreover, we obtain estimates for the general coefficients $|b_n|$ ($n \geq 1$) for functions in the subclass $\Sigma_B(\lambda, \mu, \beta)$ by making use of the Faber polynomial expansion method. The results presented in this paper would generalize and improve several recent works on the subject.

Keywords: analytic functions; univalent and bi-univalent functions; meromorphic bi-univalent functions; coefficient estimates; Faber polynomial expansion; meromorphic bi-Bazilevič functions of order β and type μ ; meromorphic bi-starlike functions of order β



Citation: Srivastava, H.M.; Motamednezhad, A.; Salehian, S. Coefficients of a Comprehensive Subclass of Meromorphic Bi-Univalent Functions Associated with the Faber Polynomial Expansion. *Axioms* **2021**, *10*, 27. <https://doi.org/10.3390/axioms10010027>

Academic Editor: Clemente Cesarano

Received: 10 February 2021

Accepted: 25 February 2021

Published: 27 February 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Let \mathcal{A} denote the class of functions f of the form:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1)$$

which are analytic in the open unit disk

$$\mathbb{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}.$$

We also let \mathcal{S} be the class of functions $f \in \mathcal{A}$ which are univalent in \mathbb{U} .

It is well known that every function $f \in \mathcal{S}$ has an inverse f^{-1} , which is defined by

$$f^{-1}(f(z)) = z \quad (z \in \mathbb{U})$$

and

$$f(f^{-1}(w)) = w \quad \left(|w| < r_0(f); r_0(f) \geq \frac{1}{4}\right).$$

If f and f^{-1} are univalent in \mathbb{U} , then f is said to be bi-univalent in \mathbb{U} . We denote by σ_B the class of bi-univalent functions in \mathbb{U} . For a brief history and interesting examples of functions in the class σ_B , see the pioneering work [1]. In fact, this widely-cited work

by Srivastava et al. [1] actually revived the study of analytic and bi-univalent functions in recent years, and it has also led to a flood of papers on the subject by (for example) Srivastava et al. [2–14] and by others [15,16].

In this paper, let Σ be the family of meromorphic univalent functions f of the following form:

$$f(z) = z + b_0 + \sum_{n=1}^{\infty} \frac{b_n}{z^n}, \tag{2}$$

which are defined on the domain

$$\Delta = \{z : z \in \mathbb{C} \text{ and } 1 < |z| < \infty\}.$$

Since a function $f \in \Sigma$ is univalent, it has an inverse f^{-1} that satisfies the following relationship:

$$f^{-1}(f(z)) = z \quad (z \in \Delta)$$

and

$$f(f^{-1}(w)) = w \quad (M < |w| < \infty; M > 0).$$

Furthermore, the inverse function f^{-1} has a series expansion of the form [17]:

$$g(w) = f^{-1}(w) = w + \sum_{n=0}^{\infty} \frac{B_n}{w^n} \quad (M < |w| < \infty).$$

A function $f \in \Sigma$ is said to be meromorphic bi-univalent if both f and f^{-1} are meromorphic univalent in Δ . The family of all meromorphic bi-univalent functions in Δ of the form (2) is denoted by $\Sigma_{\mathcal{M}}$. A simple calculation shows that (see also [18,19])

$$g(w) = f^{-1}(w) = w - b_0 - \frac{b_1}{w} - \frac{b_2 + b_0 b_1}{w^2} - \dots \tag{3}$$

Moreover, the coefficients of $g = f^{-1}$ can be given in terms of the *Faber polynomial* [20] (see also [21–23]) as follows:

$$g(w) = f^{-1}(w) = w - b_0 - \sum_{n=1}^{\infty} \frac{1}{n} K_{n+1} \frac{1}{w^n} \quad (w \in \Delta), \tag{4}$$

where

$$K_{n+1} = nb_0^{n-1}b_1 + n(n-1)b_0^{n-2}b_2 + \frac{1}{2}n(n-1)(n-2)b_0^{n-3}(b_3 + b_1^2) + \frac{n(n-1)(n-2)(n-3)}{3!}b_0^{n-4}(b_4 + 3b_1b_2) + \sum_{j \geq 5} b_0^{n-j}V_j$$

and V_j (with $5 \leq j \leq n$) is a homogeneous polynomial of degree j in the variables b_1, b_2, \dots, b_n .

Estimates on the coefficients of meromorphic univalent functions were widely investigated in the literature. For example, Schiffer [24] obtained the estimate $|b_2| \leq 2/3$ for meromorphic univalent functions $f \in \Sigma$ with $b_0 = 0$ and Duren [25] proved that

$$|b_n| \leq \frac{2}{n+1} \quad \left(f \in \Sigma; b_k = 0; 1 \leq k < \frac{n}{2}\right).$$

Many researchers introduced and studied subclasses of meromorphic bi-univalent functions (see, for instance, Janani et al. [26], Orhan et al. [27] and others [28–30]).

Recently, Srivastava et al. [31] introduced a new class $\Sigma_{B^*}(\lambda, \beta)$ of meromorphic bi-univalent functions and obtained the estimates on the initial Taylor–Maclaurin coefficients $|b_0|$ and $|b_1|$ for functions in this class.

Definition 1 (see [31]). A function $f \in \Sigma_{\mathcal{M}}$, given by (2), is said to be in the class $\Sigma_{B^*}(\lambda, \beta)$ ($\lambda \geq 1$; $0 \leq \beta < 1$), if the following conditions are satisfied:

$$\Re\left(\frac{z(f'(z))^\lambda}{f(z)}\right) > \beta$$

and

$$\Re\left(\frac{w(g'(w))^\lambda}{g(w)}\right) > \beta,$$

where the function g , given by (3) is the inverse of f and $z, w \in \Delta$.

Theorem 1 (see [31]). Let the function $f \in \Sigma_{\mathcal{M}}$, given by (2), be in the class $\Sigma_{B^*}(\lambda, \beta)$. Then,

$$|b_0| \leq 2(1 - \beta) \quad \text{and} \quad |b_1| \leq \frac{2(1 - \beta)\sqrt{4\beta^2 - 8\beta + 5}}{1 + \lambda}.$$

In this paper, we introduce a new comprehensive subclass $\Sigma_B(\lambda, \mu, \beta)$ of the meromorphic bi-univalent function class $\Sigma_{\mathcal{M}}$. We also obtain estimates for the initial Taylor–Maclaurin coefficients b_0, b_1 and b_2 for functions in this subclass. Furthermore, we find estimates for the general coefficients b_n ($n \geq 1$) for functions in this comprehensive subclass $\Sigma_B(\lambda, \mu, \beta)$ by using the Faber polynomials [20]. Our results for the meromorphic bi-univalent function subclass $\Sigma_B(\lambda, \mu, \beta)$ would generalize and improve some recent works by Srivastava et al. [31], Hamidi et al. [32] and Jahangiri et al. [33] (see also the recent works [34,35]).

2. Preliminary Results

For finding the coefficients of functions belonging to the function class $\Sigma_B(\lambda, \mu, \beta)$, we need the following lemmas and remarks.

Lemma 1 (see [21,22]). Let f be the function given by

$$f(z) = z + b_0 + \frac{b_1}{z} + \frac{b_2}{z^2} + \dots$$

be a meromorphic univalent function defined on the domain Δ . Then, for any $\rho \in \mathbb{R}$, there are polynomials K_n^ρ such that

$$\left(\frac{f(z)}{z}\right)^\rho = 1 + \sum_{n=1}^{\infty} \frac{K_n^\rho(b_0, b_1, \dots, b_{n-1})}{z^n},$$

where

$$K_n^\rho(b_0, b_1, \dots, b_{n-1}) = \rho b_{n-1} + \frac{\rho(\rho - 1)}{2} D_n^2 + \frac{\rho!}{(\rho - 3)!3!} D_n^3 + \dots + \frac{\rho!}{(\rho - n)!n!} D_n^n$$

and

$$D_n^k(x_1, x_2, \dots, x_{n-k+1}) = \sum \frac{k!(x_1)^{\mu_1} \dots (x_{n-k+1})^{\mu_{n-k+1}}}{\mu_1! \dots \mu_{n-k+1}!},$$

in which the sum is taken over all non-negative integers $\mu_1, \dots, \mu_{n-k+1}$ such that

$$\begin{cases} \mu_1 + \mu_2 + \dots + \mu_{n-k+1} = k \\ \mu_1 + 2\mu_2 + \dots + (n - k + 1)\mu_{n-k+1} = n. \end{cases}$$

The first three terms of K_n^ρ are given by

$$K_1^\rho(b_0) = \rho b_0,$$

$$K_2^\rho(b_0, b_1) = \rho b_1 + \frac{\rho(\rho - 1)}{2} b_0^2$$

and

$$K_3^\rho(b_0, b_1, b_2) = \rho b_2 + \rho(\rho - 1)b_0 b_1 + \frac{\rho(\rho - 1)(\rho - 2)}{3!} b_0^3.$$

Remark 1. In the special case when

$$b_0 = b_1 = \dots = b_{n-1} = 0,$$

it is easily seen that

$$K_i^\rho(b_0, \dots, b_{i-1}) = 0 \quad (1 \leq i \leq n)$$

and

$$K_{n+1}^\rho(b_0, b_1, \dots, b_n) = \rho b_n.$$

Lemma 2 (see [21,22]). Let f be the function given by

$$f(z) = z + b_0 + \frac{b_1}{z} + \frac{b_2}{z^2} + \dots$$

be a meromorphic univalent function defined on the domain Δ . Then, the Faber polynomials F_n of $f(z)$ are given by

$$\frac{zf'(z)}{f(z)} = 1 + \sum_{n=1}^{\infty} \frac{F_n(b_0, b_1, \dots, b_{n-1})}{z^n}, \tag{5}$$

where $F_n(b_0, b_1, \dots, b_{n-1})$ is a homogeneous polynomial of degree n .

Remark 2 (see [36]). For any integer $n \geq 1$, the polynomials $F_n(b_0, b_1, \dots, b_{n-1})$ are given by

$$F_n(b_0, b_1, \dots, b_{n-1}) = \sum_{i_1+2i_2+\dots+ni_n=n} A_{(i_1,i_2,\dots,i_n)} b_0^{i_1} b_1^{i_2} \dots b_{n-1}^{i_n},$$

where

$$A_{(i_1,i_2,\dots,i_n)} := (-1)^{n+2i_1+3i_2+\dots+(n+1)i_n} \frac{(i_1 + i_2 + \dots + i_n - 1)! n}{i_1! i_2! \dots i_n!}.$$

The first three terms of F_n are given by

$$F_1(b_0) = -b_0,$$

$$F_2(b_0, b_1) = b_0^2 - 2b_1$$

and

$$F_3(b_0, b_1, b_2) = -b_0^3 + 3b_0 b_1 - 3b_2.$$

Remark 3. In the special case when $b_0 = b_1 = \dots = b_{n-1} = 0$, it is readily observed that

$$F_i(b_0, \dots, b_{i-1}) = 0 \quad (1 \leq i \leq n)$$

and

$$F_{n+1}(b_0, b_1, \dots, b_n) = (-1)^{2n+3}(n+1)b_n = -(n+1)b_n.$$

Lemma 3. Let f be the function given by

$$f(z) = z + b_0 + \frac{b_1}{z} + \frac{b_2}{z^2} + \dots$$

be a meromorphic univalent function defined on the domain Δ . Then, for $\lambda \geq 1$ and $\mu \geq 0$,

$$\left(\frac{zf'(z)}{f(z)}\right)^\lambda \left(\frac{f(z)}{z}\right)^\mu = 1 + \sum_{n=1}^\infty \frac{L_n(b_0, b_1, \dots, b_{n-1})}{z^n},$$

where

$$L_n(b_0, b_1, \dots, b_{n-1}) = \sum_{i=0}^n K_{n-i}^\lambda(F_1, \dots, F_{n-i})K_i^\mu(b_0, \dots, b_{i-1}) \quad (K_0^\lambda = K_0^\mu = 1)$$

and $F_n = F_n(b_0, b_1, \dots, b_{n-1})$ is given by (5).

Proof. By using Lemmas 1 and 2, we have

$$\begin{aligned} \left(\frac{zf'(z)}{f(z)}\right)^\lambda \left(\frac{f(z)}{z}\right)^\mu &= \left(1 + \sum_{m=1}^\infty \frac{F_m(b_0, b_1, \dots, b_{m-1})}{z^m}\right)^\lambda \\ &\cdot \left(1 + \sum_{m=1}^\infty \frac{K_m^\mu(b_0, b_1, \dots, b_{m-1})}{z^m}\right). \end{aligned}$$

In addition, by applying Lemma 1 once again, we obtain

$$\begin{aligned} \left(\frac{zf'(z)}{f(z)}\right)^\lambda \left(\frac{f(z)}{z}\right)^\mu &= \left(1 + \sum_{m=1}^\infty \frac{K_m^\lambda(F_1, \dots, F_m)}{z^m}\right) \\ &\cdot \left(1 + \sum_{m=1}^\infty \frac{K_m^\mu(b_0, \dots, b_{m-1})}{z^m}\right) \\ &= 1 + \sum_{n=1}^\infty \sum_{i=0}^n K_{n-i}^\lambda(F_1, \dots, F_{n-i})K_i^\mu(b_0, \dots, b_{i-1}) \frac{1}{z^n} \\ &\quad (K_0^\lambda = K_0^\mu = 1). \end{aligned}$$

Our demonstration of Lemma 3 is thus completed. \square

The first three terms of L_n are given by

$$L_1(b_0) = (\mu - \lambda)b_0,$$

$$L_2(b_0, b_1) = \frac{\lambda(1 + \lambda - 2\mu) + \mu(\mu - 1)}{2}b_0^2 + (\mu - 2\lambda)b_1$$

and

$$\begin{aligned} L_3(b_0, b_1, b_2) &= \left(\frac{\lambda(2 - \mu)(\mu - \lambda)}{2} + \frac{\mu(\mu - 1)(\mu - 2) - \lambda(\lambda - 1)(\lambda - 2)}{6}\right)b_0^3 \\ &\quad + [\lambda(2\lambda + 1) + \mu(\mu - 3\lambda - 1)]b_0b_1 + (\mu - 3\lambda)b_2. \end{aligned}$$

Remark 4. In the special case when $b_0 = b_1 = \dots = b_{n-1} = 0$, we easily find that

$$L_i(b_0, \dots, b_{i-1}) = 0 \quad (1 \leq i \leq n)$$

and

$$L_{n+1}(b_0, b_1, \dots, b_n) = (\mu - (n + 1)\lambda)b_n.$$

Lemma 4 (see [37]). *If the function $p \in \mathcal{P}$, then $|c_k| \leq 2$ for each k , where \mathcal{P} is the family of all functions p , which are analytic in the domain Δ given by*

$$\Delta = \{z : z \in \mathbb{C} \text{ and } 1 < |z| < \infty\}$$

for which

$$\Re(p(z)) > 0 \quad (z \in \Delta),$$

where

$$p(z) = 1 + \frac{c_1}{z} + \frac{c_2}{z^2} + \frac{c_3}{z^3} + \dots.$$

3. The Comprehensive Class $\Sigma_B(\lambda, \mu, \beta)$

In this section, we introduce and investigate the comprehensive class $\Sigma_B(\lambda, \mu, \beta)$ of meromorphic bi-univalent functions defined on the domain Δ .

Definition 2. *A function $f \in \Sigma_{\mathcal{M}}$, given by (2), is said to be in the class*

$$\Sigma_B(\lambda, \mu, \beta) \quad (\lambda \geq 1; \mu \geq 0; 0 \leq \beta < 1)$$

of meromorphic bi-univalent functions of order β and type μ , if the following conditions are satisfied:

$$\Re\left(\left(\frac{zf'(z)}{f(z)}\right)^\lambda \left(\frac{f(z)}{z}\right)^\mu\right) > \beta$$

and

$$\Re\left(\left(\frac{wg'(w)}{g(w)}\right)^\lambda \left(\frac{g(w)}{w}\right)^\mu\right) > \beta,$$

where the function g given by (4), is the inverse of f and $z, w \in \Delta$.

Remark 5. *There are several choices of the parameters λ and μ which would provide interesting subclasses of meromorphic bi-univalent functions. For example, we have the following special cases:*

- *By putting $\lambda = 1$ and $0 \leq \mu < 1$, the class $\Sigma_B(\lambda, \mu, \beta)$ reduces to the subclass $B(\beta, \mu)$ of meromorphic bi-Bazilevič functions of order β and type μ , which was considered by Jahangiri et al. [33].*
- *By putting $\lambda = 1$ and $\mu = 0$, the class $\Sigma_B(\lambda, \mu, \beta)$ reduces to the subclass $\Sigma_B^*(\beta)$ of meromorphic bi-starlike functions of order β , which was considered by Hamidi et al. [32].*
- *By putting $\mu = \lambda - 1$, the class $\Sigma_B(\lambda, \mu, \beta)$ reduces to the class $\Sigma_{B^*}(\lambda, \beta)$ in Definition 1.*

Theorem 2. *Let $f \in \Sigma_B(\lambda, \mu, \beta)$. If $b_0 = b_1 = \dots = b_{n-1} = 0$, then*

$$|b_n| \leq \frac{2(1 - \beta)}{|(n + 1)\lambda - \mu|} \quad (n \geq 1).$$

Proof. By using Lemma 3 for the meromorphic bi-univalent function f given by

$$f(z) = z + b_0 + \sum_{n=1}^{\infty} \frac{b_n}{z^n},$$

we have

$$\left(\frac{zf'(z)}{f(z)}\right)^\lambda \left(\frac{f(z)}{z}\right)^\mu = 1 + \sum_{n=0}^{\infty} \frac{L_{n+1}(b_0, b_1, \dots, b_n)}{z^{n+1}}. \tag{6}$$

Similarly, for its inverse map g given by

$$g(w) = f^{-1}(w) = w + B_0 + \sum_{n=1}^{\infty} \frac{B_n}{w^n},$$

we find that

$$\left(\frac{wg'(w)}{g(w)}\right)^{\lambda} \left(\frac{g(w)}{w}\right)^{\mu} = 1 + \sum_{n=0}^{\infty} \frac{L_{n+1}(B_0, B_1, \dots, B_n)}{w^{n+1}}. \tag{7}$$

Furthermore, since $f \in \Sigma_B(\lambda, \mu, \beta)$, by using Definition 2, there exist two positive real-part functions

$$c(z) = 1 + \sum_{n=1}^{\infty} c_n z^{-n}$$

and

$$d(w) = 1 + \sum_{n=1}^{\infty} d_n w^{-n}$$

for which

$$\Re(c(z)) > 0 \quad \text{and} \quad \Re(d(w)) > 0 \quad (z, w \in \Delta),$$

such that

$$\left(\frac{zf'(z)}{f(z)}\right)^{\lambda} \left(\frac{f(z)}{z}\right)^{\mu} = 1 + (1 - \beta) \sum_{n=0}^{\infty} K_{n+1}^1(c_1, c_2, \dots, c_{n+1}) \frac{1}{z^{n+1}} \tag{8}$$

and

$$\left(\frac{wg'(w)}{g(w)}\right)^{\lambda} \left(\frac{g(w)}{w}\right)^{\mu} = 1 + (1 - \beta) \sum_{n=0}^{\infty} K_{n+1}^1(d_1, d_2, \dots, d_{n+1}) \frac{1}{w^{n+1}}. \tag{9}$$

Upon equating the corresponding coefficients in (6) and (8), we get

$$L_{n+1}(b_0, b_1, \dots, b_n) = (1 - \beta) K_{n+1}^1(c_1, c_2, \dots, c_{n+1}). \tag{10}$$

Similarly, from (7) and (9), we obtain

$$L_{n+1}(B_0, B_1, \dots, B_n) = (1 - \beta) K_{n+1}^1(d_1, d_2, \dots, d_{n+1}). \tag{11}$$

Now, since $b_i = 0$ ($0 \leq i \leq n - 1$), we have

$$B_i = 0 \quad (0 \leq i \leq n - 1) \quad \text{and} \quad B_n = -b_n.$$

Hence, by using Remark 4, Equations (10) and (11) can be rewritten as follows:

$$(\mu - (n + 1)\lambda)b_n = (1 - \beta)c_{n+1} \tag{12}$$

and

$$-(\mu - (n + 1)\lambda)b_n = (1 - \beta)d_{n+1}, \tag{13}$$

respectively. Thus, from (12) and (13), we find that

$$2(\mu - (n + 1)\lambda)b_n = (1 - \beta)(c_{n+1} - d_{n+1}).$$

Finally, by applying Lemma 4, we get

$$|b_n| = \frac{(1 - \beta)|c_{n+1} - d_{n+1}|}{2|(n + 1)\lambda - \mu|} \leq \frac{2(1 - \beta)}{|(n + 1)\lambda - \mu|},$$

which completes the proof of Theorem 2 \square

Theorem 3. Let the function $f \in \mathcal{M}$, given by (2), be in the class

$$\Sigma_B(\lambda, \mu, \beta) \quad (\lambda \geq 1; \mu \geq 0; 0 \leq \beta < 1).$$

Then,

$$|b_0| \leq \min \left\{ \frac{2(1-\beta)}{|\mu-\lambda|}, 2\sqrt{\frac{1-\beta}{|\lambda(1+\lambda-2\mu)+\mu(\mu-1)|}} \right\},$$

$$|b_1| \leq \frac{2(1-\beta)}{|\mu-2\lambda|}$$

and

$$|b_2| \leq \frac{2\{|\lambda(2\lambda+4)+\mu(\mu-3\lambda-2)|+|\lambda(2\lambda+1)+\mu(\mu-3\lambda-1)|\}(1-\beta)}{|\mu-3\lambda|\lambda(4\lambda+5)+\mu(2\mu-6\lambda-3)|}$$

$$+ \frac{8|T(\mu, \lambda)|(1-\beta)^3}{|(\mu-3\lambda)(\mu-\lambda)^3|},$$

where

$$T(\mu, \lambda) = \frac{\lambda(2-\mu)(\mu-\lambda)}{2} + \frac{\mu(\mu-1)(\mu-2) - \lambda(\lambda-1)(\lambda-2)}{6}.$$

Proof. By putting $n = 0, 1, 2$ in (10), we get

$$(\mu-\lambda)b_0 = (1-\beta)c_1, \tag{14}$$

$$\frac{\lambda(1+\lambda-2\mu)+\mu(\mu-1)}{2}b_0^2 + (\mu-2\lambda)b_1 = (1-\beta)c_2 \tag{15}$$

and

$$T(\mu, \lambda)b_0^3 + [\lambda(2\lambda+1)+\mu(\mu-3\lambda-1)]b_0b_1 + (\mu-3\lambda)b_2 = (1-\beta)c_3. \tag{16}$$

Similarly, by putting $n = 0, 1, 2$ in (11), we have

$$-(\mu-\lambda)b_0 = (1-\beta)d_1, \tag{17}$$

$$\frac{\lambda(1+\lambda-2\mu)+\mu(\mu-1)}{2}b_0^2 - (\mu-2\lambda)b_1 = (1-\beta)d_2 \tag{18}$$

and

$$-T(\mu, \lambda)b_0^3 + (\lambda(2\lambda+4)+\mu(\mu-3\lambda-2))b_0b_1 - (\mu-3\lambda)b_2 = (1-\beta)d_3. \tag{19}$$

Clearly, from (14) and (17), we get

$$c_1 = -d_1 \tag{20}$$

and

$$b_0 = \frac{(1-\beta)c_1}{\mu-\lambda}. \tag{21}$$

Adding (15) and (18), we obtain

$$b_0^2 = \frac{(1-\beta)(c_2+d_2)}{\lambda(1+\lambda-2\mu)+\mu(\mu-1)}. \tag{22}$$

In view of the Equations (21) and (22), by applying Lemma 4, we get

$$|b_0| \leq \frac{2(1-\beta)}{|\mu-\lambda|} \quad \text{and} \quad |b_0|^2 \leq \frac{4(1-\beta)}{|\lambda(1+\lambda-2\mu) + \mu(\mu-1)|},$$

respectively. Thus, we get the desired estimate on the coefficient $|b_0|$.

Next, in order to find the bound on the coefficient $|b_1|$, we subtract (18) from (15). We thus obtain

$$b_1 = \frac{(1-\beta)(c_2-d_2)}{2(\mu-2\lambda)}. \tag{23}$$

Applying Lemma 4 once again, we get

$$|b_1| \leq \frac{2(1-\beta)}{|\mu-2\lambda|}.$$

Finally, in order to determine the bound on $|b_2|$, we consider the sum of the Equations (16) and (19) with $c_1 = -d_1$. This yields

$$b_0b_1 = \frac{(1-\beta)(c_3+d_3)}{\lambda(4\lambda+5) + \mu(2\mu-6\lambda-3)}. \tag{24}$$

Subtracting (19) from (16) with $c_1 = -d_1$, we obtain

$$2(\mu-3\lambda)b_2 + (\mu-3\lambda)b_0b_1 + 2T(\mu,\lambda)b_0^3 = (1-\beta)(c_3-d_3). \tag{25}$$

In addition, by using (21) and (24) in (25), we get

$$b_2 = \frac{(1-\beta)(c_3-d_3)}{2(\mu-3\lambda)} - \frac{(1-\beta)(c_3+d_3)}{2[\lambda(4\lambda+5) + \mu(2\mu-6\lambda-3)]} - \frac{T(\mu,\lambda)(1-\beta)^3c_1^3}{(\mu-3\lambda)(\mu-\lambda)^3}.$$

Hence,

$$b_2 = \frac{\{[\lambda(2\lambda+4) + \mu(\mu-3\lambda-2)]c_3 - [\lambda(2\lambda+1) + \mu(\mu-3\lambda-1)]d_3\}(1-\beta)}{(\mu-3\lambda)[\lambda(4\lambda+5) + \mu(2\mu-6\lambda-3)]} - \frac{T(\mu,\lambda)(1-\beta)^3c_1^3}{(\mu-3\lambda)(\mu-\lambda)^3}.$$

Thus, by applying Lemma 4 once again, we get

$$|b_2| \leq \frac{2\{|\lambda(2\lambda+4) + \mu(\mu-3\lambda-2)| + |\lambda(2\lambda+1) + \mu(\mu-3\lambda-1)|\}(1-\beta)}{|\mu-3\lambda|[\lambda(4\lambda+5) + \mu(2\mu-6\lambda-3)]} + \frac{8|T(\mu,\lambda)|(1-\beta)^3}{|(\mu-3\lambda)(\mu-\lambda)^3|}.$$

This completes the proof of Theorem 3. \square

4. A Set of Corollaries and Consequences

By setting $\lambda = 1$ and $0 \leq \mu < 1$ in Theorem 2, we have the following result.

Corollary 1. *Let the function $f \in \mathcal{M}$, given by (2), be in the subclass $B(\beta, \mu)$ of meromorphic bi-Bazilevič functions of order β and type μ . If*

$$b_0 = b_1 = \dots = b_{n-1} = 0,$$

then

$$|b_n| \leq \frac{2(1-\beta)}{n+1-\mu} \quad (n \geq 1).$$

Remark 6. The estimate of $|b_n|$, given in Corollary 1, is the same as the corresponding estimate given by Hamidi et al. [38] Corollary 3.3.

By setting $\mu = 0$ in Corollary 1, we have the following result.

Corollary 2. Let the function $f \in \mathcal{M}$, given by (2), be in the subclass $\Sigma_B^*(\beta)$ of meromorphic bi-starlike functions of order β . If

$$b_0 = b_1 = \dots = b_{n-1} = 0,$$

then

$$|b_n| \leq \frac{2(1-\beta)}{n+1} \quad (n \geq 1).$$

Remark 7. The estimate of $|b_n|$, given in Corollary 2, is the same as the corresponding estimate given by Hamidi et al. [38] Corollary 3.4.

By setting $\mu = \lambda - 1$ in Theorem 2, we have the following result.

Corollary 3. Let the function $f \in \mathcal{M}$, given by (2), be in the subclass $\Sigma_{B^*}(\lambda, \beta)$. If

$$b_0 = b_1 = \dots = b_{n-1} = 0,$$

then

$$|b_n| \leq \frac{2(1-\beta)}{n\lambda+1} \quad (n \geq 1).$$

Remark 8. Corollary 3 is a generalization of a result presented in Theorem 1, which was proved by Srivastava et al. [31].

By setting $\lambda = 1$ and $0 \leq \mu < 1$ in Theorem 3, we have the following result.

Corollary 4. Let the function $f \in \mathcal{M}$, given by (2), be in the subclass $B(\beta, \mu)$ of meromorphic bi-Bazilevič functions of order β and type μ . Then,

$$|b_0| \leq \begin{cases} \sqrt{\frac{4(1-\beta)}{(1-\mu)(2-\mu)}} & \left(0 \leq \beta \leq \frac{1}{2-\mu}\right) \\ \frac{2(1-\beta)}{1-\mu} & \left(\frac{1}{2-\mu} \leq \beta < 1\right), \end{cases}$$

$$|b_1| \leq \frac{2(1-\beta)}{2-\mu}$$

and

$$|b_2| \leq \frac{2(1-\beta)}{3-\mu} + \frac{4(2-\mu)(1-\beta)^3}{3(1-\mu)^2}.$$

Remark 9. Corollary 4 also contains the estimate of the Taylor–Maclaurin coefficient $|b_2|$ of functions in the subclass $B(\beta, \mu)$ (see [33]).

By setting $\mu = 0$ in Corollary 4, we have the following result.

Corollary 5. Let the function $f \in \mathcal{M}$, given by (2), be in the subclass $\Sigma_B^*(\beta)$ of meromorphic bi-starlike functions of order β . Then,

$$|b_0| \leq \begin{cases} \sqrt{2(1-\beta)} & (0 \leq \beta \leq \frac{1}{2}) \\ 2(1-\beta) & (\frac{1}{2} \leq \beta < 1), \end{cases}$$

$$|b_1| \leq 1 - \beta$$

and

$$|b_2| \leq \frac{2(1-\beta)}{3} + \frac{8(1-\beta)^3}{3}.$$

Remark 10. Corollary 5 not only improves the estimate of the Taylor–Maclaurin coefficient $|b_0|$, which was given by Hamidi et al. [32] Theorem 2, but it also provides an improvement of the known estimate of the Taylor–Maclaurin coefficient $|b_2|$ of functions in the subclass $\Sigma_B^*(\beta)$. Furthermore, the estimate of $|b_0|$, presented in Corollary 5, is the same as the corresponding estimate given by Hamidi et al. [38] Corollary 3.5.

By setting $\mu = \lambda - 1$ in Theorem 3, we have the following result.

Corollary 6. Let the function $f \in \mathcal{M}$, given by (2), be in the subclass $\Sigma_{B^*}(\lambda, \beta)$. Then,

$$|b_0| \leq \begin{cases} \sqrt{2(1-\beta)} & (0 \leq \beta \leq \frac{1}{2}) \\ 2(1-\beta) & (\frac{1}{2} \leq \beta < 1), \end{cases}$$

$$|b_1| \leq \frac{2(1-\beta)}{\lambda + 1}$$

and

$$|b_2| \leq \frac{2(1-\beta)}{2\lambda + 1} + \frac{8(1-\beta)^3}{2\lambda + 1}.$$

Remark 11. Corollary 6 improves the estimates of the Taylor–Maclaurin coefficients $|b_0|$ and $|b_1|$ in Theorem 1 of Srivastava et al. [31]. In fact, it also provides an improvement of the known estimate of the Taylor–Maclaurin coefficient $|b_2|$ of functions in the subclass $\Sigma_{B^*}(\lambda, \beta)$.

Remark 12. In his recently-published survey-cum-expository review article, Srivastava [39] demonstrated how the theories of the basic (or q -) calculus and the fractional q -calculus have significantly encouraged and motivated further developments in Geometric Function Theory of Complex Analysis (see, for example, [8,40–42]). This direction of research is applicable also to the results which we have presented in this article. However, as pointed out by Srivastava [39] (p. 340), any further attempts to easily (and possibly trivially) translate the suggested q -results into the corresponding (p, q) -results (with $0 < |q| < p \leq 1$) would obviously be inconsequential because the additional parameter p is redundant.

Author Contributions: All three authors contributed equally to this investigation. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Srivastava, H.M.; Mishra, A.K.; Gochhayat, P. Certain subclasses of analytic and bi-univalent functions. *Appl. Math. Lett.* **2010**, *23*, 1188–1192. [[CrossRef](#)]
2. Çağlar, M.; Deniz, E.; Srivastava, H.M. Second Hankel determinant for certain subclasses of bi-univalent functions. *Turk. J. Math.* **2017**, *41*, 694–706. [[CrossRef](#)]
3. Srivastava, H.M.; Bansal, D. Coefficient estimates for a subclass of analytic and bi-univalent functions. *J. Egypt. Math. Soc.* **2015**, *23*, 242–246. [[CrossRef](#)]
4. Srivastava, H.M.; Bulut, S.; Çağlar, M.; Yagmur, N. Coefficient estimates for a general subclass of analytic and bi-univalent functions. *Filomat* **2013**, *27*, 831–842. [[CrossRef](#)]
5. Srivastava, H.M.; Gaboury, S.; Ghanim, F. Coefficient estimates for some general subclasses of analytic and bi-univalent functions. *Afrika Mat.* **2017**, *28*, 693–706. [[CrossRef](#)]
6. Srivastava, H.M.; Gaboury, S.; Ghanim, F. Initial coefficient estimates for some subclasses of m -fold symmetric bi-univalent functions. *Acta Math. Sci. Ser. B Engl. Ed.* **2016**, *36*, 863–871. [[CrossRef](#)]
7. Srivastava, H.M.; Gaboury, S.; Ghanim, F. Coefficient estimates for some subclasses of m -fold symmetric bi-univalent functions. *Acta Univ. Apulensis Math. Inform.* **2015**, *23*, 153–164.
8. Srivastava, H.M.; Khan, S.; Ahmad, Q.Z.; Khan, N.; Hussain, S. The Faber polynomial expansion method and its application to the general coefficient problem for some subclasses of bi-univalent functions associated with a certain q -integral operator. *Stud. Univ. Babeş-Bolyai Math.* **2018**, *63*, 419–436. [[CrossRef](#)]
9. Srivastava, H.M.; Sakar, F.M.; Ö. Güney, H. Some general coefficient estimates for a new class of analytic and bi-univalent functions defined by a linear combination. *Filomat* **2018**, *34*, 1313–1322. [[CrossRef](#)]
10. Srivastava, H.M.; Sivasubramanian, S.; Sivakumar, R. Initial coefficient bounds for a subclass of m -fold symmetric bi-univalent functions. *Tbilisi Math. J.* **2014**, *7*, 1–10. [[CrossRef](#)]
11. Srivastava, H.M.; Sümer Eker, S.; Ali, R.M. Coefficient bounds for a certain class of analytic and bi-univalent functions. *Filomat* **2015**, *29*, 1839–1845. [[CrossRef](#)]
12. Srivastava, H.M.; Sümer Eker, S.; Hamidi, S.G.; Jahangiri, J.M. Faber polynomial coefficient estimates for bi-univalent functions defined by the Tremblay fractional derivative operator. *Bull. Iran. Math. Soc.* **2018**, *44*, 149–157. [[CrossRef](#)]
13. Srivastava, H.M.; Wanas, A.K. Initial Maclaurin coefficient bounds for new subclasses of analytic and m -fold symmetric bi-univalent functions defined by a linear combination. *Kyungpook Math. J.* **2019**, *59*, 493–503.
14. Srivastava, H.M.; Wanas, A.K.; Murugusundaramoorthy, G. A certain family of bi-univalent functions associated with the Pascal distribution series based upon the Horadam polynomials. *Surveys Math. Appl.* **2021**, *16*, 193–205.
15. Zireh, A.; Adegani, E.A.; Bulut, S. Faber polynomial coefficient estimates for a comprehensive subclass of analytic bi-univalent functions defined by subordination. *Bull. Belg. Math. Soc. Simon Stevin* **2016**, *23*, 487–504. [[CrossRef](#)]
16. Zireh, A.; Adegani, E.A.; Bidkham, M. Faber polynomial coefficient estimates for subclass of bi-univalent functions defined by quasi-subordinate. *Math. Slovaca* **2018**, *68*, 369–378. [[CrossRef](#)]
17. Panigrahi, T. Coefficient bounds for certain subclasses of meromorphic and bi-univalent functions. *Bull. Korean Math. Soc.* **2013**, *50*, 1531–1538. [[CrossRef](#)]
18. Schober, G. Coefficients of inverses of meromorphic univalent functions. *Proc. Am. Math. Soc.* **1977**, *67*, 111–116. [[CrossRef](#)]
19. Xiao, H.-G.; Xu, Q.-H. Coefficient estimates for three generalized classes of meromorphic and bi-univalent functions. *Filomat* **2015**, *29*, 1601–1612. [[CrossRef](#)]
20. Faber, G. Über polynomische Entwicklungen. *Math. Ann.* **1903**, *57*, 389–408. [[CrossRef](#)]
21. Airault, H.; Bouali, A. Differential calculus on the Faber polynomials. *Bull. Sci. Math.* **2006**, *130*, 179–222. [[CrossRef](#)]
22. Airault, H.; Ren, J. An algebra of differential operators and generating functions on the set of univalent functions. *Bull. Sci. Math.* **2002**, *126*, 343–367. [[CrossRef](#)]
23. Todorov, P.G. On the Faber polynomials of the univalent functions of class Σ . *J. Math. Anal. Appl.* **1991**, *162*, 268–276. [[CrossRef](#)]
24. Schiffer, M. Sur un problème extrémum de la représentation conforme. *Bull. Soc. Math. Fr.* **1938**, *66*, 48–55.
25. Duren, P.L. Coefficients of meromorphic schlicht functions. *Am. Math. Soc.* **1971**, *28*, 169–172. [[CrossRef](#)]
26. Janani, T.; Murugusundaramoorthy, G. Coefficient estimates of meromorphic bi-starlike functions of complex order. *Int. J. Anal. Appl.* **2014**, *4*, 68–77.
27. Orhan, H.; Magesh, N.; Balaji, V.K. Initial coefficient bounds for certain classes of Meromorphic bi-univalent functions. *Asian-Eur. J. Math.* **2014**, *7*, 1–9. [[CrossRef](#)]
28. Motamednezhad, A.; Salehian, S. Faber polynomial coefficient estimates for certain subclass of meromorphic bi-univalent functions. *Commun. Korean Math. Soc.* **2018**, *33*, 1229–1237. [[CrossRef](#)]
29. Salehian, S.; Zireh, A. Coefficient estimate for certain subclass of meromorphic and bi-univalent functions. *Commun. Korean Math. Soc.* **2017**, *32*, 389–397. [[CrossRef](#)]
30. Zireh, A.; Salehian, S. Initial coefficient bounds for certain class of meromorphic bi-univalent functions. *Acta Univ. Sapient. Math.* **2019**, *11*, 224–235. [[CrossRef](#)]

31. Srivastava, H.M.; Joshi, S.B.; Joshi, S.S.; Pawar, H. Coefficient estimates for certain subclasses of meromorphically bi-univalent functions. *Palest. J. Math.* **2016**, *5*, 250–258.
32. Hamidi, S.G.; Halim, S.A.; Jahangiri, J.M. Faber polynomials coefficient estimates for meromorphic bi-starlike functions. *Int. J. Math. Math. Sci.* **2013**, *2013*, 498159. [[CrossRef](#)]
33. Jahangiri, J.M.; Hamidi, S.G. Coefficients of meromorphic bi-Bazilevič functions. *J. Complex Anal.* **2014**, *2014*, 63917. [[CrossRef](#)]
34. Srivastava, H.M.; Motamednezhad, A.; Adegan, E.A. Faber polynomial coefficient estimates for bi-univalent functions defined by using differential subordination and a certain fractional derivative operator. *Mathematics* **2020**, *8*, 172. [[CrossRef](#)]
35. Srivastava, H.M.; Murugusundaramoorthy, G.; El-Deeb, S.M. Faber polynomial coefficient estimates of bi-close-to-convex functions connected with the Borel distribution of the Mittag-Leffler type. *J. Nonlinear Var. Anal.* **2021**, *5*, 103–118.
36. Bouali, A. Faber polynomials, Cayley-Hamilton equation and Newton symmetric functions. *Bull. Sci. Math.* **2006**, *130*, 49–70. [[CrossRef](#)]
37. Pommerenke, C. *Univalent Functions*, 1st ed.; Vandenhoeck und Ruprecht: Göttingen, Germany, 1975.
38. Hamidi, S.G.; Janani, T.; Murugusundaramoorthy, G.; Jahangiri, J.M. Coefficient estimates for certain classes of meromorphic bi-univalent functions. *C. R. Acad. Sci. Paris. Ser. I* **2014**, *352*, 277–282. [[CrossRef](#)]
39. Srivastava, H.M. Operators of basic (or q -) calculus and fractional q -calculus and their applications in geometric function theory of complex analysis. *Iran. J. Sci. Technol. Trans. A Sci.* **2020**, *44*, 327–344. [[CrossRef](#)]
40. Khan, B.; Srivastava, H.M.; Tahir, M.; Darus, H.; Ahmad, Q.Z.; Khan, N. Applications of a certain q -integral operator to the subclasses of analytic and bi-univalent functions. *AIMS Math.* **2021**, *6*, 1024–1039. [[CrossRef](#)]
41. Srivastava, H.M.; Altınkaya, S.; Yalcin, S. Hankel determinant for a subclass of bi-univalent functions defined by using a symmetric q -derivative operator. *Filomat* **2018**, *32*, 503–516. [[CrossRef](#)]
42. Srivastava, H.M.; El-Deeb, S.M. The Faber polynomial expansion method and the Taylor–Maclaurin coefficient estimates of bi-close-to-convex functions connected with the q -convolution. *AIMS Math.* **2020**, *5*, 7087–7106. [[CrossRef](#)]