

# Identifying signals of memory from observations of animal movements

Dongmin Kim, Peter R. Thompson, David W. Wolfson, Jerod A. Merkle, L. G. R. Oliveira-Santos, James D. Forester, Tal Avgar, Mark A. Lewis & John Fieberg

2024

Faculty of Science

Faculty Publications

© 2024 Kim et al. This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 license: <https://creativecommons.org/licenses/by/4.0/>

Original citation:

Kim, D., Thompson, P. R., Wolfson, D. W., Merkle, J. A., Oliveira-Santos, L. G. R., Forester, J. D., Avgar, T., Lewis, M. A., & Fieberg, J. (2024). Identifying signals of memory from observations of animal movements. *Movement Ecology*, 12(1). <https://doi.org/10.1186/s40462-024-00510-9>

---

Downloaded from UVicSpace Research & Learning Repository

dspace.library.uvic.ca



University  
of Victoria

Libraries

REVIEW

Open Access



# Identifying signals of memory from observations of animal movements

Dongmin Kim<sup>1,2\*</sup>, Peter R. Thompson<sup>3</sup>, David W. Wolfson<sup>4</sup>, Jerod A. Merkle<sup>5</sup>, L. G. R. Oliveira-Santos<sup>6</sup>, James D. Forester<sup>4</sup>, Tal Avgar<sup>7</sup>, Mark A. Lewis<sup>8</sup> and John Fieberg<sup>4</sup>

## Abstract

Incorporating memory (i.e., some notion of familiarity or experience with the landscape) into models of animal movement is a rising challenge in the field of movement ecology. The recent proliferation of new methods offers new opportunities to understand how memory influences movement. However, there are no clear guidelines for practitioners wishing to parameterize the effects of memory on moving animals. We review approaches for incorporating memory into step-selection analyses (SSAs), a frequently used movement modeling framework. Memory-informed SSAs can be constructed by including spatial-temporal covariates (or maps) that define some aspect of familiarity (e.g., whether, how often, or how long ago the animal visited different spatial locations) derived from long-term telemetry data. We demonstrate how various familiarity covariates can be included in SSAs using a series of coded examples in which we fit models to wildlife tracking data from a wide range of taxa. We discuss how these different approaches can be used to address questions related to whether and how animals use information from past experiences to inform their future movements. We also highlight challenges and decisions that the user must make when applying these methods to their tracking data. By reviewing different approaches and providing code templates for their implementation, we hope to inspire practitioners to investigate further the importance of memory in animal movements using wildlife tracking data.

**Keywords** Animal movement, Cognitive map, Familiarity, Memory, Occurrence distribution, Space use, Step-selection analyses, Wildlife tracking data

## \*Correspondence:

Dongmin Kim

kimx3725@umn.edu

<sup>1</sup> Department of Ecology, Evolution and Behavior, University of Minnesota, St. Paul, MN, USA

<sup>2</sup> Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA

<sup>3</sup> School of Environmental Science, Simon Fraser University, Burnaby, BC, Canada

<sup>4</sup> Department of Fisheries, Wildlife and Conservation Biology, University of Minnesota, St. Paul, MN, USA

<sup>5</sup> Department of Zoology and Physiology, University of Wyoming, Laramie, WY, USA

<sup>6</sup> Departamento de Ecologia, Universidade Federal do Mato Grosso do Sul, Campo Grande, Brazil

<sup>7</sup> Department of Biology, University of British Columbia—Okanagan and Wildlife Science Centre, Biodiversity Pathways Ltd, Kelowna, BC, Canada

<sup>8</sup> Department of Mathematics and Statistics and Department of Biology, University of Victoria, Victoria, BC, Canada



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## Introduction

Animal movement impacts ecological processes at all levels, including individual foraging efficiencies [16, 108], population persistence [24, 69], species distributions [38, 63], connectivity [26] and ecosystem functioning [12, 72, 99]. While much research has explored how animal movements are influenced by environmental conditions [38], intra- and inter-specific social interactions [63], and internal states such as hunger levels [49], the importance of past experience and memory is also increasingly recognized as a key component of animal movement [32, 60, 80]. For example, by remembering the location and outcomes of previously visited locations, many species can increase energy intake rates [50, 105] and lifespan [112], and avoid areas that might increase mortality risk [17, 43, 47]. Further, by remembering average environmental conditions, such as the average timing of resource waves, animals can better time migratory movements [2, 18]. Thus, models that incorporate memory are important for both developing and testing ecological theory, and they are likely to lead to improved predictions of how animals will respond to changes in their environment [33, 44].

One straightforward approach to integrating such complex types of memories into a model is to assume that past experiences can be encoded into a spatially-referenced system in the animal's brain (sometimes referred to as a "cognitive map"), which is then accessed during the retrieval phase to inform movements. Although hidden from direct observation, a spatially referenced map can be mathematically modeled as a surface changing dynamically over time as memories are lost, reinforced, or replaced. These constructs are central to key empirical models for memory, including "time since last visit" to a location [90] as a determinant of wolf movement [91], and episodic returns of brown bears to ephemeral seasonal resources [102, 103]. Overall, the map is a latent, spatially-referenced variable, whose dynamics are inferred indirectly from animal movement patterns. Although the existence of cognitive maps in animals remains a subject of ongoing debate among researchers [6, 111], we use the term "cognitive map" here to broadly describe neurological or psychological mechanisms that allow animals to store and process spatial information. This definition is commonly used in much of the movement ecology literature [32, 60].

The information that animals gather, and perhaps memorize, as they move can be divided into three categories: (1) spatial information (i.e., locations animals have visited), (2) site attributes, including resource quality or quantity, and (3) temporal information (i.e., about how long-ago animals visited a previous site or when a site peaks in forage quality) [32, 55, 60, 102, 103, 106]. Movement ecologists often distinguish between 'spatial

memory' that encodes spatial configurations (#1 from the description above) and 'attribute memory', which describes the attributes of local features (#2 and #3 from the description above) [32]. Food-caching blue jays (*Cyanocitta cristata*) use all three kinds of information: they remember the locations of many caches, the type of seed in each cache, and how long it has been since making the cache [23]. Owing to the temporal variability present in most environments, it can be advantageous to rely more heavily on recent experiences and to discount memories from long ago [25, 101]. For example, roe deer (*Capreolus capreolus*) primarily base their foraging decisions on recent experiences due to rapid changes in resource availability within their home ranges [85]. Note, however, that memory is also expected to temporally decay due to the limitations of the neurological infrastructure that holds it, and distinguishing such decay from an adaptive discounting may be particularly challenging [9]. By revisiting sites, animals can update their knowledge of site attributes, and optimal return times may depend on how quickly the reliability of past information decays due to environmental change as well as resource renewal rates [85, 98]. For example, wolves (*Canis lupus*) and brown bears (*Ursus arctos*) delay returning to previously visited kill sites so that prey numbers may recover [45, 92].

Ecologists have developed theoretical models to explore how past experiences and memory might influence animal movements [32, 39, 55, 60, 110]. The simplest encodable memory attribute is familiarity with a given location, either whether an individual has ever visited the site [75, 86] or how frequently it has visited the site in the past [17]. When coupled to spatial movement models, preference for familiar locations is sufficient for the formation of stable home ranges [42, 47, 66, 71, 84, 108]. More complex memory attributes include locations of resources or past conflicts, allowing animals to integrate spatial and attribute memory (i.e., memory of where positive and negative experiences occurred). Attraction to previously discovered resources can lead to resource-driven patterns of nonterritorial spatial segregation [1, 87]. By way of contrast, memory and avoidance of locations where past conflicts with conspecifics occurred can give rise to spontaneous territorial pattern formation [40, 82]).

Much of our understanding of the role of memory has come from ethologists and cognitive scientists studying animal behavior [53, 54, 95]. Early studies relied on observational data from experimental settings and found that food-storing birds frequently revisited sites to store and retrieve their resources (cache sites) [58, 59]. These behaviors may reflect the use of memory, but they may also be explained by birds preferring to store and revisit sites with certain characteristics [57].

To address this potential issue (which is still debated today, [79]), later ethologists used experimentation to separate out the role of memory-based and preference-based navigation by testing whether birds revisit their cache sites under different conditions (e.g., with and without food present) [94]. In addition, Balda and Kamil [11] forced food-caching Clark's nutcrackers (*Nucifraga columbiana*) to store food at non-preferred locations and found these locations were still revisited.

Recent advances in animal tracking technology and statistical modeling approaches have motivated ecologists to explore the potential for memory-informed movements in a wide range of animal taxa (although terrestrial mammals, mainly ungulates, remain by far the most-studied group; see Table 1). By tracking individual animals over consecutive years, ecologists can identify whether current movements can be explained from observations of previously visited locations. These studies face challenges similar to those faced by early ethologists, namely that animals may revisit sites primarily because those sites have characteristics the animals prefer [79]. To overcome this inferential challenge, ecologists fit statistical models that include covariates capturing habitat preferences along with covariates reflecting familiarity [65, 67]. This approach attempts to use statistical controls rather than experimental controls to infer whether animals revisit sites more frequently than expected based on site characteristics alone. However, we must be cautious about attributing revisits to memory, even after adjusting for known habitat preferences, because we will rarely know all the environmental features that influence animal movements [60]. Animals may be responding to unmeasured environmental cues that we do not include in our model.

Popular analytical frameworks, such as Step-Selection Analyses (SSAs) [10, 35, 38, 104] have been used to identify signals of memory from observations of animal movements [67, 76, 84, 86, 90]. We focus our review on SSAs because of their flexibility and ease of use due to readily available statistical software [96], but also because of their continuous methodological development [56, 68, 81, 97]. Nonetheless, several additional challenges remain before these approaches can be widely adopted. These include technological challenges associated with managing tracking data and creating models with different "familiarity" or "memory" covariates. Here, we provide an overview of methods for parameterizing memory effects in SSAs to help guide practitioners wishing to identify or quantify the effects of memory on animal movement. We also offer several examples with annotated code and then discuss the strengths and limitations

of current approaches and future directions for memory-informed movement research.

### Exploring how memory influences animal movements using SSAs

SSAs are widely used to quantify influences on animal movement [10, 35, 38, 104]. SSAs model movement in discrete time using two model components: (1) a selection-free movement kernel describing how animals move in the absence of habitat selection, defined using distributions of step length and turning angle, and (2) a selection function describing animal preferences concerning the habitat attributes at each step's endpoint (Box 1). Model parameters in SSAs can be estimated using commonly available statistical software that implements conditional logistic regression [10, 35]. Because they allow one to model and predict dynamic space-use patterns using accessible and available software, SSAs are attractive to movement ecologists and are widely used to analyze animal tracking data [96]. Further, SSAs have strong connections to other popular methods for modeling animal movement, they have been shown to be equivalent to biased correlated random walks [22], and they can be approximated by diffusion-taxis models [83]. In addition, certain continuous time movement models can be recast as SSAs [31].

A variety of spatiotemporal familiarity covariates can be included in an SSA to model the effects of memory (Boxes 2, 3). For example, one may estimate an occurrence distribution (OD; [5, 36]) describing the relative use of the landscape over a specific period in the past (e.g., a continuous surface of either the relative intensity of use or a binary presence/absence variable). By including the OD as a spatial predictor in an SSA, one can evaluate whether an animal's current landscape usage is biased toward or away from previously visited locations [76, 86, 117]. A notable limitation of this approach is the risk of overestimating the importance of familiarity due to unaccounted-for habitat attributes (e.g. if the OD reflects the distribution of an unobserved resource, see [79]). The OD approach further necessitates that users choose an appropriate time window in the past for calculating the OD (Fig. 1). Using a limited time window for calculating the familiarity covariate implies an abrupt memory decay function where past locations are memorialized for a fixed amount of time and then forgotten completely. Alternatively, one could choose to continually update the OD from the first to the last location, which would imply the animal never forgets its past experiences. With either approach, the OD effectively weights all previously visited areas within the specified time window equally, regardless of how long ago the animal visited the location. Another option is to allow more recent

**Table 1** Empirical modeling for memory

Paper	Species	Memory representation	Types	Methods	Model fit	Habitat-selection predictors
Aygar et al. [7]	Woodland caribou	Dynamics in animals' habitat selection over landscape and time	Forage	Likelihood of animals moving with cognitive selection functions	Bayesian (MCMC)	<ol style="list-style-type: none"> <li>1) Foraging qualities</li> <li>2) Predator density (Wolf)</li> <li>3) Moose habitat</li> <li>4) Snow depth</li> </ol>
Rheault et al. [86]	Mule deer	Differences between animals' occurrences throughout time	Forage	SSF framework with an OD as a response variable in the model	Step Selection Functions (SSFs) with conditional logistic regression	<ol style="list-style-type: none"> <li>1) Terrain ruggedness index</li> <li>2) elevation</li> <li>3) distance to treed edges</li> <li>4) snow depth</li> <li>5) land covers</li> <li>6) NDVI</li> <li>7) distance to well pads</li> <li>8) distance to roads</li> <li>9) previous ODs (past yr)</li> <li>10) current ODs (prev 7 days)</li> </ol>
Falcón-Cortés et al. [34]	Elk	Number of visits to the foraging path	Forage	A Markovian model with the probability of moving to patches	Bayesian (MCMC)	<p>memory decay &amp; memory use probability based on...</p> <ol style="list-style-type: none"> <li>1) distance between patches</li> <li>2) time since last visit to the patches</li> </ol>
Thompson et al [102, 103]	Brown Bear	Current resource quality + time since the last visit	Forage	Hidden Markov Model with SSF Framework Likelihood of animals moving in non-stationary states with a cognitive map	Maximum-likelihood estimates (MLE)	<ol style="list-style-type: none"> <li>1) distance from turbid water to riparian areas where bears selected</li> <li>2) vegetation classes</li> <li>3) density of ground squirrels and alpine sweet vetch</li> <li>4) distance from human settlements</li> </ol>
Oliveira-Santos et al. [76]	Feral hog	Utilization distribution of an animal's trajectory	Home range	SSF framework with an OD as a response variable in the model	Step Selection Functions (SSFs) with conditional logistic regression	<ol style="list-style-type: none"> <li>1) Land cover types</li> <li>2) Time of day</li> <li>3) BRB densities</li> <li>4) Residence time</li> </ol>
Northrup et al. [75]	Mule deer	Differences between animals' occurrences throughout time	Home range	SSF framework with an OD as a response variable in the model	Hierarchical Bayesian regression (MCMC)	<ol style="list-style-type: none"> <li>1) Tree</li> <li>2) density of drilling well pads</li> <li>3) average NDVI difference</li> <li>4) major road density</li> <li>5) terrain ruggedness</li> <li>6) fat</li> <li>7) age</li> <li>8) natural gas facilities density</li> <li>9) total snow depth difference</li> <li>10) pipeline density difference</li> </ol>
Schlägel et al. [91]	Wolf	Time since the last visit	Territorial defense	Spatially explicit random walk model similar framework of... + selection-free movement kernel + selection functions	Maximum-likelihood estimates (MLE)	<ol style="list-style-type: none"> <li>1) Prey density</li> <li>2) Distance to territory boundary</li> </ol>

**Table 1** (continued)

Paper	Species	Memory representation	Types	Methods	Model fit	Habitat-selection predictors
Ranc et al. [85]	Roe deer	Food availability	Home range	Likelihood of animals moving with cognitive selection functions	Maximum-likelihood estimates (MLE)	<ol style="list-style-type: none"> <li>1) Min. daily temperature</li> <li>2) within state resource access</li> <li>3) between-state resource access</li> <li>4) illumination index (dawn &amp; dusk)</li> <li>5) changes in the illumination index</li> </ol>
Ranc et al. [84]	Roe deer	Surrounding available habitats	Home range	Likelihood of animals moving with cognitive selection functions	Maximum-likelihood estimates (MLE)	<ol style="list-style-type: none"> <li>1) Landcover - reforested</li> <li>2) Landcover - agriculture</li> <li>3) step lengths</li> <li>4) Memory</li> <li>5) Memory decay</li> </ol>
Gurarie et al. [45]	Wolf	Predation success & Territorial marking	Territorial movement	A discrete choice modeling framework	Maximum-likelihood estimates (MLE)	<ol style="list-style-type: none"> <li>1) selected zone per each trip</li> <li>2) predation quality</li> <li>3) boundary coverage</li> <li>4) mass of prey item</li> <li>5) time since predation events</li> <li>6) number of kills</li> <li>7) boundary sizes</li> </ol>
Merkle et al. [67]	Mule deer	Differences between animals' occurrences throughout time	Migration	SSF framework with a distance and turning angle bias as parameters in the model	Step Selection Functions (SSFs) with conditional logistic regression	<ol style="list-style-type: none"> <li>1) Elevation</li> <li>2) % tree cover</li> <li>3) Distance to roads</li> <li>4) Terrain position index</li> <li>5) Integrated NDVI</li> <li>6) Rate of Green-up</li> <li>7) Distance to the previous route</li> <li>8) Direction to the previous range</li> </ol>

(or distant) memories to have more influence on current movements by replacing the OD with a covariate representing the length of time since the animal last visited a location (TSLV). Or, one can create multiple ODs reflecting space use during the recent or the more distant past and allow the model to determine optimal weights given to short-term and long-term memories represented by these covariates (e.g., [76]).

For migratory species that navigate relatively long distances, familiarity predictors could include distances between current and previous migratory trajectories or angular covariates that compare the direction of an animal's movement in relation to a previous seasonal range (Fig. 2). These methods can describe an animal's use of memory for navigation and capture its tendency to use familiar migration routes and consistent but seasonally varying home ranges [67].

Unless it is reasonable to assume that the animal lacked any memory at the onset of tracking, it is necessary to 'sacrifice' some early positional data to calculate the familiarity predictor. For some animals, including those that have long lifespans or live in highly seasonal environments, the 'memory build-up' period may need to be one year or more, unless there are reasons to believe memory (or its effect) decays over much shorter time scales. For example, Avgar et al. [7] used a full year of 'memory build-up' data to model woodland caribou (*Rangifer tarandus caribou*) space-use patterns over a subsequent year but have found no indication of memory decay within that time (indicating the need to use an even longer 'memory build up' period). Alternatively, studies of young or dispersing individuals and data from translocation experiments offer opportunities to model memory formation as individuals enter new and unfamiliar environments and learn how to navigate them efficiently [3, 15, 21, 32, 84, 85, 117]. However, datasets containing movements of individuals in unfamiliar landscapes are rare and difficult to obtain due to the high cost of translocation experiments and a tendency to avoid tracking juveniles due to often high mortality rates [93, 113].

## Case studies

In this section, we review memory-informed movement models for animal tracking data using 4 case studies. The first two examples, involving data from sandhill cranes (*Antigone canadensis*) and feral hogs (*Sus scrofa*), demonstrate how one can use a spatial familiarity predictor calculated from a past OD to explore whether animals retain information from their past experiences and for how long. The third example considers the migratory movements of mule deer and illustrates the use of two familiarity predictors formed using the minimum distance between the current and last year's migratory paths and the cosine of the angle between the direction of a current movement step and the previous year's centroid of locations. The last example, involving data from a brown bear, uses a spatiotemporal covariate to quantify how long it has been since the individual last visited spatial locations, and thus, how memory may influence revisitation rates. We provide a workflow to reproduce the main components of the habitat- and memory-based SSF from all the case studies, demonstrating the relative importance of memory-based and habitat metrics in an SSF of animal movement (see Appendix codes for the case studies). We also highlight challenges and decisions that the user must make when applying these methods to their tracking data (Box 2).

### Sandhill crane—'fixed-time' OD

Sandhill cranes breed throughout North America during the summer and migrate south for the winter. During their first year, juveniles migrate to overwintering areas with their parents and then disperse from the family group either during the spring migration or upon arrival to the natal territory the following spring [46, 100]. During the first few years of independence, subadult cranes typically make long-distance movements across the landscape during the summer; in contrast, the movements of breeding adults are largely constrained to their breeding territories [119]. Once cranes become successful breeding adults, typically between 4 and 6 years old, they use

## Box 1 Step-Selection Analyses (SSAs)

Step-Selection Analyses (SSAs) model the conditional probability,  $p(s_t | H_{t-1}; \beta_m, \beta_w)$ , of finding an individual at a location  $s_t$  at the time  $t$  given a set of previously visited locations,  $H_{t-1}$ , using a selection-free movement kernel,  $k(s_t | H_{t-1}; \beta_m)$ , which describes how animals would move in the absence of habitat selection, and a movement-free habitat-selection function,  $w(s_t; t, \beta_w)$ , which describes the animals' preferences for certain environmental features (e.g., variables representing resources, risks, and/or other conditions; [62]):

$$p(s_t | H_{t-1}; \beta_m, \beta_w) = \frac{k(s_t | H_{t-1}; \beta_m) \cdot w(s_t; t, \beta_w)}{\int_{s' \in U} k(s' | H_{t-1}; \beta_m) \cdot w(s'; t, \beta_w) ds'} \quad (1)$$

$$H_{t-1} = s_{t-1}, s_{t-2}, \dots, s_{t-j}, \dots, s_0 \quad (2)$$

$\beta_m$  contains parameters in the step-length and turn angle distributions ( $\beta_{m1}, \dots, \beta_{mq}$ ), and  $\beta_w$  contains resource-selection parameters that quantify the attractiveness of different locations using a vector of selection coefficients ( $\beta_{w1}, \dots, \beta_{wp}$ ) for each environmental covariate ( $r_1(s_t), \dots, r_p(s_t)$ ).  $s_t \in U$  describes all the locations within the spatial domain  $U$ . To calculate a step length,  $s_t$ , two locations are required,  $(s_t, s_{t-1})$ . Similarly, a turning angle,  $ta_t$ , is calculated using the current and the past two locations  $(s_t, s_{t-1}, s_{t-2})$ .

**Box 2** Guide for the memory-informed movement modeling within SSF framework


Question	Biological example	Method	Caveats, challenges, limitations	Implementation parameters or choices
Are animals more likely to visit areas that are more familiar to them?	Feral hogs primarily rely on recent experiences when making night time foraging decisions [76].	Occurrence Distribution (OD)	The effects of unmodeled environmental drivers may be wrongly attributed to memory and vice versa [79].	Time window for measuring past space use. Length of burn-in periods (how much early positional data to sacrifice to estimate the familiarity predictor?)
Do animals prefer to visit areas they have not visited recently? (and if so, is there an optimal return time based on foraging or predation experience?)	Predators like wolves and brown bears delay returning to previously visited kill sites so that prey numbers may rebound to normal levels [45, 92].	Time Since Last Visit (TSLV)	The effects of unmodeled environmental drivers may be wrongly attributed to memory and vice versa [79]. TSLV may be relatively homogeneous within an animal's movement kernel.	"Patch" size (what is the typical spatial grain of the animal's cognitive map?) Length of burn-in periods (how much early positional data to sacrifice to estimate the familiarity predictor?)
Do migratory animals use consistent migration routes to return to the same seasonal ranges each year?	Mule deer ( <i>Odocoileus hemionus</i> ) have been shown to use the exact same migration routes and seasonal ranges, likely relying on their memory of those spatial locations to help them navigate [67].	Distance and directional bias toward previously used areas across large spatial scales	The effects of unmodeled environmental drivers and social cues may be wrongly attributed to memory and vice versa [79].	Whether to use angular covariates or distance-to covariates. Time window for measuring past space use.

**Box 3** Familiarity function in SSFs

Movement ecologists have described familiarity with different parts of the landscape that animals experienced using familiarity covariates, which we formalize via a familiarity function,  $f$ . Including an exponential familiarity function allows the attractiveness of different locations to be governed by both environmental and familiarity covariates within the traditional SSF framework ( $e^{\beta_w r(s)} \cdot e^{\beta_f f(s)} = e^{\beta_w r(s) + \beta_f f(s)}$ ). Examples of familiarity covariates include occurrence distributions (ODs) reflecting the intensity of past space use in the study area, time since last visit (TSLV), migratory distances between current and previously used paths, and angular covariates used to capture bias toward previously used migratory ranges:

$$p(s_t | H_{t-1}; \beta_m, \beta_w, \beta_f) = \frac{k(s_t | H_{t-1}; \beta_m) \cdot w(s_t; t, \beta_w) \cdot f(s_t; t, \beta_f)}{\sum_{s' \in U} k(s' | H_{t-1}; \beta_m) \cdot w(s'; t, \beta_w) \cdot f(s'; t, \beta_f)} \tag{3}$$

$$f(s_t; t, \beta_f) = \{e^{\beta_f \cdot f(s_t)}\} \tag{4}$$



**Fig. 1** Spatial memory can be quantified using an animal’s occurrence distribution (OD) measured over some prior period (colored areas: 5-day periods at time 6 and time 8 = orange vs green) which captures an animal’s movement path and its uncertainty. A time-varying covariate can be constructed by updating the OD at regular time intervals. This updating step ensures that distant experiences are eventually forgotten and no longer play a role in driving animal movement. Users must choose an appropriate time window in the past for calculating the memory covariate and how often to update it


their accumulated knowledge of the landscape to return and nest in the same breeding area in subsequent years [74, 100].

As an illustrative example of how a sandhill crane’s space use constricts each year as the crane learns its landscape and develops a breeding territory, we consider a 5-year dataset of global positioning system (GPS) telemetry locations of a sandhill crane (with a 15-minute fix interval), starting from the time of fledging [118, 119]. A visualization of summer locations shows that the spatial coverage visited by the crane decreases each year as it ages, selecting locations it had previously visited (Fig. 3a). This pattern suggests that the crane may be using its past experience to decide where to establish a breeding territory.

To quantify the potential effects of familiarity on the crane’s summer movements, we calculated areas associated with the 95% contour of an OD estimated from the previous year’s summer locations using conditional probability density functions from a continuous-time movement model (where the conditioning ensures movements pass through observed locations; [36]). We then

fit a separate SSF to each year of data, including the previous year’s OD as a predictor in the model. A positive coefficient associated with the previous year’s OD would suggest the crane prefers to revisit sites it visited in the previous year (compared to *equally accessible and otherwise identical sites* that it did not visit during the previous year, where accessibility is determined by the selection-free movement kernel; [35]). If the coefficient is negative, the opposite interpretation holds.

Coefficients associated with the previous year’s OD were positive in all years (Fig. 3b), even though the crane rarely revisited previously used locations in the first two years (Fig. 3c). Interpretation of the coefficients associated with categorical predictors can be difficult, as they reflect a ratio of ratios (use: availability ratio for one class versus the use: availability ratio of a reference class; [35]). The positive coefficients for the previous year’s OD in 2016 and 2017 reflect the fact that the use: availability ratio associated with grid cells visited in the previous year was higher than the use: availability ratio for grid cells that were not visited in the previous year. Thus, we end up with a positive coefficient for the OD predictor even



**Fig. 2** Familiarity covariates used in applications involving migratory animals. The red bounding box in panel (a) displays the area zoomed into for boxes b and c. Memory can be quantified using a distance predictor calculated as the minimum distance between current and past migratory paths (panel b). Specifically, we can calculate the distances (dashed lines) between Year 2 (observed [navy], random [pink]) locations and Year 1 (green) migratory paths. Memory can also be included as a directional bias predictor by comparing whether the current year's steps are biased toward the previous season's range (panel c). This bias predictor can be calculated using the angles,  $\theta_1, \theta_2$ , between the step (observed [navy], random [pink]) and the centroid of the previous year's winter range

though most of the areas encountered and used by the crane represent areas that were not visited in the prior year. Recall our interpretation of a positive coefficient, *if presented with two equally accessible locations* (same step length and turn angle required to reach both locations) that only differ in whether they had been visited in the previous year (i.e., they had the same landcover class), the crane would be more likely to select the location it was familiar with. A limitation of this model and application is that the crane was rarely presented with this type of choice. In the first two years, it was rarely found near sites it had previously visited whereas in years 3–5 it rarely visited areas it was unfamiliar with. This example highlights a limitation of the OD approach and the need for new methods that can capture tradeoffs in exploratory and informed movements as young individuals learn to navigate the landscape. Another limitation is that the birds may be selecting for an unobserved resource, one that we did not include in our analysis, which introduces

a hidden correlation between their past and current space use.


**Feral Hog – OD with temporal variation: short- and long-term memories stratified by time of day**

Hogs were introduced to the Pantanal wetland about 300 years ago, and currently represent the highest wild mammal biomass in this region. They are crepuscular-nocturnal, social, long-lived, cooperative animals that forage at the edges of water bodies and in ephemeral pools that become increasingly rare during the winter dry season. Hogs lack sweat glands and behaviorally mitigate heat stress by spending the hot hours of the day resting in forest patches.

As animals move, they can access and update their reference memory (long-term acquisition) and working memory (short-term acquisition) to navigate through space. Still, the stored spatial information may or may not be used depending on the current animal needs and

(See figure on next page.)

**Fig. 3** Visualization of Sandhill crane tracking data and coefficients from fitted step-selection functions with a memory covariate formed using an OD capturing previously visited locations. **a** Movement patterns of an individual sandhill crane during summer seasons (06/15-09/22) from 2015 to 2020. **b** Parameter estimates including those for the memory covariate (Odprev: OD from the previous year). **c** Distribution of each year's used and available locations for sites (i.e., grid cells) that were (OD = 1) and were not (OD = 0) visited in the previous year. Numbers at the top of the bars indicate the number of locations in each group



**Fig. 3** (See legend on previous page.)

context, which generates temporal heterogeneity (e.g., within the day, and seasons) in the use of that information. Oliveira-Santos et al. [76] modeled the hidden process underlying spatial memory acquisition using spatiotemporal covariates generated from Biased Random Bridge kernel density estimates based on residence time [13]. Specifically, for each individual step, they used previous locations to build multiple spatial maps, formed using different time windows, that could affect the hogs' future movement decisions. These maps were then continuously updated as the individual moved through the landscape.

Oliveira-Santos et al. [76] considered 4 conceptually different hypotheses regarding how hogs process and use spatial information, combining long- and short-term memory with differential use of stored memory within the daily cycle. To represent long-term memory, familiarity covariates were constructed and constantly updated with all previous locations as the individual moved, whereas short-term memory was represented using a spatiotemporal covariate that kept track of just the last 3 days. Additionally, long- and short-term familiarity covariates were also built considering only daytime or only nighttime locations, which they referred to as long-term temporal and recent-temporal memory, respectively. When fitting models for these last two cases, movement steps taken at night or during the day were paired with familiarity covariates generated from previous locations collected only at night or day, respectively.

All tracked hogs strongly selected for previously visited areas, mainly those associated with short-term memory. Most of these individuals (65% of the tracked hogs) appeared to use working memory as part of their movement process, as covariates generated from recent nocturnal locations were better at predicting future nocturnal use than covariates generated from all time periods. Importantly, the effect of familiarity also varied within the day, being more important during the daylight hours when individuals were sleeping in well-known places than at night when animals were foraging and were more willing to take risks by walking through less familiar areas. Although hogs are acknowledged to have high cognitive skills and memory retention, Oliveira-Santos et al. [76] concluded that they relied mainly on recent spatial information because the distribution of prime food resources in the study area responds quickly to foraging pressure and changes in water levels.

#### **Mule Deer—migratory paths and angles**

Many large ungulates are migratory, capitalizing on seasonal and spatial variation in food, predation, and hospitable conditions [9]. Mule deer are a concentrate forager (i.e., prefer to consume high-quality food) that

display some of the longest terrestrial migrations in North America [51], spending winters in arid, low-elevation sagebrush, grassland, and desert ecosystems, and then migrating (up to 400 km) into montane ecosystems at higher elevations for summer.

Unlike some other migratory species (e.g., Sierra Nevada Bighorn Sheep; [14]), Mule deer migrate well outside their perceptual range (e.g., what they see, hear, and smell at a given moment), yet they display relatively strong fidelity to seasonal ranges [70] and migration routes [88]. On average, mule deer migrate on the same path during spring year after year 81% of the time [88]. In some cases, such migrations occur across relatively vast expanses of flat deserts, rolling hills, and thick forests, where sensory abilities such as vision may provide limited cues for navigation [89].

Evidence suggests that mule deer may navigate during migration by memorizing the path of their previous migration route and the general location of their seasonal ranges [67, 107]. The relative influence of these memorized spatial locations can be assessed in a movement model that first considers several other habitat features that may influence ungulate movement and space use. Merkle et al. [67] examined the relative role of memory usage versus local variation in habitat on mule deer navigation during migration. They found that variables indexing past experience (distance to the previous migration route and direction to the previous seasonal range) were 2–28 times more predictive of migratory movements than local variation in habitat. Alternative explanations for these long-distance migrations include following scent trails or other conspecifics; however, those explanations are not well supported due to the fact that these deer spend on average 81% of their migration walking on the exact same path as in the previous year [88].

#### **Brown Bear—time since last visit (TSLV)**

Brown bears are opportunistic omnivores found in North America, Europe, and Asia [78]. Their life history strategies and dietary compositions vary greatly depending on the environment they live in [37, 41, 61], but their ability to navigate towards previously visited food patches is ubiquitous throughout their natural range [92, 102, 103, 116]. The “barren-ground grizzly bears” found in the Canadian Arctic are unique for many reasons, including their foraging and denning behavior [30, 64]. These bears' food resources are only available for a short, albeit predictable, portion of their active seasons [30], so in addition to returning to the correct spatial location where food was present, bears must identify the temporal pattern of this resource and revisit the patch at the correct time. Ecologists interested in understanding how these bears incorporate memory into their movement patterns


must implement models that account for these nonlinear temporal dynamics.

We modified the model developed by Schlägel and Lewis [90] so that it could be implemented in the step-selection framework and fit using conditional logistic regression. This required using distributions from the exponential family (gamma, von Mises) to model the distribution of step lengths and turn angles, respectively [10, 35]. In addition, we included both linear and quadratic terms to model the influence of time since last visit (TSLV) on the movement patterns of an adult female brown bear in the Mackenzie River Delta region of the Northwest Territories, Canada. This bear was immobilized and fitted with a GPS collar that recorded its location every 4h during the active season (the time in which the bear was out of its den).

We calculated TSLV as the difference between the current timestamp and the last time the bear visited a series of 2×2 km grid cells, updating the map at each observation time. This familiarity covariate allowed “revisitation” to occur when the animal was in the perceptual vicinity of an area it previously visited (i.e., within the same grid cell), without necessitating that the animal returns exactly to its previous coordinates [90]. We discarded the first year of location data as a “burn-in,” and set TSLV to 365 for any grid cells where TSLV was missing, effectively assuming that these cells had been visited just prior to

the first observed location. The need for a burn-in period is a limitation of this approach but is necessary since we have no history of the bear’s past visits prior to the start of data collection. A sensitivity analysis can be performed to evaluate whether results change if the length of the burn-in period is increased or decreased. To analyze how selection strength varied with TSLV, we calculated the relative selection strength (RSS) at different TSLV values [10]. As with other habitat selection models, by including both linear and quadratic terms, we were able to identify a non-linear response to TSLV with intermediate TSLV values of approximately 350 days displaying the strongest selection (Fig. 4). These results agree with those of Thompson et al. [103], who also found that bears tended to revisit sites seasonally.

Although the TSLV approach can capture the potential memory effects of wildlife over time, the interpretation of the model may depend on the user’s choice of resource covariates. For example, we used berries as a seasonal resource covariate. However, the importance of the memory covariate (TSLV) might change if additional environmental covariates are added to the model [79]. Similar to Merkle et al. [67], Thompson et al. [103] compared their models with memory to “resource-only” models without memory, finding that the former models better fit the data. Yet, it is nearly impossible to perfectly quantify the distribution of food resources on the landscape, and



**Fig. 4** Visualization of the relative attractiveness of previously visited locations as a function of time since last visit (TSLV) modeled using linear and quadratic terms in a step-selection analysis of brown bear location data with the distribution of used (green, bottom) and available (brown, top) locations shown along the x-axis

“resource-only” models might have performed better if more environmental data could be acquired. Even when alternative models are considered and compared rigorously to a “memory model”, it is important to consider how these alternative models may fall short in quantifying their desired hypotheses.

## Discussion

Modern GPS tracking systems generate massive telemetry datasets by following individual animals over a long time on a global scale. With this abundance of available data, it is now possible to develop models that evaluate how memory relates to animal movement [73], which has inspired the recent development and application of many such models (see Supporting Information 1). Our review focused on approaches that account for familiarity with different areas of the landscape by including spatial, spatiotemporal, or angular covariates as predictors in step-selection analyses. These frameworks can be implemented using available statistical software for fitting conditional logistic regression models, which we demonstrate using multiple tracking data sets and annotated code examples. Although some data development is necessary before fitting the models (e.g., to create the memory predictors and generate random steps), users can leverage R packages to make these steps easier. For example, the *ctmm* package [19] can be used to calculate ODs, and the *amt* package [96] can be used to generate random steps necessary for parameterizing the model.

Individual-based models of animal movement are increasingly used to inform conservation and management at the population or even species levels [4, 48, 114]. Prime among these applications is the use of habitat-selection models to identify critical habitats, delineate range boundaries, and project spatial distribution across space or time [77]. A strong selection for certain habitats or environmental features and conditions may occur with or without memory, and it is important to consider how models that incorporate familiarity covariates may alter inferences. On the one hand, we might expect mild to moderate collinearity between familiarity covariates and other important environmental drivers, which can make it challenging to quantify their unique contributions [79]. On the other hand, accounting for familiarity and memory effects should reduce bias associated with estimators of habitat-selection strength. Consider an animal that is both attracted to certain habitats and to places it has used in the past; if we ignore the latter, we will consequently overestimate the former, putting more weight on habitat attraction than it truly has [79]. Similarly, it is important to consider the effect of environmental drivers when looking for influences of memory on animal movements [109]. Memory-free movement models that allow

for habitat selection have been suggested as null models for evaluating evidence of memory [79]. An important caveat, however, is that it will be difficult in most cases to identify and measure all environmental features that animals select for. Thus, movement ecologists need to be cautious with attributing the effects of familiarity covariates to memory, even when the inclusion of familiarity covariates improves model fit relative to a null model containing known predictors of habitat quality. An additional benefit to considering memory effects in SSAs is the option to simulate these effects under various management scenarios [39]. For example, managers might be interested in assessing the likelihood of successfully relocating an animal and having it establish a home range in a new site [14, 85]. Explicitly modeling the process of building and responding to increasing knowledge of the landscape may be critical for obtaining reliable predictions.

Information gathering, an essential prerequisite for memory, is governed by sensory ecology [20, 27–29, 52], and it is in this context that we must also consider the limitations associated with the use of SSA to infer memory. The information available to the animal about any given spatial location (the ‘signal’) is a function of the animal’s position in relation to that location (the source of the signal), the strength of the signal (e.g., the intensity of odor, light, or sound), the overlap with similar signals coming from elsewhere on the landscape, and the animal’s sensory capacity to perceive and process the signal (which in itself may be a complex function of the animal’s morphology and physiology). Information can only be committed into memory if it is being perceived. Moreover, it is very likely that the weight given to memorized information (or its retention time) is modulated by the signal-to-noise ratio at which the information was perceived, and perhaps even by the desirability (‘good’ or ‘bad’) of the information (valence-dependent learning,[8]). The models we described here assume, for the most part, that animals sense and retain information from only ‘visited localities’, which are arbitrarily sized spatial units (typically corresponding to a single pixel of the available environmental data). Furthermore, most SSFs only assess what the animal can perceive within local ranges (i.e., what is nearby), though angular and distance-to-covariates allow modeling perception over larger spatial scales. Thus, SSFs are a useful, but extreme simplification of the true underlying sensory ecology.

More mechanistically inclined frameworks for explicitly modeling the sensory processes involved in memory buildup have been proposed (e.g., [7, 9, 65, 84, 102, 103]); however, fitting these models to wildlife tracking data is still a challenge. These frameworks typically include additional free parameters used to construct spatiotemporal

covariates that determine how perceived habitat quality varies over time (e.g., by modeling how perception decays with distance from the individual, and memory decays temporally). As a result, it is not possible to fit these models using standard statistical software developed for SSAs. Instead, the parameters governing the spatio-temporal covariate must be estimated simultaneously with other habitat-selection and movement parameters through a custom-written likelihood function that can be optimized using Markov chain Monte Carlo or other numerical optimization methods. We include an example from Thompson et al. [103] in our supplementary material to demonstrate this approach. Methods that more realistically model the process of memory formation should be pursued, but we suspect that most practitioners will continue to explore the role of memory on animal movements using simpler models that can fit within the standard SSA framework using conditional logistic regression. The strength of this approach is that it can be easily and widely applied to tracking data. Still, these efforts should be complemented by additional experiments and more realistic mechanistic models to better understand the multifaceted ways that animals use memory to navigate their landscapes [84, 115].

### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40462-024-00510-9>.

Supplementary Material 1.

### Author contributions

DK and JF conceived of the topic, and DK led the writing of the manuscript. Case studies were led by DK, DW, and JF (sandhill crane), DK, PT, and JF (brown bear), JM (mule deer), and L O-S and JDF (hog). All authors discussed, edited, read, and approved the final manuscript.

### Funding

DK was supported by the National Science Foundation INTERN award 1654609, the Smithsonian Institution Fellowship Program (SIFP), and the University of Minnesota Doctoral Dissertation Fellowship (DDF). JF was supported by the National Aeronautics and Space Administration award 80NSSC21K1182. JF and JDF received partial salary support from the Minnesota Agricultural Experimental Station.

### Availability of data and materials

Data is provided within the supplementary information files and the datasets and code associated with the case studies can be found in a Github repository [https://github.com/kimx3725/Memory\\_Movement/tree/main](https://github.com/kimx3725/Memory_Movement/tree/main)

### Declarations

#### Ethics approval and consent to participate

Not applicable.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare no competing interests.

Received: 23 May 2024 Accepted: 3 October 2024

Published online: 18 November 2024

### References

- Aarts G, Mul E, Fieberg J, Brasseur S, van Gils JA, Matthiopoulos J, Riotte-Lambert L. Individual-level memory is sufficient to create spatial segregation among neighboring colonies of central place foragers. *Am Nat.* 2021;198(2):E37–52.
- Abrahms B, Hazen EL, Aikens EO, Savoca MS, Goldbogen JA, Bograd SJ, Jacox MG, Irvine LM, Palacios DM, Mate BR. Memory and resource tracking drive blue whale migrations. *Proc Natl Acad Sci.* 2019;116(12):5582–7.
- Aikens EO, Nourani E, Fiedler W, Wikelski M, Flack A. Learning shapes the development of migratory behavior. *Proc Natl Acad Sci.* 2024;121(12): e2306389121.
- Aiello CM, Galloway NL, Prentice PR, Darby NW, Hughson D, Epps CW. Movement models and simulation reveal highway impacts and mitigation opportunities for a metapopulation-distributed species. *Landscape Ecol.* 2023;38(4):1085–103.
- Alston JM, Fleming CH, Noonan MJ, Tucker MA, Silva J, Folta C, Akre TS, Ali AH, Belant JL, Beyer D, Blaum N. Clarifying space use concepts in ecology: range vs. occurrence distributions. *BioRxiv.* 2022;3:155.
- Asensio N, Brockelman WY, Malaivijitnond S, Reichard UH. Gibbon travel paths are goal oriented. *Animal Cogn.* 2011;14:395–405.
- Avgar T, Baker JA, Brown GS, Hagens JS, Kittle AM, Mallon EE, McGreer MT, Mosser A, Newmaster SG, Patterson BR, Reid DE. Space-use behaviour of woodland caribou based on a cognitive movement model. *J Animal Ecol.* 2015;84(4):1059–70.
- Avgar T, Berger-Tal O. Biased learning as a simple adaptive foraging mechanism. *Front Ecol Evol.* 2022;8(9): 759133.
- Avgar T, Deardon R, Fryxell JM. An empirically parameterized individual based model of animal movement, perception, and memory. *Ecol Model.* 2013;251:158–72.
- Avgar T, Potts JR, Lewis MA, Boyce MS. Integrated step selection analysis: bridging the gap between resource selection and animal movement. *Method Ecol Evol.* 2016;7(5):619–30.
- Balda RP, Kamil AC. Long-term spatial memory in Clark's nutcracker *Nucifraga columbiana*. *Animal Behav.* 1992;44(4):761–9.
- Bauer S, Hoyer BJ. Migratory animals couple biodiversity and ecosystem functioning worldwide. *Science.* 2014;344(6179):1242552.
- Benhamou S. Dynamic approach to space and habitat use based on biased random bridges. *PLoS ONE.* 2011;6(1): e14592.
- Berger DJ, German DW, John C, Hart R, Stephenson TR, Avgar T. Seeing is be-leaving: perception informs migratory decisions in Sierra Nevada bighorn sheep (*Ovis canadensis sierrae*). *Front Ecol Evol.* 2022;10: 742275.
- Brønsvik H, Nourani E, Fiedler W, Flack A. Experience reduces route selection for conspecifics by the collectively migrating white stork. *Current Biol.* 2024;34(9):2030–7.
- Bonte D, Van Dyck H, Bullock JM, Coulon A, Delgado M, Gibbs M, Lehouck V, Matthysen E, Mustin K, Saastamoinen M, Schtickzelle N. Costs of dispersal. *Biol Rev.* 2012;87(2):290–312.
- Bracis C, Bildstein KL, Mueller T. Revisitation analysis uncovers spatio-temporal patterns in animal movement data. *Ecography.* 2018;41(11):1801–11.
- Bracis C, Mueller T. Memory, not just perception, plays an important role in terrestrial mammalian migration. *Proc Royal Soc B Biol Sci.* 1855;284(1855):20170449.
- Calabrese JM, Fleming CH, Gurarie E. ctmm: an R package for analyzing animal relocation data as a continuous-time stochastic process. *Method Ecol Evol.* 2016;7(9):1124–32.
- Dall SR, Giraldeau LA, Olsson O, McNamara JM, Stephens DW. Information and its use by animals in evolutionary ecology. *Trends Ecol Evol.* 2005;20(4):187–93.
- Dalziel BD, Morales JM, Fryxell JM. Fitting probability distributions to animal movement trajectories: using artificial neural networks to link distance, resources, and memory. *Am Nat.* 2008;172(2):248–58.

22. Duchesne T, Fortin D, Rivest LP. Equivalence between step selection functions and biased correlated random walks for statistical inference on animal movement. *PLoS ONE*. 2015;10(4): e0122947.
23. Darley-Hill S, Johnson WC. Acorn dispersal by the blue jay (*Cyanocitta cristata*). *Oecologia*. 1981;50:231–2.
24. Delgado Sánchez MM, Miranda M, Álvarez SJ, Gurarie E, Fagan WF, Penneriani Dragone V, di Virgilio A, Morales JM. The importance of individual variation in the dynamics of animal collective movements. *Philos Trans Royal Soc Biol Sci*. 2018;373:20170008.
25. Devenport LD. Spontaneous recovery without interference: why remembering is adaptive. *Animal Learn Behav*. 1998;26(2):172–81.
26. Doherty TS, Driscoll DA. Coupling movement and landscape ecology for animal conservation in production landscapes. *Proc Royal Soc B Biol Sci*. 1870;2018(285):20172272.
27. Dominoni DM, Halfwerk W, Baird E, Buxton RT, Fernández-Juricic E, Fristrup KM, McKenna BF, Mennitt DJ, Perkin EK, Seymoure BM, Stoner DC. Why conservation biology can benefit from sensory ecology. *Nat Ecol Evol*. 2020;4(4):502–11.
28. Dukas R, editor. *Cognitive ecology: the evolutionary ecology of information processing and decision making*. University of Chicago Press; 1998.
29. Dusenbery DB. *Sensory ecology: how organisms acquire and respond to information*. New York: WH Freeman; 1992.
30. Edwards MA, Derocher AE. Mating-related behaviour of grizzly bears inhabiting marginal habitat at the periphery of their North American range. *Behav Process*. 2015;111:75–83.
31. Eisaguirre JM, Williams PJ, Hooten MB. Rayleigh step-selection functions and connections to continuous-time mechanistic movement models. *Mov Ecol*. 2024;12(1):14.
32. Fagan WF, Lewis MA, Auger-Méthé M, Avgar T, Benhamou S, Breed G, LaDage L, Schlägel UE, Tang WW, Papastamatiou YP, Forester J. Spatial memory and animal movement. *Ecol Lett*. 2013;16(10):1316–29.
33. Fagan WF, McBride F, Korolov L. Reinforced diffusions as models of memory-mediated animal movement. *J Royal Soc Interf*. 2023;20(200):20220700.
34. Falcón-Cortés A, Boyer D, Merrill E, Frair JL, Morales JM. Hierarchical, memory-based movement models for translocated elk (*Cervus canadensis*). *Frontiers in Ecology and Evolution*. 2021;9:702925.
35. Fieberg J, Signer J, Smith B, Avgar T. A 'how to' guide for interpreting parameters in habitat-selection analyses. *J Animal Ecol*. 2021;90(5):1027–43.
36. Fleming CH, Fagan WF, Mueller T, Olson KA, Leimgruber P, Calabrese JM. Rigorous home range estimation with movement data: a new autocorrelated kernel density estimator. *Ecology*. 2015;96(5):1182–8.
37. Finnegan SP, Svoboda NJ, Fowler NL, Schooler SL, Belant JL. Variable intraspecific space use supports optimality in an apex predator. *Sci Rep*. 2021;11(1):21115.
38. Fortin D, Beyer HL, Boyce MS, Smith DW, Duchesne T, Mao JS. Wolves influence elk movements: behavior shapes a trophic cascade in Yellowstone National Park. *Ecology*. 2005;86(5):1320–30.
39. Forrest SW, Pagendam D, Bode M, Drovandi C, Potts JR, Perry J, Vanderduys E, Hoskins AJ. Simulating animal movement trajectories from temporally dynamic step selection functions. *bioRxiv*. 2024:2024-03.
40. Ellison N, Hatchwell BJ, Biddiscombe SJ, Napper CJ, Potts JR. Mechanistic home range analysis reveals drivers of space use patterns for a non-territorial passerine. *J Animal Ecol*. 2020;89(12):2763–76.
41. García-Rodríguez A, Selva N, Zwiłjac-Kozica T, Albrecht J, Lionnet C, Rioux D, Taberlet P, De Barba M. The bear-berry connection: ecological and management implications of brown bears' food habits in a highly touristic protected area. *Biol Conserv*. 2021;264: 109376.
42. Gaustestad AO, Loe LE, Mysterud A. Inferring spatial memory and spatiotemporal scaling from GPS data: comparing red deer *Cervus elaphus* movements with simulation models. *J Animal Ecol*. 2013;82(3):572–86.
43. Gehr B, Bonnot NC, Heurich M, Cagnacci F, Ciuti S, Hewison AM, Gailard JM, Ranc N, Premier J, Vogt K, Hofer E. Stay home, stay safe—site familiarity reduces predation risk in a large herbivore in two contrasting study sites. *J Animal Ecol*. 2020;89(6):1329–39.
44. Gurarie E, Avgar T. Cognitive movement ecology. *Front Ecol Evol*. 2024;12:1360427.
45. Gurarie E, Bracis C, Brilliantova A, Kojola I, Suutarinen J, Ovaskainen O, Potluri S, Fagan WF. Spatial memory drives foraging strategies of wolves, but in highly individual ways. *Front Ecol Evol*. 2022;10: 768478.
46. Hayes MA. *Dispersal and population genetic structure in two flyways of sandhill cranes (*Grus canadensis*)* (Doctoral dissertation, The University of Wisconsin-Madison). 2015
47. Heathcote RJ, Whiteside MA, Beardsworth CE, Van Horik JO, Laker PR, Toledo S, Orchan Y, Nathan R, Madden JR. Spatial memory predicts home range size and predation risk in pheasants. *Nat Ecol Evol*. 2023;7(3):461–71.
48. Hofmann DD, Cozzi G, McNutt JW, Ozgul A, Behr DM. A three-step approach for assessing landscape connectivity via simulated dispersal: African wild dog case study. *Landsc Ecol*. 2023;38(4):981–98.
49. Hooten MB, Scharf HR, Morales JM. Running on empty: recharge dynamics from animal movement data. *Ecol Lett*. 2019;22(2):377–89.
50. Janson C, Byrne RW. Resource cognition in wild primates: opening up the black box. *Animal Cogn*. 2007;10:357–67.
51. Joly K, Gurarie E, Sorum MS, Kaczensky P, Cameron MD, Jakes AF, Borg BL, Nandintsetseg D, Hopcraft JG, Buuveibaatar B, Jones PF. Longest terrestrial migrations and movements around the world. *Sci Rep*. 2019;9(1):15333.
52. Jordan LA, Ryan MJ. The sensory ecology of adaptive landscapes. *Biol Lett*. 2015;11(5):20141054.
53. Kamil AC. On the proper definition of cognitive ethology. In: *Animal cognition in nature 1998* (pp. 1–28). Academic Press.
54. Kandel ER. The biology of memory: a forty-year perspective. *J Neurosci*. 2009;29(41):12748–56.
55. Kashetsky T, Avgar T, Dukas R. The cognitive ecology of animal movement: evidence from birds and mammals. *Front Ecol Evol*. 2021;9: 724887.
56. Klappstein NJ, Michelot T, Fieberg J, Pedersen EJ, Mills FJ. Step selection functions with non-linear and random effects. *Method Ecol Evol*. 2024;15(8):1332–46.
57. Krebs JR, Healy SD, Shettleworth SJ. Spatial memory of Paridae: comparison of a storing and a non-storing species, the coal tit, *Parus ater*, and the great tit *P. major*. *Animal Behav*. 1990;39(6):1127–37.
58. Krebs JR, Sherry DF, Healy SD, Perry VH, Vaccarino AL. Hippocampal specialization of food-storing birds. *Proc Natl Acad Sci*. 1989;86(4):1388–92.
59. Lorenz KZ. The companion in the bird's world. *Auk*. 1937;54(3):245–73.
60. Lewis MA, Fagan WF, Auger-Méthé M, Frair J, Fryxell JM, Gros C, Gurarie E, Healy SD, Merkle JA. Learning and animal movement. *Front Ecol Evol*. 2021;9: 681704.
61. MacHutchon AG, Wellwood DW. Grizzly bear food habits in the northern Yukon, Canada. *Ursus*. 2003;225–35.
62. Matthiopoulos J, Fieberg JR, Aarts G. Species-habitat associations: spatial data, predictive models, and ecological insights. 2023
63. Matthysen E. Density-dependent dispersal in birds and mammals. *Ecography*. 2005;28(3):403–16.
64. McLoughlin C. Learner support in distance and networked learning environments: ten dimensions for successful design. *Dist Edu*. 2002;23(2):149–62.
65. Merkle JA, Fortin D, Morales JM. A memory-based foraging tactic reveals an adaptive mechanism for restricted space use. *Ecol Lett*. 2014;17(8):924–31.
66. Merkle JA, Potts JR, Fortin D. Energy benefits and emergent space use patterns of an empirically parameterized model of memory-based patch selection. *Oikos*. 2017. <https://doi.org/10.1111/oik.03356>.
67. Merkle JA, Sawyer H, Monteith KL, Dwinell SP, Fralick GL, Kauffman MJ. Spatial memory shapes migration and its benefits: evidence from a large herbivore. *Ecol Lett*. 2019;22(11):1797–805.
68. Michelot T, Klappstein NJ, Potts JR, Fieberg J. Understanding step selection analysis through numerical integration. *Method Ecol Evol*. 2024;15(1):24–35.
69. Miles D, Stedman M, Heald A. Living with COVID-19: balancing costs against benefits in the face of the virus. *Natl Inst Econ Rev*. 2020;253:R60-76.
70. Morrison TA, Merkle JA, Hopcraft JG, Aikens EO, Beck JL, Boone RB, Courtemanch AB, Dwinell SP, Fairbanks WS, Griffith B, Middleton AD. Drivers of site fidelity in ungulates. *J Animal Ecol*. 2021;90(4):955–66.
71. Nabe-Nielsen J, Tougaard J, Teilmann J, Lucke K, Forchhammer MC. How a simple adaptive foraging strategy can lead to emergent home ranges and increased food intake. *Oikos*. 2013;122(9):1307–16.

72. Nathan R, Getz WM, Revilla E, Holyoak M, Kadmon R, Saltz D, Smouse PE. A movement ecology paradigm for unifying organismal movement research. *Proc Natl Acad Sci*. 2008;105(49):19052–9.
73. Nathan R, Monk CT, Arlinghaus R, Adam T, Alós J, Assaf M, Baktoft H, Beardsworth CE, Bertram MG, Bijleveld AI, Brodin T. Big-data approaches lead to an increased understanding of the ecology of animal movement. *Science*. 2022;375(6582):1780.
74. Nesbitt SA. First reproductive success and individual productivity in Sandhill cranes. *J Wildlife Manag*. 1992;56:573.
75. Northrup JM, Anderson CR Jr, Wittemyer G. Environmental dynamics and anthropogenic development alter philopatry and space-use in a North American cervid. *Div Distribut*. 2016;22(5):547–57.
76. Oliveira-Santos LG, Forester JD, Piovezan U, Tomas WM, Fernandez FA. Incorporating animal spatial memory in step selection functions. *J Animal Ecol*. 2016;85(2):516–24.
77. Osipova L, Okello MM, Njumbi SJ, Ngene S, Western D, Hayward MW, Balkenhol N. Using step-selection functions to model landscape connectivity for African elephants: accounting for variability across individuals and seasons. *Animal Conserv*. 2019;22(1):35–48.
78. Pasitschniak-Arts M. *Ursus arctos*. *Mammal Spec*. 1993;23(439):1.
79. Picardi S, Abrahms B, Gelzer E, Morrison TA, Verzuh T, Merkle JA. Defining null expectations for animal site fidelity. *Ecol Lett*. 2023;26(1):157–69.
80. Piper WH. Making habitat selection more “familiar”: a review. *Behav Ecol Sociobiol*. 2011;65:1329–51.
81. Potts JR, Börger L. How to scale up from animal movement decisions to spatiotemporal patterns: an approach via step selection. *J Animal Ecol*. 2023;92(1):16–29.
82. Potts JR, Lewis MA. How memory of direct animal interactions can lead to territorial pattern formation. *J Royal Soc Interf*. 2016;13(118):20160059.
83. Potts JR, Schlägel UE. Parametrizing diffusion-taxis equations from animal movement trajectories using step selection analysis. *Method Ecol Evol*. 2020;11(9):1092–105.
84. Ranc N, Cagnacci F, Moorcroft PR. Memory drives the formation of animal home ranges: evidence from a reintroduction. *Ecol Lett*. 2022;25(4):716–28.
85. Ranc N, Moorcroft PR, Ossi F, Cagnacci F. Experimental evidence of memory-based foraging decisions in a large wild mammal. *Proc Natl Acad Sci*. 2021;118(15):e2014856118.
86. Rheault H, Anderson CR Jr, Bonar M, Marrotte RR, Ross TR, Wittemyer G, Northrup JM. Some memories never fade: Inferring multi-scale memory effects on habitat selection of a migratory ungulate using step-selection functions. *Front Ecol Evol*. 2021;9:702818.
87. Riotte-Lambert L, Benhamou S, Chamaillé-Jammes S. How memory-based movement leads to nonterritorial spatial segregation. *Am Nat*. 2015;185(4):E103–16.
88. Sawyer H, LeBeau CW, McDonald TL, Xu W, Middleton AD. All routes are not created equal: an ungulate’s choice of migration route can influence its survival. *J Appl Ecol*. 2019;56(8):1860–9.
89. Sawyer H, Middleton AD, Hayes MM, Kauffman MJ, Monteith KL. The extra mile: ungulate migration distance alters the use of seasonal range and exposure to anthropogenic risk. *Ecosphere*. 2016;7(10):e01534.
90. Schlägel UE, Lewis MA. Detecting effects of spatial memory and dynamic information on animal movement decisions. *Method Ecol Evol*. 2014;5(11):1236–46.
91. Schlägel UE, Merrill EH, Lewis MA. Territory surveillance and prey management: wolves keep track of space and time. *Ecol Evol*. 2017;7(20):8388–405.
92. Selva N, Teitelbaum CS, Sergiel A, Zwijacz-Kozica T, Zięba F, Bojarska K, Mueller T. Supplementary ungulate feeding affects movement behavior of brown bears. *Basic Appl Ecol*. 2017;24:68–76.
93. Sergio F, Tavecchia G, Blas J, López L, Tanferna A, Hiraldo F. Variation in age-structured vital rates of a long-lived raptor: implications for population growth. *Basic Appl Ecol*. 2011;12(2):107–15.
94. Sherry DF, Vaccarino AL, Buckenham K, Herz RS. The hippocampal complex of food-storing birds. *Brain, Behav Evol*. 1989;34(5):308–17.
95. Shettleworth SJ. *Cognition, evolution, and behavior*. Oxford university press; 2009.
96. Signer J, Fieberg J, Avgar T. Animal movement tools (amt): R package for managing tracking data and conducting habitat selection analyses. *Ecol Evol*. 2019;9(2):880–90.
97. Signer J, Fieberg J, Reineking B, Schlägel U, Smith B, Balkenhol N, Avgar T. Simulating animal space use from fitted integrated step-selection functions (iSSF). *Method Ecol Evol*. 2024;15(1):43–50.
98. Spencer WD. Home ranges and the value of spatial information. *J Mammal*. 2012;93(4):929–47.
99. Subaluskus AL, Dutton CL, Rosi EJ, Post DM. Annual mass drownings of the Serengeti wildebeest migration influence nutrient cycling and storage in the Mara River. *Proc Natl Acad Sci*. 2017;114(29):7647–52.
100. Tacha TC, Haley DE, Vohs PA. Age of sexual maturity of Sandhill cranes from mid-continental North America. *J Wildlife Manag*. 1989;1:43–6.
101. Tello-Ramos MC, Branch CL, Kozlovsky DY, Pitera AM, Pravosudov VV. Spatial memory and cognitive flexibility trade-offs: to be or not to be flexible, that is the question. *Animal Behav*. 2019;147:129–36.
102. Thompson PR, Derocher AE, Edwards MA, Lewis MA. Detecting seasonal episodic-like spatio-temporal memory patterns using animal movement modelling. *Method Ecol Evol*. 2022;13(1):105–20.
103. Thompson PR, Lewis MA, Edwards MA, Derocher AE. Time-dependent memory and individual variation in Arctic brown bears (*Ursus arctos*). *Mov Ecol*. 2022;10(1):18.
104. Thurfjell H, Ciuti S, Boyce MS. Applications of step-selection functions in ecology and conservation. *Mov Ecol*. 2014;2:1–2.
105. Trapanese C, Meunier H, Masi S. What, where and when: spatial foraging decisions in primates. *Biol Rev*. 2019;94(2):483–502.
106. Tulving E, Craik FI, editors. *The Oxford handbook of memory*. Oxford University Press; 2005.
107. van de Kerk M, Larsen RT, Olson DD, Hersey KR, McMillan BR. Variation in movement patterns of mule deer: have we oversimplified migration? *Mov Ecol*. 2021;9:1–2.
108. Van Moorter B, Visscher D, Benhamou S, Börger L, Boyce MS, Gaillard JM. Memory keeps you at home: a mechanistic model for home range emergence. *Oikos*. 2009;118(5):641–52.
109. Van Moorter B, Visscher D, Herfindal I, Basille M, Mysterud A. Inferring behavioural mechanisms in habitat selection studies getting the null-hypothesis right for functional and familiarity responses. *Ecography*. 2013;36(3):323–30.
110. Wang H, Salmani Y. Open problems in PDE models for knowledge-based animal movement via nonlocal perception and cognitive mapping. *J Mathem Biol*. 2023;86(5):71.
111. Wehner R, Hoinville T, Cruse H. On the ‘cognitive map debate’ in insect navigation. *Stud Hist Philos Sci*. 2023;102:87–9.
112. Welkin JF, Sonnenberg BR, Branch CL, Heinen VK, Pitera AM, Benedict LM, Whitenack LE, Bridge ES, Pravosudov VV. Spatial cognitive ability is associated with longevity in food-caching chickadees. *Science*. 2024;385(6713):1111–5.
113. Weise FJ, Stratford KJ, van Vuuren RJ. Financial costs of large carnivore translocations—accounting for conservation. *PLoS ONE*. 2014;9(8):e105042.
114. Whittington J, Hebblewhite M, Baron RW, Ford AT, Paczkowski J. Towns and trails drive carnivore movement behaviour, resource selection, and connectivity. *Mov Ecol*. 2022;10(1):17.
115. Wild S, Alarcón-Nieto G, Chimento M, Aplin LM. Manipulating actions: a selective two-option device for cognitive experiments in wild animals. *J Animal Ecol*. 2023;92(8):1509–19.
116. Wirsing AJ, Quinn TP, Cunningham CJ, Adams JR, Craig AD, Waits LP. Alaskan brown bears (*Ursus arctos*) aggregate and display fidelity to foraging neighborhoods while preying on Pacific salmon along small streams. *Ecol Evol*. 2018;8(17):9048–61.
117. Wolf M, Frair J, Merrill E, Turchin P. The attraction of the known: the importance of spatial familiarity in habitat selection in wapiti *Cervus elaphus*. *Ecography*. 2009;32(3):401–10.
118. Wolfson DW. *Migratory ecology and movement patterns of mid-continent and eastern sandhill cranes* (Master’s thesis, University of Minnesota) 2018
119. Wolfson DW, Fieberg JR, Andersen DE. Juvenile Sandhill Cranes exhibit wider ranging and more exploratory movements than adults during the breeding season. *Ibis*. 2020;162(2):556–62.

## Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.