

---

Faculty of Science

Faculty Publications

---

Approximation Properties of an Extended Family of the Szász–Mirakjan Beta-Type Operators

Hari Mohan Srivastava, Gürhan İçöz and Bayram Çekim

October 2019

© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license ( <http://creativecommons.org/licenses/by/4.0/> ).

This article was originally published at:

<http://dx.doi.org/10.3390/axioms8040111>


---

Citation for this paper:

Srivastava, H.M., İçöz, G. & Çekim, B. (2019). Approximation Properties of an Extended Family of the Szász–Mirakjan Beta-Type Operators. *Axioms*, 8(4), 111. <https://doi.org/10.3390/axioms8040111>

Article

# Approximation Properties of an Extended Family of the Szász–Mirakjan Beta-Type Operators

Hari Mohan Srivastava <sup>1,2,3,\*</sup> , Gürhan İçöz <sup>4</sup> and Bayram Çekim <sup>4</sup> <sup>1</sup> Department of Mathematics and Statistics, University of Victoria, Victoria, BC V8W 3R4, Canada<sup>2</sup> Department of Medical Research, China Medical University Hospital, China Medical University, Taichung 40402, Taiwan<sup>3</sup> Department of Mathematics and Informatics, Azerbaijan University, 71 Jeyhun Hajibeyli Street, AZ1007 Baku, Azerbaijan<sup>4</sup> Department of Mathematics, Gazi University, Ankara TR-06500, Turkey; gurhanicoz@gazi.edu.tr (G.İ.); bayramcekim@gazi.edu.tr (B.Ç.)

\* Correspondence: harimsri@math.uvic.ca

Received: 14 September 2019; Accepted: 30 September 2019; Published: 10 October 2019



**Abstract:** Approximation and some other basic properties of various linear and nonlinear operators are potentially useful in many different areas of researches in the mathematical, physical, and engineering sciences. Motivated essentially by this aspect of approximation theory, our present study systematically investigates the approximation and other associated properties of a class of the Szász–Mirakjan-type operators, which are introduced here by using an extension of the familiar Beta function. We propose to establish moments of these extended Szász–Mirakjan Beta-type operators and estimate various convergence results with the help of the second modulus of smoothness and the classical modulus of continuity. We also investigate convergence via functions which belong to the Lipschitz class. Finally, we prove a Voronovskaja-type approximation theorem for the extended Szász–Mirakjan Beta-type operators.

**Keywords:** gamma and beta functions; Szász–Mirakjan operators; Szász–Mirakjan Beta type operators; extended Gamma and Beta functions; confluent hypergeometric function; Modulus of smoothness; modulus of continuity; Lipschitz class; local approximation; Voronovskaja type approximation theorem

**2010 Mathematics Subject Classification:** Primary 33B15; 33C05; 41A25; 41A35; Secondary 33C20

## 1. Introduction, Definitions and Preliminaries

In approximation theory and its related fields, approximation and other basic properties of various linear and nonlinear operators are investigated, because mainly of the potential for their usefulness in many areas of researches in the mathematical, physical, and engineering sciences. Our study in this article is motivated essentially by the demonstrated applications of such results as those associated with various approximation operators. With this objective in view, we begin by providing the following definitions and other (chiefly historical) background material related to our presentation here.

For a given continuous function,  $f \in C[0, \infty)$ , and for  $x \in [0, \infty)$ , Otto Szász [1] defined a family of operators in the year 1950, which we recall here as follows,

$$S_n(f; x) = e^{-nx} \sum_{k=0}^{\infty} \frac{(nx)^k}{k!} f\left(\frac{k}{n}\right) \quad (n \in \mathbb{N} := \{1, 2, 3, \dots\}), \quad (1)$$

This family of operators was considered earlier in 1941 by G. M. Mirakjan (see [2]). There are several integral and other modifications, variations, and basic (or  $q$ -) extensions of the Szász-Mirakjan-type operators. These include the Bézier, Kantorovich, Durrmeyer, and other types of modifications and extensions of the Szász-Mirakjan operators (see, for details, [3–15]). In particular, Gupta and Noor [6] introduced an integral modification of the Szász-Mirakjan operators in Equation (1) by considering a weight function in terms of the Beta basis functions as given below.

$$T_n(f; x) = \sum_{k=1}^{\infty} s_{n,k}(x) \int_0^{\infty} b_{n,k}(t) f(t) dt + s_{n,0}(x) f(0), \tag{2}$$

where

$$s_{n,k}(x) = e^{-nx} \frac{(nx)^k}{k!},$$

$$b_{n,k}(t) = \frac{1}{B(k, n+1)} \frac{t^{k-1}}{(1+t)^{n+k+1}} = \frac{(n+1)_k}{(k-1)!} \frac{t^{k-1}}{(1+t)^{n+k+1}},$$

$$B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)} = B(\beta, \alpha),$$

and  $(\lambda)_\ell$  ( $\ell \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$ ) represents the Pochhammer symbol given by

$$(\lambda)_\ell := \frac{\Gamma(\lambda+\ell)}{\Gamma(\lambda)} = \begin{cases} 1 & (\ell = 0) \\ \lambda(\lambda+1)(\lambda+2) \cdots (\lambda+\ell-1) & (\ell \in \mathbb{N}) \end{cases}$$

in terms of the classical (Euler’s) Gamma function  $\Gamma(z)$  and the classical Beta function  $B(\alpha, \beta)$ .

Gupta and Noor [6] observed that the operators in Equation (2) reproduce not only the constant function, but linear functions as well. Owing to this valuable property of the operators in Equation (2), many authors investigated the different approximation properties of the summation-integral operators in Equation (2) (see, for example, [16–18]). Gupta and Noor [6] also derived some direct results for the operators  $T_n$ , a pointwise rate of convergence, a Voronovskaja-type asymptotic formula, and an error estimate in simultaneous approximation.

In recent years, some extensions of such well-known special functions as, for example, the classical Gamma and Beta functions, have been considered by several authors. For example, in 1994, Chaudhry and Zubair [19] introduced the following extension of the Gamma function,

$$\Gamma_p(x) := \int_0^{\infty} t^{x-1} \exp\left(-t - \frac{p}{t}\right) dt \quad (\Re(p) > 0). \tag{3}$$

Subsequently, in 1997, Chaudhry et al. [20] presented the following extension of Euler’s Beta function,

$$B_p(x, y) := \int_0^1 t^{x-1} (1-t)^{y-1} \exp\left(-\frac{p}{t(1-t)}\right) dt \tag{4}$$

$$(\Re(p) > 0; \Re(x) > 0; \Re(y) > 0).$$

Obviously, each of the definitions in Equations (3) and (4) also remains valid when  $p = 0$ , in which case we have the following relationships,

$$\Gamma_0(x) = \Gamma(x) \quad \text{and} \quad B_0(x, y) = B(x, y).$$

Özergin et al. [21] considered the following generalizations of the Gamma and Beta functions,

$$\Gamma_p^{(\alpha,\beta)}(x) := \int_0^\infty t^{x-1} {}_1F_1\left(\alpha; \beta; -t - \frac{p}{t}\right) dt$$

$$(\Re(\alpha) > 0; \Re(\beta) > 0; \Re(p) > 0; \Re(x) > 0)$$

and

$$B_p^{(\alpha,\beta)}(x,y) := \int_0^1 t^{x-1} (1-t)^{y-1} {}_1F_1\left(\alpha; \beta; -\frac{p}{t(1-t)}\right) dt$$

$$(\Re(\alpha) > 0; \Re(\beta) > 0; \Re(p) > 0; \Re(x) > 0; \Re(y) > 0),$$

respectively. Here, as usual,  ${}_1F_1$  denotes the (Kummer’s) confluent hypergeometric function. It is obvious that

$$\Gamma_p^{(\alpha,\alpha)}(x) = \Gamma_p(x) \quad \text{and} \quad \Gamma_0^{(\alpha,\alpha)}(x) = \Gamma(x)$$

and that

$$B_p^{(\alpha,\alpha)}(x,y) = B_p(x,y) \quad \text{and} \quad B_0^{(\alpha,\alpha)}(x,y) = B(x,y).$$

Finding different integral representations of the generalized Beta function is important and useful for later use. It is also useful to discuss the relationships between the classical Gamma and Beta functions and their generalizations. In fact, by definition, it is easily seen that

$$\int_0^\infty \Gamma_p(x) dp = \Gamma(x+1) \quad (\Re(x) > -1).$$

and that (see, for example, [21])

$$\int_0^\infty B_p(x,y) dp = B(x+1,y+1) \tag{5}$$

$$(\Re(x) > -1; \Re(y) > -1).$$

Note that various further extensions and generalizations of the classical Gamma and Beta functions, as well as their corresponding hypergeometric and related functions, were introduced and studied by, among others, Lin et al. [22] and Srivastava et al. [23].

We now introduce the following generalization of Szász-Mirakjan Beta-type operators via the above extension of the Beta function as follows,

$$S_n^*(f,x) = \sum_{k=0}^\infty e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2,n+1)} \cdot \int_0^\infty \int_0^\infty \frac{t^k}{(1+t)^{n+k+1}} \exp\left(-\frac{p(1+t)^2}{t}\right) f(t) dt dp \tag{6}$$

for  $x \in [0, \infty)$ , and for a function  $f \in C_\nu[0, \infty)$ , provided that the double integral in Equation (6) is convergent when  $n > \nu$ . Here, and in what follows, we have

$$C_\nu[0, \infty) := \{f : f \in C[0, \infty) \quad \text{and} \quad |f(t)| \leq M(1+t)^\nu \quad (M > 0; \nu > 0)\}.$$

We note that, by setting  $t = \frac{u}{1+u}$  in Equation (4), we get

$$B_p(x, y) = \int_0^\infty \frac{u^{x-1}}{(1+u)^{x+y}} \exp\left(-\frac{p(1+u)^2}{u}\right) du. \tag{7}$$

So, if we take in consideration Equations (5) and (7) in the definition Equation (6), then we can say that the operators,  $S_n^*$ , are a generalization of the operators,  $T_n$ , given by Equation (2).

In this article, we investigate the moments of the general Szász-Mirakjan Beta-type operators  $S_n^*$  and find the rate of convergence with the help of the classical and second moduli of continuity. We also derive a Voronovskaja-type approximation theorem associated with these general operators,  $S_n^*$ .

### 2. A Set of Auxiliary Results

In this section, we give the moments of Szász-Mirakjan Beta-type operators,  $S_n^*$ , defined by Equation (6). We first recall for the  $S_n$  that

$$S_n(1, x) = 1, \quad S_n(t, x) = x \quad \text{and} \quad S_n(t^2, x) = x^2 + \frac{x}{n}, \tag{8}$$

just as in [1].

**Lemma 1.** *The moments of the Szász-Mirakjan Beta-type operators,  $S_n^*$ , defined by (6) are given by*

$$S_n^*(1, x) = 1, \tag{9}$$

$$S_n^*(t, x) = x + \frac{2}{n} \tag{10}$$

and

$$S_n^*(t^2, x) = \frac{n}{n-1} x^2 + \frac{6}{n-1} x + \frac{6}{n(n-1)}. \tag{11}$$

**Proof.** By using the known formulas in Equation (8), we find from the definition (6) that

$$\begin{aligned} S_n^*(1, x) &= \sum_{k=0}^\infty e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2, n+1)} \\ &\quad \cdot \int_0^\infty \int_0^\infty \frac{t^k}{(1+t)^{n+k+1}} \exp\left(-\frac{p(1+t)^2}{t}\right) dt dp \\ &= \sum_{k=0}^\infty e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2, n+1)} \int_0^\infty B_p(k+1, n) dp \\ &= \sum_{k=0}^\infty e^{-nx} \frac{(nx)^k}{k!} \frac{B(k+2, n+1)}{B(k+2, n+1)} \\ &= S_n(1, x) = 1. \end{aligned}$$

For  $n > 1$ , we have

$$\begin{aligned}
 S_n^*(t, x) &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2, n+1)} \\
 &\quad \cdot \int_0^{\infty} \int_0^{\infty} \frac{t^{k+1}}{(1+t)^{n+k+1}} \exp\left(-\frac{p(1+t)^2}{t}\right) dt dp \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2, n+1)} \\
 &\quad \cdot \int_0^{\infty} B_p(k+2, n-1) dp \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{B(k+3, n)}{B(k+2, n+1)} \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{\Gamma(k+3)\Gamma(n)}{\Gamma(n+k+3)} \frac{\Gamma(n+k+3)}{\Gamma(k+2)\Gamma(n+1)} \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{k+2}{n} \\
 &= S_n(t, x) + \frac{2}{n} S_n(1, x) = x + \frac{2}{n}
 \end{aligned}$$

and, for  $n > 2$ , we find that

$$\begin{aligned}
 S_n^*(t^2, x) &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2, n+1)} \\
 &\quad \cdot \int_0^{\infty} \int_0^{\infty} \frac{t^{k+2}}{(1+t)^{n+k+1}} \exp\left(-\frac{p(1+t)^2}{t}\right) dt dp \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2, n+1)} \\
 &\quad \cdot \int_{p=0}^{\infty} B_p(k+3, n-2) dp \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{B(k+4, n-1)}{B(k+2, n+1)} \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{\Gamma(k+4)\Gamma(n-1)}{\Gamma(n+k+3)} \frac{\Gamma(n+k+3)}{\Gamma(k+2)\Gamma(n+1)} \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{(k+3)(k+2)}{n(n-1)} \\
 &= \frac{n}{n-1} S_n(t^2, x) + \frac{5}{n-1} S_n(t, x) + \frac{6}{n(n-1)} S_n(1, x) \\
 &= \frac{n}{n-1} x^2 + \frac{6}{n-1} x + \frac{6}{n(n-1)}.
 \end{aligned}$$

The proof of Lemma 1 is thus completed.  $\square$

**Lemma 2.** The central moments of the Szász-Mirakjan Beta-type operators,  $S_n^*$ , defined by Equation (6) are given by

$$S_n^*(t-x, x) = \frac{2}{n} \tag{12}$$

and

$$S_n^*((t-x)^2, x) = \frac{1}{n-1} x^2 + \frac{2(n+2)}{n(n-1)} x + \frac{6}{n(n-1)} =: \varepsilon_n(x). \tag{13}$$

**Proof.** The assertions (12) and (13) of Lemma 2 follow easily from those of Lemma 1, so we omit the details involved.  $\square$

### 3. Local Approximation

Let  $C_B[0, \infty)$  be the set of all real-valued continuous and bounded functions  $f$  on  $[0, \infty)$ , which is endowed with the norm given by

$$\|f\| = \sup_{x \in [0, \infty)} |f(x)|.$$

Then Peetre’s  $K$ -functional is defined by

$$K_2(f; \delta) = \inf \{ \|f - g\| + \delta \|g''\| : g \in C_B^2[0, \infty) \},$$

where

$$C_B^2[0, \infty) := \{g : g \in C_B[0, \infty) \text{ and } g', g'' \in C_B[0, \infty)\}.$$

There exists a positive constant  $C > 0$  such that (see, for example, [24])

$$K_2(f, \delta) \leq C \omega_2(f, \sqrt{\delta}), \tag{14}$$

where  $\delta > 0$  and  $\omega_2$  denotes the second-order modulus of smoothness for  $f \in C_B[0, \infty)$ , which is defined by

$$\omega_2(f; \sqrt{\delta}) = \sup_{0 < h \leq \delta} \sup_{x \in [0, \infty)} |f(x+2h) - 2f(x+h) + f(x)|.$$

The usual modulus of continuity for  $f \in C_B[0, \infty)$  is given by

$$\omega(f; \delta) = \sup_{0 < h \leq \delta} \sup_{x \in [0, \infty)} |f(x+h) - f(x)|.$$

Lemma 3 below provides an auxiliary inequality which is useful in proving our next theorem (see Theorem 1).

**Lemma 3.** For all  $g \in C_B^2[0, \infty)$ , it is asserted that

$$|\tilde{S}_n^*(g, x) - g(x)| \leq \delta_n(x) \|g''(x)\|, \tag{15}$$

where

$$\delta_n(x) = \frac{1}{n-1} x^2 + \frac{2(n+2)}{n(n-1)} x + \frac{2(5n-2)}{n^2(n-1)} \tag{16}$$

and

$$\tilde{S}_n^*(f, x) = S_n^*(f, x) + f(x) - f\left(x + \frac{2}{n}\right) \tag{17}$$

for  $f \in C_B[0, \infty)$ .

**Proof.** First of all, we find from (17) that

$$\tilde{S}_n^*(t-x, x) = S_n^*(t-x, x) - \frac{2}{n} = \frac{2}{n} - \frac{2}{n} = 0. \tag{18}$$

Now, by using the Taylor’s formula, we have

$$g(t) - g(x) = (t - x)g'(x) + \int_x^t (t - u)g''(u) du,$$

which, in view of Equation (18), yields

$$\begin{aligned} \tilde{S}_n^*(g, x) - g(x) &= \tilde{S}_n^*(t - x, x)g'(x) + \tilde{S}_n^*\left(\int_x^t (t - u)g''(u) du, x\right) \\ &= S_n^*\left(\int_x^t (t - u)g''(u)du, x\right) - \int_x^{x+\frac{2}{n}} \left(x + \frac{2}{n} - u\right) g''(u) du. \end{aligned}$$

On the other hand, as

$$\begin{aligned} \left| \int_x^t (t - u)g''(u) du \right| &\leq \int_x^t |t - u| \cdot |g''(u)| du \\ &\leq \|g''\| \int_x^t |t - u| du \\ &\leq (t - x)^2 \|g''\| \end{aligned}$$

and

$$\left| \int_x^{x+\frac{2}{n}} \left(x + \frac{2}{n} - u\right) g''(u) du \right| \leq \left(\frac{2}{n}\right)^2 \|g''\|,$$

we conclude that

$$\begin{aligned} \left| \tilde{S}_n^*(g, x) - g(x) \right| &\leq S_n^*((t - x)^2, x) \|g''\| + \frac{4}{n^2} \|g''\| \\ &= \left(\frac{1}{n - 1} x^2 + \frac{2(n + 2)}{n(n - 1)} x + \frac{6}{n(n - 1)} + \frac{4}{n^2}\right) \|g''\| \\ &= \left(\frac{1}{n - 1} x^2 + \frac{2(n + 2)}{n(n - 1)} x + \frac{2(5n - 2)}{n^2(n - 1)}\right) \|g''\| = \delta_n(x) \|g''\|. \end{aligned}$$

This is the result asserted by Lemma 3. □

We now state and prove our main results in this section.

**Theorem 1.** Let  $f \in C_B[0, \infty)$ . Then, for every  $x \in [0, \infty)$ , there exists a constant  $L > 0$ , such that

$$|S_n^*(f, x) - f(x)| \leq L\omega_2\left(f; \sqrt{\delta_n(x)}\right) + \omega\left(f; \frac{2}{n}\right),$$

where  $\omega_2(f; \delta)$  is the second-order modulus of smoothness,  $\omega(f; \delta)$  is the usual modulus of continuity, and  $\delta_n(x)$  is given by Equation (16).

**Proof.** We observe from Equation (17) that

$$\begin{aligned} |S_n^*(f, x) - f(x)| &\leq \left| \tilde{S}_n^*(f, x) - f(x) \right| + \left| f(x) - f\left(x + \frac{2}{n}\right) \right| \\ &\leq \left| \tilde{S}_n^*(f - g, x) - (f - g)(x) \right| + \left| f(x) - f\left(x + \frac{2}{n}\right) \right| \\ &\quad + \left| \tilde{S}_n^*(g, x) - g(x) \right| \end{aligned}$$



for  $g \in C_B^2[0, \infty)$ . Thus, by applying Lemma 3 for  $g \in C_B^2[0, \infty)$ , we get

$$|S_n^*(f, x) - f(x)| \leq 4 \|f - g\| + \delta_n(x) \|g''\| + \omega\left(f; \frac{2}{n}\right),$$

which, by taking the infimum on the right-hand side over all  $g \in C_B^2[0, \infty)$  and using (14), yields

$$\begin{aligned} |S_n^*(f, x) - f(x)| &\leq 4K_2(f; \delta_n(x)) + \omega\left(f; \frac{2}{n}\right) \\ &\leq L\omega_2\left(f; \sqrt{\delta_n(x)}\right) + \omega\left(f; \frac{2}{n}\right). \end{aligned}$$

where  $L = 4M > 0$ . This evidently completes the demonstration of Theorem 1.  $\square$

**Theorem 2.** Let  $E$  be any bounded subset of the interval  $[0, \infty)$ , and suppose that  $0 < \alpha \leq 1$ . If  $f \in C_B[0, \infty)$  is locally  $\text{Lip}_M(\alpha)$ , that is, if the following inequality holds true,

$$|f(y) - f(x)| \leq M|y - x|^\alpha \quad (y \in E; x \in [0, \infty)),$$

then, for each  $x \in [0, \infty)$ ,

$$|S_n^*(f, x) - f(x)| \leq M([\varepsilon_n(x)]^{\frac{\alpha}{2}} + 2[d(x, E)]^\alpha), \tag{19}$$

$\varepsilon_n(x)$  is given by Equation (13),  $M$  is a constant depending on  $\alpha$  and  $f$ , and  $d(x, E)$  is the distance between  $x$  and  $E$  defined as follows:

$$d(x, E) = \inf \{ |y - x| : y \in E \}.$$

**Proof.** Let  $\bar{E}$  denote the closure of  $E$  in  $[0, \infty)$ . Then there exists a point  $x_0 \in \bar{E}$  such that

$$|x - x_0| = d(x, E).$$

By the above-mentioned definition of  $\text{Lip}_M(\alpha)$ , we get

$$\begin{aligned} |S_n^*(f, x) - f(x)| &\leq S_n^*(|f(y) - f(x)|, x) \\ &\leq S_n^*(|f(y) - f(x_0)|, x) + S_n^*(|f(x) - f(x_0)|, x) \\ &\leq M\{S_n^*(|y - x_0|^\alpha, x) + |x - x_0|^\alpha\} \\ &\leq M\{S_n^*(|y - x|^\alpha + |x - x_0|^\alpha, x) + |x - x_0|^\alpha\} \\ &\leq M\{S_n^*(|y - x_0|^\alpha, x) + 2|x - x_0|^\alpha\}. \end{aligned}$$

Now, if we use the Hölder inequality with

$$p = \frac{2}{\alpha} \quad \text{and} \quad q = \frac{2}{2 - \alpha},$$

we find that

$$\begin{aligned} |S_n^*(f, x) - f(x)| &\leq M\left(\left[S_n^*((y - x_0)^2, x)\right]^{\frac{\alpha}{2}} + 2[d(x, E)]^\alpha\right) \\ &= M\left([\varepsilon_n(x)]^{\frac{\alpha}{2}} + 2[d(x, E)]^\alpha\right). \end{aligned}$$

We have thus completed our demonstration of the result asserted by Theorem 2.  $\square$

#### 4. A Voronovskaja-Type Approximation Theorem

By applying Equations (5) to (7), as well as Lemma 1, we first prove the following result.

**Lemma 4.** *It is asserted that*

$$S_n^*(t^3, x) = \frac{n^2}{(n-1)(n-2)} x^3 + \frac{12n}{(n-1)(n-2)} x^2 + \frac{36}{(n-1)(n-2)} x + \frac{24}{n(n-1)(n-2)} \tag{20}$$

and

$$S_n^*(t^4, x) = \frac{n^3}{(n-1)(n-2)(n-3)} x^4 + \frac{20n^2}{(n-1)(n-2)(n-3)} x^3 + \frac{120n}{(n-1)(n-2)(n-3)} x^2 + \frac{240}{(n-1)(n-2)(n-3)} x + \frac{120}{n(n-1)(n-2)(n-3)}. \tag{21}$$

Furthermore, the following result holds true,

$$S_n^*((t-x)^4, x) = \frac{3(n+6)}{(n-1)(n-2)(n-3)} x^4 + \frac{4(3n^2+32n+12)}{n(n-1)(n-2)(n-3)} x^3 + \frac{12(n^2+21n+18)}{n(n-1)(n-2)(n-3)} x^2 + \frac{144(n+2)}{n(n-1)(n-2)(n-3)} x + \frac{120}{n(n-1)(n-2)(n-3)}. \tag{22}$$

**Proof.** We begin by recalling the following moments of the Szász-Mirakjan operators,

$$S_n(t^3, x) = x^3 + \frac{3x^2}{n} + \frac{x}{n^2} \tag{23}$$

and

$$S_n(t^4, x) = x^4 + \frac{6x^3}{n} + \frac{7x^2}{n^2} + \frac{x}{n^3}. \tag{24}$$

Using Equations (8) and the above formulas (23) and (24), we thus find for  $n > 3$  that

$$\begin{aligned}
 S_n^*(t^3, x) &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2, n+1)} \\
 &\quad \cdot \int_0^{\infty} \int_0^{\infty} \frac{t^{k+3}}{(1+t)^{n+k+1}} \exp\left(-\frac{p(1+t)^2}{t}\right) dt dp \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2, n+1)} \int_0^{\infty} B_p(k+4, n-3) dp \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{B(k+5, n-2)}{B(k+2, n+1)} \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{\Gamma(k+5)\Gamma(n-2)}{\Gamma(n+k+3)} \frac{\Gamma(n+k+3)}{\Gamma(k+2)\Gamma(n+1)} \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{(k+4)(k+3)(k+2)}{n(n-1)(n-2)} \\
 &= \frac{n^2}{(n-1)(n-2)} S_n(t^3, x) + \frac{9n}{(n-1)(n-2)} S_n(t^2, x) \\
 &\quad + \frac{26}{(n-1)(n-2)} S_n(t, x) + \frac{24}{n(n-1)(n-2)} S_n(1, x) \\
 &= \frac{n^2}{(n-1)(n-2)} \left(x^3 + \frac{3x^2}{n} + \frac{x}{n^2}\right) + \frac{9n}{(n-1)(n-2)} \left(x^2 + \frac{x}{n}\right) \\
 &\quad + \frac{26}{(n-1)(n-2)} x + \frac{24}{n(n-1)(n-2)} \\
 &= \frac{n^2}{(n-1)(n-2)} x^3 + \frac{12n}{(n-1)(n-2)} x^2 + \frac{36}{(n-1)(n-2)} x \\
 &\quad + \frac{24}{n(n-1)(n-2)}.
 \end{aligned}$$

On the other hand, for  $n > 4$ , we find that

$$\begin{aligned}
 S_n^*(t^4, x) &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2, n+1)} \\
 &\quad \cdot \int_0^{\infty} \int_0^{\infty} \frac{t^{k+4}}{(1+t)^{n+k+1}} \exp\left(-\frac{p(1+t)^2}{t}\right) dt dp \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{1}{B(k+2, n+1)} \int_0^{\infty} B_p(k+5, n-4) dp \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{B(k+6, n-3)}{B(k+2, n+1)} \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{\Gamma(k+6)\Gamma(n-3)}{\Gamma(n+k+3)} \frac{\Gamma(n+k+3)}{\Gamma(k+2)\Gamma(n+1)} \\
 &= \sum_{k=0}^{\infty} e^{-nx} \frac{(nx)^k}{k!} \frac{(k+5)(k+4)(k+3)(k+2)}{n(n-1)(n-2)(n-3)},
 \end{aligned}$$

that is, that

$$\begin{aligned}
 S_n^*(t^4, x) &= \frac{n^3}{(n-1)(n-2)(n-3)} S_n(t^4, x) + \frac{14n^2}{(n-1)(n-2)(n-3)} S_n(t^3, x) \\
 &\quad + \frac{71n}{(n-1)(n-2)(n-3)} S_n(t^2, x) + \frac{154}{(n-1)(n-2)(n-3)} S_n(t, x) \\
 &\quad + \frac{120}{n(n-1)(n-2)(n-3)} S_n(1, x) \\
 &= \frac{n^3}{(n-1)(n-2)(n-3)} x^4 + \frac{20n^2}{(n-1)(n-2)(n-3)} x^3 \\
 &\quad + \frac{120n}{(n-1)(n-2)(n-3)} x^2 + \frac{240}{(n-1)(n-2)(n-3)} x \\
 &\quad + \frac{120}{n(n-1)(n-2)(n-3)},
 \end{aligned}$$

which, together, complete the proof of Lemma 4.  $\square$

**Theorem 3.** Let  $f, f', f'' \in C_v[0, \infty)$  for  $v \geq 4$ . Then, the following Voronovskaja-type approximation result holds true,

$$\lim_{n \rightarrow \infty} \{n [S_n^*(f, x) - f(x)]\} = 2f'(x) + \left(\frac{x^2}{2} + x\right) f''(x). \tag{25}$$

**Proof.** By Taylor’s expansion of  $f(t)$  at the point  $t = x$ , we have

$$f(t) = f(x) + f'(x)(t - x) + \frac{1}{2} f''(x)(t - x)^2 + \Psi(t, x)(t - x)^2, \tag{26}$$

where  $\Psi(t, x)$  is remainder term,  $\Psi(\cdot, x) \in C_v[0, \infty)$  and  $\Psi(t, x) \rightarrow 0$  as  $t \rightarrow x$ .

Applying the Szász-Mirakjan Beta-type operators  $S_n^*$  to Equation (26) and, using Lemma 2, we obtain

$$\begin{aligned}
 S_n^*(f, x) - f(x) &= f'(x) S_n^*(t - x, x) + \frac{1}{2} f''(x) S_n^*((t - x)^2, x) \\
 &\quad + S_n^*(\Psi(t, x)(t - x)^2, x) \\
 &= \frac{2}{n} f'(x) + \frac{1}{2} \left(\frac{1}{n-1} x^2 + \frac{2(n+2)}{n(n-1)} x + \frac{6}{n(n-1)}\right) f''(x) \\
 &\quad + S_n^*(\Psi(t, x)(t - x)^2, x).
 \end{aligned} \tag{27}$$

We now apply the Cauchy-Schwarz inequality to the third term on the right-hand side of Equation (27). We thus find that

$$n \left| S_n^*(\Psi(t, x)(t - x)^2, x) \right| \leq \sqrt{n^2 S_n^*((t - x)^4, x)} \cdot \sqrt{S_n^*([\Psi(t, x)]^2, x)}.$$

Let

$$\eta(t, x) := [\Psi(t, x)]^2.$$

In this case, we observe that  $\eta(x, x) = 0$  and also that  $\eta(\cdot, x) \in C_v[0, \infty)$ . Then, it follows that

$$\lim_{n \rightarrow \infty} \{S_n^*([\Psi(t, x)]^2, x)\} = \lim_{n \rightarrow \infty} \{S_n^*(\eta(t, x), x)\} = \eta(x, x) = 0$$

uniformly with respect to  $x \in [0, b]$  ( $b > 0$ ) and the following limit,

$$\lim_{n \rightarrow \infty} \{n^2 S_n^*((t-x)^4, x)\}$$

is finite. Consequently, we have

$$\lim_{n \rightarrow \infty} \{n S_n^*(\Psi(t, x) (t-x)^2, x)\} = 0.$$

Thus, in the limit when  $n \rightarrow \infty$  in Equation (27), we obtain

$$\lim_{n \rightarrow \infty} \{n[S_n^*(f, x) - f(x)]\} = 2f'(x) + \left(\frac{x^2}{2} + x\right) f''(x).$$

The proof of Theorem 3 is thus completed.  $\square$

## 5. Concluding Remarks and Observations

We find it worthwhile to reiterate the fact that, in approximation theory and related fields, the approximation and some other basic properties of various linear and nonlinear operators are investigated because mainly of the potential for their usefulness in many areas of researches in the mathematical, physical, and engineering sciences. This article has been motivated essentially by the demonstrated applications of such results as those associated with various approximation operators.

In our present investigation, we have systematically studied a number of approximation properties of a class of the Szász-Mirakjan Beta-type operators, which we have introduced here by using an extension of the familiar Beta function  $B(\alpha, \beta)$ . We have established the moments of these extended Szász-Mirakjan Beta-type operators and estimated several convergence results with the help of the second modulus of smoothness and the classical modulus of continuity. We have also investigated convergence via functions belonging to the Lipschitz class. Finally, we have proved a Voronovskaja-type approximation theorem for the general Szász-Mirakjan Beta-type operators.

Using the other substantially more general forms of the classical Beta function  $B(\alpha, \beta)$ , which we have indicated in Section 1 of this article (see, for example, [22,23]), one can analogously develop further extensions and generalizations of the various results which we have presented here. In many of these suggested areas of further researches on the subject of this article, some other, possibly deeper, mathematical analytic tools and techniques will have to be called for.

**Author Contributions:** All three authors contributed equally to this investigation.

**Funding:** This research received no external funding.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Szász, O. Generalizations of S. Bernstein's polynomials to the infinite interval. *J. Res. Nat. Bur. Stand.* **1950**, *45*, 239–245. [[CrossRef](#)]
2. Mirakjan, G.M. Approximation des fonctions continues au moyen de polynômes de la forme  $e^{-nx} \sum_{k=0}^m C_{k,n} x^k$ . *C. R. (Doklady) Acad. Sci. URSS (New Ser.)* **1941**, *31*, 201–205.
3. Duman, O.; Özarlan, M.A. Szász-Mirakjan-type operators providing a better error estimation. *Appl. Math. Lett.* **2007**, *20*, 1184–1188. [[CrossRef](#)]
4. Gal, S.G. Approximation with an arbitrary order by generalized Szász-Mirakjan operators. *Stud. Univ. Babeş-Bolyai Math.* **2014**, *59*, 77–81.
5. Gupta, V.; Mahmudov, N. Approximation properties of the  $q$ -Szász-Mirakjan-Beta operators. *Indian J. Ind. Appl. Math.* **2012**, *3*, 41–53.
6. Gupta, V.; Noor, M.A. Convergence of derivatives for certain mixed Szász-Beta operators. *J. Math. Anal. Appl.* **2006**, *321*, 1–9. [[CrossRef](#)]

7. İçöz, G.; Mohapatra, R.N. Approximation properties by  $q$ -Durrmeyer-Stancu operators. *Anal. Theory Appl.* **2013**, *29*, 373–383.
8. Gupta, V.; Srivastava, H.M. A general family of the Srivastava-Gupta operators preserving linear functions. *Eur. J. Pure Appl. Math.* **2018**, *11*, 575–579. [[CrossRef](#)]
9. Gupta, V.; Srivastava, G.S.; Sahai, A. On simultaneous approximation by Szász-beta operators. *Soochow J. Math.* **1995**, *21*, 1–11.
10. Păltănea, R. Modified Szász-Mirakjan operators of integral form. *Carpathian J. Math.* **2008**, *24*, 378–385.
11. Srivastava, H.M.; Mursaleen, M.; Alotaibi, A.M.; Nasiruzzaman, M.; Al-Abied, A.A.H. Some approximation results involving the  $q$ -Szász-Mirakjan-Kantorovich type operators via Dunkl's generalization. *Math. Methods Appl. Sci.* **2017**, *40*, 5437–5452. [[CrossRef](#)]
12. Srivastava, H.M.; Özger, F.; Mohiuddine, S.A. Construction of Stancu-type Bernstein operators based on Bézier bases with shape parameter  $\lambda$ . *Symmetry* **2019**, *11*, 316. [[CrossRef](#)]
13. Srivastava, H.M.; Zeng, X.-M. Approximation by means of the Szász-Bézier integral operators. *Int. J. Pure Appl. Math.* **2004**, *14*, 283–294.
14. Xie, L.-S.; Xie, T.-F. Approximation theorems for localized Szász-Mirakjan operators. *J. Approx. Theory* **2008**, *152*, 125–134. [[CrossRef](#)]
15. Zeng, X.-M. Approximation of absolutely continuous functions by Stancu Beta operators. *Ukrainian Math. J.* **2012**, *63*, 1787–1794. [[CrossRef](#)]
16. Duman, O.; Özarşlan, M.A.; Aktuğlu, H. Better error estimates for Szász-Mirakjan-Beta operators. *J. Comput. Anal. Appl.* **2008**, *10*, 53–59.
17. Özarşlan, M.A.; Aktuğlu, H.  $A$ -Statistical approximation of generalized Szász-Mirakjan-Beta operators. *Appl. Math. Lett.* **2011**, *24*, 1785–1790. [[CrossRef](#)]
18. Qi, Q.-L.; Zhang, Y.-P. Pointwise approximation for certain mixed Szász-Beta operators. In *Further Progress in Analysis, Proceedings of the Sixth International Conference (ISAAC 2002) on Clifford Algebras and Their Applications in Mathematical Physics, Tennessee Technological University, Cookeville, TN, USA, 20–25 May 2002*; World Scientific Publishing Company: Singapore, 2009; pp. 152–164.
19. Chaudhry, M.A.; Zubair, S.M. Generalized incomplete gamma functions with applications. *J. Comput. Appl. Math.* **1994**, *55*, 99–124. [[CrossRef](#)]
20. Chaudhry, M.A.; Qadir, A.; Rafique, M.; Zubair, S.M. Extension of Euler's beta function. *J. Comput. Appl. Math.* **1997**, *78*, 19–32. [[CrossRef](#)]
21. Özergin, E.; Özarşlan, M.A.; Altın, A. Extension of gamma, beta and hypergeometric functions. *J. Comput. Appl. Math.* **2011**, *235*, 4601–4610. [[CrossRef](#)]
22. Lin, S.-D.; Srivastava, H.M.; Yao, J.-C. Some classes of generating relations associated with a family of the generalized Gauss type hypergeometric functions. *Appl. Math. Inform. Sci.* **2015**, *9*, 1731–1738.
23. Srivastava, H.M.; Parmar, R.K.; Chopra, P. A class of extended fractional derivative operators and associated generating relations involving hypergeometric functions. *Axioms* **2012**, *1*, 238–258. [[CrossRef](#)]
24. DeVore, R.A.; Lorentz, G.G. *Constructive Approximation*; Springer: Berlin/Heidelberg, Germany; New York, NY, USA, 1993.

