

***THE FEKETE-SZEGÖ PROBLEM FOR
A SUBCLASS OF CLOSE-TO-CONVEX
FUNCTIONS***

H.M. SRIVASTAVA, A.K. MISHRA, & M.K. DAS

DMS-832-IR

June 1999

THE FEKETE-SZEGŐ PROBLEM FOR A SUBCLASS OF CLOSE-TO-CONVEX FUNCTIONS

H.M. Srivastava

Department of Mathematics and Statistics
University of Victoria
Victoria, British Columbia V8W 3P4, Canada
E-Mail: hmsri@uvvm.uvic.ca

A.K. Mishra

Department of Mathematics
Berhampur University
Berhampur 760 007, Orissa, India
and

M.K. Das

Department of Mathematics
Government Science College
Chatrapur 761 020, Orissa, India

Abstract

Let $\mathcal{C}_1(\beta)$ be the class of normalized functions f , which are analytic in the open unit disk \mathcal{U} , given by the power series:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n,$$

and satisfy the inequality:

$$\operatorname{Re} \left\{ \frac{z f'(z)}{\phi(z)} e^{i\beta} \right\} > 0 \quad \left(z \in \mathcal{U}; -\frac{\pi}{2} < \beta < \frac{\pi}{2} \right)$$

for some normalized univalent and convex function ϕ . In this paper we solve the Fekete-Szegő problem for the family:

$$\mathcal{C}_1 := \bigcup_{\beta} \mathcal{C}_1(\beta) \left(-\frac{\pi}{2} < \beta < \frac{\pi}{2} \right)$$

1991 *Mathematics Subject Classification*. Primary 30C45.

Key words and phrases. Analytic functions, close-to-convex functions, convex functions, Fekete-Szegő problem, starlike functions, univalent functions.

by proving that

$$\max_{f \in \mathcal{C}_1} |a_3 - \lambda a_2^2| = \begin{cases} \frac{5}{3} - \frac{9\lambda}{4} & \left(0 \leq \lambda \leq \frac{2}{9}\right) \\ \frac{2}{3} + \frac{1}{9\lambda} & \left(\frac{2}{9} \leq \lambda \leq \frac{2}{3}\right) \\ \frac{5}{6} & \left(\frac{2}{3} \leq \lambda \leq 1\right). \end{cases}$$

1. Introduction

Let \mathcal{S} be the class of (normalized) analytic and univalent functions in the open unit disk

$$\mathcal{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$$

that are given by the Taylor series:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (z \in \mathcal{U}). \quad (1.1)$$

Let \mathcal{S}^* and \mathcal{K} denote, respectively, the subsets of \mathcal{S} consisting of starlike and convex functions in \mathcal{U} . A function f , analytic in \mathcal{U} and given by the series (1.1), is said to be close-to-convex in \mathcal{U} if

$$\operatorname{Re} \left\{ \frac{z f'(z)}{\phi(z)} e^{i\beta} \right\} > 0 \quad \left(z \in \mathcal{U}; \phi \in \mathcal{S}^*; -\frac{\pi}{2} < \beta < \frac{\pi}{2} \right). \quad (1.2)$$

We denote the family of close-to-convex functions in \mathcal{U} by \mathcal{C} . This class was introduced and studied by Kaplan [10]. The number $e^{i\beta}$ is necessary in (1.2) for the definition of close-to-convex functions.

It is well known, for $f \in \mathcal{S}$ and given by (1.1), that

$$|a_3 - a_2^2| \leq 1, \quad (1.3)$$

where the equality holds true for the Koebe function:

$$k(z) = \frac{z}{(1-z)^2} = z + \sum_{n=2}^{\infty} n z^n.$$

Since $k \in \mathcal{S}^* (\supset \mathcal{C})$, the estimate (1.3) is true for \mathcal{S}^* and \mathcal{C} without any further refinement. However, Trimble [18] has shown that, for $f \in \mathcal{K}$ and given by (1.1),

$$|a_3 - a_2^2| \leq \frac{1}{3}. \quad (1.4)$$

Earlier in 1933, Fekete and Szegő [4] made use of Löwner's parametric method in order to prove that, if $f \in \mathcal{S}$ and is given by (1.1),

$$|a_3 - \lambda a_2^2| \leq 1 + 2e - \frac{2\lambda}{1-\lambda} \quad (0 \leq \lambda \leq 1). \quad (1.5)$$

Equality in (1.5) holds true for the Koebe function $k(z)$ only for $\lambda = 0$ and $\lambda = 1$. The case $0 < \lambda < 1$ provides an example of an extremal problem over \mathcal{S} in which the Koebe function $k(z)$ fails to be extremal. The determination of sharp upper bound for the nonlinear functional $|a_3 - \lambda a_2^2|$

for any given family \mathcal{F} of normalized analytic functions is popularly known as the Fekete-Szegő problem for \mathcal{F} . The result (1.5) was also proved later by Goluzin [7], Jenkins [9], Pfluger [14], Shaffer and Spencer [15], and others.

The Fekete-Szegő problem for the families \mathcal{K} , \mathcal{S}^* , and \mathcal{C} has been completely solved in the literature. Thus we have (*cf.* [10] and [11])

$$\max_{f \in \mathcal{K}} |a_3 - \lambda a_2^2| = \max \left\{ \frac{1}{3}, |\lambda - 1| \right\},$$

$$\max_{f \in \mathcal{S}^*} |a_3 - \lambda a_2^2| = \begin{cases} |3 - 4\lambda| & \left(\lambda \leq \frac{1}{2} \text{ and } \lambda \leq 1 \right) \\ 1 & \left(\frac{1}{2} \leq \lambda \leq 1 \right), \end{cases}$$

and

$$\max_{f \in \mathcal{C}} |a_3 - \lambda a_2^2| = \begin{cases} 3 - 4\lambda & \left(0 \leq \lambda \leq \frac{1}{3} \right) \\ \frac{1}{3} + \frac{4}{9\lambda} & \left(\frac{1}{3} \leq \lambda \leq \frac{2}{3} \right) \\ 1 & \left(\frac{2}{3} \leq \lambda \leq 1 \right). \end{cases}$$

Many other recent works on the Fekete-Szegő problem include, for example, [1], [3], [6], [13], [14], and [17].

We now introduce the class $\mathcal{C}_1(\beta)$ of (normalized) analytic functions f in \mathcal{U} , which are given by (1.1) and satisfy the inequality (1.2) with $\phi \in \mathcal{K}$ (instead of $\phi \in \mathcal{S}^*$), and let

$$\mathcal{C}_1 := \bigcup_{\beta} \mathcal{C}_1(\beta) \quad \left(-\frac{\pi}{2} < \beta < \frac{\pi}{2} \right).$$

Since $\mathcal{K} \subset \mathcal{S}^*$, it follows that $\mathcal{C}_1 \subset \mathcal{C}$. Also, by taking $f = \phi$ in (1.2), we have $\mathcal{K} \subset \mathcal{C}_1$. Furthermore, the choice $\phi(z) = z$ in (1.2) exhibits the fact that the class of (normalized) analytic functions satisfying the inequality:

$$\operatorname{Re} \left\{ e^{i\beta} f'(z) \right\} > 0 \quad \left(z \in \mathcal{U}; -\frac{\pi}{2} < \beta < \frac{\pi}{2} \right)$$

is contained in the class \mathcal{C}_1 .

Problems involving growth and distortion inequalities, coefficient estimates, convex hull, extreme points, and so on, for the family \mathcal{C}_1 were investigated by Silverman and Telage [16]. In this paper, we completely solve the Fekete-Szegő problem for the family \mathcal{C}_1 . In particular, one of our results (Theorem 4 below) gives a refinement of (1.3) for the smaller set \mathcal{C}_1 , and it also includes some recent results of Abdel-Gawad and Thomas [2].

Theorem 1. *Let $f \in \mathcal{C}_1$ and be given by (1.1). Then*

$$\left| a_3 - \frac{2}{9} a_2^2 \right| \leq \frac{7}{6}$$

The result is sharp.

Theorem 2. Let $f \in \mathcal{C}_1$ and be given by (1.1). Then

$$|a_3 - \lambda a_2^2| \leq \frac{5}{3} - \frac{9\lambda}{4} \quad \left(\lambda \leq \frac{2}{9} \right).$$

The result is sharp.

Theorem 3. Let $f \in \mathcal{C}_1$ and be given by (1.1). Then

$$|a_3 - \lambda a_2^2| \leq \frac{2}{3} + \frac{1}{9\lambda} \quad \left(\frac{2}{9} < \lambda < \frac{2}{3} \right).$$

The result is sharp.

Theorem 4. Let $f \in \mathcal{C}_1$ and be given by (1.1). Then

$$\left| a_3 - \frac{2}{3} a_2^2 \right| \leq \frac{5}{6}.$$

The result is sharp.

Theorem 5. Let $f \in \mathcal{C}_1$ and be given by (1.1). Then

$$|a_3 - a_2^2| \leq \frac{5}{6}.$$

The result is sharp.

Theorem 6. Let $f \in \mathcal{C}_1$ and be given by (1.1). Then

$$|a_3 - \lambda a_2^2| \leq \frac{5}{6} \quad \left(\frac{2}{3} \leq \lambda \leq 1 \right).$$

The result is sharp.

It follows from the definition that, if $f \in \mathcal{C}_1$, then f' can be written as

$$f'(z) = \frac{\phi(z)}{z} h(z) e^{-i\beta} \quad \left(z \in \mathcal{U}; \phi \in \mathcal{K}; -\frac{\pi}{2} < \beta < \frac{\pi}{2} \right), \quad (1.6)$$

where

$$h(z) = \left(\frac{1+w(z)}{1-w(z)} \right) \cos \beta + i \sin \beta$$

for some Schwarz function w analytic in \mathcal{U} such that

$$w(0) = 0 \quad \text{and} \quad |w(z)| < 1 \quad (z \in \mathcal{U}). \quad (1.7)$$

Thus, if f is given by (1.1),

$$\phi(z) = z + \sum_{n=2}^{\infty} b_n z^n \quad (z \in \mathcal{U}) \quad (1.8)$$

and

$$w(z) = \sum_{n=1}^{\infty} \alpha_n z^n \quad (z \in \mathcal{U}) \quad (1.9)$$

then

$$2a_2 = \left(2e^{-i\beta} \cos \beta\right) a_1 + b_2 \quad (1.10)$$

and

$$3a_3 = 2e^{-i\beta} (\alpha_2 + \alpha_1^2 + \alpha_1 b_2) \cos \beta + b_3. \quad (1.11)$$

Equations (1.9) and (1.10), together, yield

$$\begin{aligned} a_3 - \lambda a_2^2 &= \frac{1}{3} \left(b_3 - \frac{3}{4} \lambda b_2^2 \right) + \frac{2}{3} e^{-i\beta} \left[\alpha_2 + \left(1 - \frac{3}{2} \lambda e^{-i\beta} \cos \beta \right) \alpha_1^2 \right] \cos \beta \\ &\quad + \frac{2}{3} e^{-i\beta} \left(1 - \frac{3}{2} \lambda \right) \alpha_1 b_2 \cos \beta. \end{aligned} \quad (1.12)$$

The expression (1.12) shall be used throughout this paper. We shall also need the following results for the proof of Theorems 1 to 6.

Theorem A (Keogh and Merkes [11]). *Let ϕ given by (1.8) be a univalent convex function in \mathcal{U} (that is, $\phi \in \mathcal{K}$). Then, for any complex number s ,*

$$|b_3 - s b_2^2| \leq \max \left\{ \frac{1}{3}, |s - 1| \right\}. \quad (1.13)$$

Theorem B (Keogh and Merkes [11]). *Let the Schwarz function w be given by (1.7) and the power series (1.9). Then, for any complex number s ,*

$$|\alpha_2 - s \alpha_1^2| \leq 1 + (|s| - 1) |\alpha_1|^2. \quad (1.14)$$

Theorem C (Silverman and Telage [16]). *Let $f \in \mathcal{C}_1$ and be given by (1.1). Then*

$$|a_n| \leq 2 - \frac{1}{n} \quad (n \in \mathbb{N} \setminus \{1\}). \quad (1.15)$$

Equality in (1.15) holds true for the function:

$$h(z, \zeta) = \bar{\zeta} \log(1 - \zeta z) + \frac{2z}{1 - \zeta z} \quad (|\zeta| = 1).$$

2. Proofs of Theorems 1 to 6

Proof of Theorem 1. Putting $\lambda = \frac{2}{9}$ in (1.12), we get

$$\left| a_3 - \frac{2}{9} a_2^2 \right| \leq \frac{1}{3} \left| b_3 - \frac{1}{6} b_2^2 \right| + \left\{ \frac{2}{3} \left| \alpha_2 - \left(\frac{1}{3} e^{-i\beta} \cos \beta - 1 \right) \alpha_1^2 \right| + \frac{4}{9} |\alpha_1 b_2| \right\} \cos \beta$$

Using Theorem A, Theorem B, and the inequality $|b_2| \leq 1$, we get

$$\begin{aligned} \left| a_3 - \frac{2}{9} a_2^2 \right| &\leq \frac{5}{18} + \left\{ \frac{2}{3} \left[1 + \left(\left| \frac{1}{3} (\cos \beta - i \sin \beta) \cos \beta - 1 \right| - 1 \right) |\alpha_1|^2 \right] \right. \\ &\quad \left. + \frac{4}{9} |\alpha_1 b_2| \right\} \cos \beta \\ &\leq \frac{17}{18} + \left\{ \frac{2}{3} \left(\sqrt{1 - \frac{5}{9} \cos^2 \beta} - 1 \right) |\alpha_1|^2 + \frac{4}{9} |\alpha_1 b_2| \right\} \cos \beta. \end{aligned} \quad (2.1)$$

Since

$$\sqrt{1 - \frac{5}{9} \cos^2 \beta} - 1 \leq \frac{2}{3} - 1 = -\frac{1}{3},$$

we find from (2.1) that

$$\begin{aligned} \left| a_3 - \frac{2}{9} a_2^2 \right| &\leq \frac{17}{18} + \left\{ \frac{2}{3} \left(-\frac{1}{3} \right) |\alpha_1|^2 + \frac{4}{9} |\alpha_1 b_2| \right\} \cos \beta \\ &= \frac{17}{18} + \frac{2}{9} |b_2|^2 \cos \beta - \left\{ \frac{2}{9} |b_2|^2 + \frac{2}{9} |\alpha_1|^2 - \frac{4}{9} |\alpha_1 b_2| \right\} \cos \beta \\ &\leq \frac{7}{6} - \frac{2}{9} (|b_2| - |\alpha_1|)^2 \cos \beta \\ &\leq \frac{7}{6}, \end{aligned}$$

which completes the proof of Theorem 1.

Proof of Theorem 2. We begin by considering

$$\begin{aligned} |a_3 - \lambda a_2^2| &= \left| a_3 - \frac{2}{9} a_2^2 + \frac{2}{9} a_2^2 - \lambda a_2^2 \right| \\ &\leq \left| a_3 - \frac{2}{9} a_2^2 \right| + \left(\frac{2}{9} - \lambda \right) |a_2|^2, \end{aligned}$$

which, in view of Theorem 1, yields

$$\left| a_3 - \frac{2}{9} a_2^2 \right| \leq \frac{7}{6}.$$

Thus, using Theorem C, we have

$$\begin{aligned} |a_3 - \lambda a_2^2| &\leq \frac{7}{6} + \left(\frac{2}{9} - \lambda \right) \left(\frac{3}{2} \right)^2 \\ &= \frac{5}{3} - \frac{9\lambda}{4}, \end{aligned}$$

which completes the proof of Theorem 2.

Equality can be shown to hold true by putting $b_2 = b_3 = 1$, $\alpha_1 = 1$, $\alpha_2 = 0$, and $\beta = 0$ in (1.12).

Proof of Theorem 3. By (1.12), we have

$$\begin{aligned} |a_3 - \lambda a_2^2| &\leq \frac{1}{3} \left| b_3 - \frac{3}{4} \lambda b_2^2 \right| + \frac{2}{3} \left| \alpha_2 - \left(\frac{3}{2} \lambda e^{-i\beta} \cos \beta - 1 \right) \alpha_1^2 \right| \cos \beta \\ &\quad + \frac{2}{3} \left(1 - \frac{3}{2} \lambda \right) |\alpha_1| |b_2| \cos \beta. \end{aligned} \tag{2.2}$$

Now, using Theorem A, Theorem B, and the fact that $|b_2| \leq 1$, we get

$$\begin{aligned} |a_3 - \lambda a_2^2| &\leq \frac{4-3\lambda}{12} + \frac{2}{3} \left\{ 1 + \left(\left| \frac{3}{2} \lambda e^{-i\beta} \cos \beta - 1 \right| - 1 \right) |\alpha_1|^2 \right\} \cos \beta + \frac{2-3\lambda}{3} |\alpha_1| \cos \beta \\ &= \frac{4-3\lambda}{12} + \left\{ \frac{2}{3} + \frac{2}{3} |\alpha_1|^2 \left(\sqrt{1 - \left(3\lambda - \frac{9}{4} \lambda^2 \right) \cos^2 \beta} - 1 \right) \cos \beta \right. \\ &\quad \left. + \frac{2-3\lambda}{3} |\alpha_1| \right\} \end{aligned} \quad (2.3)$$

Putting $\cos \beta = y$ and $|\alpha_1| = \alpha$ in (2.3), we get

$$\begin{aligned} |a_3 - \lambda a_2^2| &\leq \frac{4-3\lambda}{12} + y \left\{ \frac{2}{3} + \frac{2}{3} \alpha^2 \left(\sqrt{1 - \left(3\lambda - \frac{9}{4} \lambda^2 \right) y^2} - 1 \right) + \frac{2-3\lambda}{3} \alpha \right\} \\ &= F_\lambda(\alpha, y) \quad (\text{say}). \end{aligned} \quad (2.4)$$

In the rest of the proof, we shall show that $F_\lambda(\alpha, y)$ attains its maximum value for $(\alpha, y) \in [0, 1] \times [0, 1]$ at the point $\left(\frac{2-3\lambda}{6\lambda}, 1 \right)$. Note that

$$F_\lambda \left(\frac{2-3\lambda}{6\lambda}, 1 \right) = \frac{2}{3} + \frac{1}{9\lambda}. \quad (2.5)$$

We shall first show that F_λ does not have a local maximum at any interior point of the open rectangle $(0, 1) \times (0, 1)$. For, if F_λ had a local maximum at some point $(\alpha_0, y_0) \in (0, 1) \times (0, 1)$, then the partial derivatives $\frac{\partial F_\lambda}{\partial \alpha}$ and $\frac{\partial F_\lambda}{\partial y}$ must vanish at (α_0, y_0) . Now

$$\frac{\partial F_\lambda}{\partial \alpha} = y \left[\frac{4\alpha}{3} \left(\sqrt{1 - y^2 \left(3\lambda - \frac{9\lambda^2}{4} \right)} - 1 \right) + \frac{2-3\lambda}{3} \right].$$

Thus the first requirement:

$$\frac{\partial F_\lambda}{\partial \alpha} \Big|_{(\alpha_0, y_0)} = 0$$

implies that

$$4\alpha_0 \left(\sqrt{1 - y_0^2 \left(3\lambda - \frac{9\lambda^2}{4} \right)} - 1 \right) = -\frac{4}{3} \gamma \quad \left(\gamma = \frac{3}{4}(2-3\lambda) \right).$$

Note that $0 < \gamma < 1$. Therefore

$$y_0^2 \left(3\lambda - \frac{9\lambda^2}{4} \right) = \frac{2\gamma}{3\alpha_0} - \frac{\gamma^2}{9\alpha_0^2}. \quad (2.6)$$

Similarly, we have

$$\frac{\partial F_\lambda}{\partial y} = \left[\frac{2}{3} + \frac{2\alpha^2}{3} \left(\sqrt{1 - y^2 \left(3\lambda - \frac{9\lambda^2}{4} \right)} - 1 \right) + \left(\frac{2 - 3\lambda}{3} \right) \alpha \right] + y \left[\frac{2\alpha^2}{3} \frac{(-y) \left(3\lambda - \frac{9\lambda^2}{4} \right)}{\sqrt{1 - y^2 \left(3\lambda - \frac{9\lambda^2}{4} \right)}} \right].$$

Thus the second requirement:

$$\left. \frac{\partial F_\lambda}{\partial y} \right|_{(\alpha_0, y_0)} = 0$$

implies that

$$\frac{2}{3} + \frac{2\alpha_0^2}{3} \left(\sqrt{1 - y_0^2 \left(3\lambda - \frac{9\lambda^2}{4} \right)} - 1 \right) + \left(\frac{2 - 3\lambda}{3} \right) \alpha_0 = \frac{2\alpha_0^2}{3} \frac{y_0^2 \left(3\lambda - \frac{9\lambda^2}{4} \right)}{\sqrt{1 - y_0^2 \left(3\lambda - \frac{9\lambda^2}{4} \right)}}. \quad (2.7)$$

Substituting the value of $y_0^2 \left(3\lambda - \frac{9\lambda^2}{4} \right)$ from (2.6) into (2.7), we find that

$$\frac{2}{3} + \frac{2\alpha_0^2}{3} \left(-\frac{\gamma}{3\alpha_0} \right) + \frac{4\gamma\alpha_0}{9} - \frac{2}{3}\alpha_0^2 \left[\frac{\frac{2\gamma}{3\alpha_0} - \frac{\gamma^2}{9\alpha_0^2}}{1 - \left(\frac{\gamma}{3\alpha_0} \right)} \right] = 0$$

or, equivalently, that

$$\alpha_0^2 \gamma - 3\alpha_0 + \gamma = 0. \quad (2.8)$$

Solving the quadratic equation (2.8) for α_0 , we have

$$\alpha_0 \gamma = \frac{3 - \sqrt{9 - 4\gamma^2}}{2}. \quad (2.9)$$

The value of $F_\lambda(\alpha_0, y_0)$ written in terms of γ becomes

$$F_\lambda(\alpha_0, y_0) = \frac{6 + 4\gamma}{36} + y_0 \left(\frac{2}{3} + \frac{2}{9}\gamma\alpha_0 \right),$$

which, upon substituting the value of $\alpha_0\gamma$ from (2.9), yields

$$\begin{aligned} F_\lambda(\alpha_0, y_0) &= \frac{6+4\gamma}{36} + y_0 \left(1 - \frac{1}{3} \sqrt{1 - \frac{4}{9}\gamma^2} \right) \\ &\leq \frac{6+4\gamma}{36} + 1 - \frac{1}{3} \sqrt{1 - \frac{4}{9}\gamma^2} \\ &= \frac{21 + 2\gamma - 6\sqrt{1 - \frac{4}{9}\gamma^2}}{18}. \end{aligned}$$

Since $y \in (0, 1)$, there exists $\eta > 0$ such that

$$\gamma = \frac{3}{2} \cos \delta \quad \text{and} \quad \sqrt{1 - \frac{4}{9}\gamma^2} = \sin \delta \quad \left(0 < \eta < \delta < \frac{\pi}{2} \right).$$

Moreover, the inequality:

$$1 < 2 \cos \delta + \sin \delta$$

gives

$$(1 - \cos \delta)(1 - \sin \delta) < \frac{\cos^2 \delta}{2},$$

so that

$$\begin{aligned} (1 - \cos \delta) \left(21 + 2\gamma - 6\sqrt{1 - \frac{4}{9}\gamma^2} \right) &= (1 - \cos \delta)(15 + 3 \cos \delta) + 6(1 - \cos \delta)(1 - \sin \delta) \\ &< 15 + 3 \cos \delta - 15 \cos \delta - 3 \cos^2 \delta + 3 \cos^2 \delta \\ &= 15 - 12 \cos \delta = 3 + 12(1 - \cos \delta). \end{aligned}$$

Therefore, we have

$$\frac{21 + 2\gamma - 6\sqrt{1 - \frac{4}{9}\gamma^2}}{18} < \frac{1}{6(1 - \cos \delta)} + \frac{2}{3} = \frac{2}{3} + \frac{1}{9\lambda},$$

which shows that

$$F_\lambda(\alpha_0, y_0) < \frac{2}{3} + \frac{1}{9\lambda}.$$

Hence $F_\lambda(\alpha, y)$ does not have a local maximum in $(0, 1) \times (0, 1)$, so that the maximum must be attained at a boundary point. Since

$$F_\lambda(\alpha, 0) = \frac{4 - 3\lambda}{12} < \frac{2}{3} + \frac{1}{9\lambda},$$

there is no maximum on the line $y = 0$. Similarly, we have

$$F_\lambda(0, y) = \frac{4 - 3\lambda}{12} + \frac{2}{3}y \leq \frac{4 - 3\lambda}{12} + \frac{2}{3} < \frac{2}{3} + \frac{1}{9\lambda},$$

so that there is no maximum on the line $\alpha = 0$ either. On the line $y = 1$, we get

$$F_\lambda(\alpha, 1) = 1 - \frac{\lambda}{4} - \lambda\alpha^2 + \left(\frac{2}{3} - \lambda\right)\alpha = G_\lambda(\alpha) \quad (\text{say}).$$

Putting $\alpha = 1$, we get

$$G_\lambda(1) = \frac{5}{3} - \frac{9\lambda}{4}.$$

Since $G_\lambda(1)$ is not maximal, the local maximum of $G_\lambda(\alpha)$ is attained at

$$\alpha_0 = \frac{2 - 3\lambda}{6\lambda} \quad \text{for which} \quad \left. \frac{dG_\lambda(\alpha)}{d\alpha} \right|_{\alpha=\alpha_0} = 0.$$

This leads to the maximal value given by (2.4). The proof will be complete if we show that

$$F_\lambda(1, y) \leq \frac{2}{3} + \frac{1}{9\lambda} \quad (0 < y < 1). \quad (2.10)$$

Since

$$\begin{aligned} F_\lambda(1, y) &= \frac{1}{3} \left[\frac{3 + 2\gamma}{6} + 2y \left(\sqrt{1 - y^2 \left(1 - \frac{4}{9}\gamma^2 \right)} + \frac{2\gamma}{3} \right) \right] \\ &= \frac{1}{3} \left[\frac{3 + 2\gamma}{6} + H_\lambda(y) \right], \end{aligned}$$

where

$$H_\lambda(y) = 2y \left(\sqrt{1 - y^2 \left(1 - \frac{4}{9}\gamma^2 \right)} + \frac{2\gamma}{3} \right),$$

the assertion (2.10) is equivalent to

$$H_\lambda(y) \leq \frac{2(3 - \gamma)^2}{3(3 - 2\gamma)} \quad (0 < y < 1). \quad (2.11)$$

Therefore, it suffices to prove (2.11) at the points $y \in (0, 1)$ for which

$$\frac{dH_\lambda(y)}{dy} = 0,$$

which implies that

$$\frac{2\gamma}{3} \sqrt{1 - y^2 \left(1 - \frac{4}{9}\gamma^2 \right)} = 2y^2 \left(1 - \frac{4}{9}\gamma^2 \right) - 1. \quad (2.12)$$

Squaring both sides in (2.12), we get

$$\left\{ 2y^2 \left(1 - \frac{4}{9}\gamma^2 \right) \right\}^2 - 2 \left\{ 2y^2 \left(1 - \frac{4}{9}\gamma^2 \right) \right\} \left(1 - \frac{\gamma^2}{9} \right) + \left(1 - \frac{4}{9}\gamma^2 \right) = 0.$$

Thus we have

$$2y^2 \left(1 - \frac{4}{9}\gamma^2 \right) = \frac{9 - \gamma^2 - \gamma\sqrt{18 + \gamma^2}}{9}, \quad (2.13)$$

since $0 \leq y \leq 1$.

We first square both sides of the inequality (2.11), substitute the value of y^2 from (2.13), and get the equivalent inequality :

$$(3 - 2\gamma) \left(9 - \gamma^2 - \gamma\sqrt{18 + \gamma^2}\right) \left(\sqrt{18 + \gamma^2} + 5\gamma\right)^2 \leq 8(3 + 2\gamma)(3 - \gamma)^4.$$

For simplicity of our calculations, we put $t = \sqrt{18 + \gamma^2}$; then (after a routine calculation) this becomes

$$18t\gamma(3 - 2\gamma) (2 - \gamma^2) \leq 729 - 486\gamma - 270\gamma^2 + 324\gamma^3 - 30\gamma^4 - 28\gamma^5. \quad (2.14)$$

The right-hand side of (2.14) turns out to be positive, since

$$\begin{aligned} &729 - 486\gamma - 270\gamma^2 + 324\gamma^3 - 30\gamma^4 - 28\gamma^5 \\ &= (1 - \gamma) (729 + 243\gamma - 27\gamma^2 + 297\gamma^3 + 267\gamma^4) + 239\gamma^5 \\ &> 0. \end{aligned}$$

Thus, squaring (2.14) once again, we get the equivalent inequality:

$$\begin{aligned} &324 (18 + \gamma^2) \gamma^2 (36 - 48\gamma - 20\gamma^2 + 48\gamma^3 - 7\gamma^4 - 12\gamma^5 + 4\gamma^6) \\ &\leq 729^2 - 2 \cdot 729 \cdot 486\gamma + (486^2 - 2 \cdot 729 \cdot 270) \gamma^2 + (2 \cdot 729 \cdot 342 + 2 \cdot 486 \cdot 270) \gamma^3 \\ &\quad + (270^2 - 2 \cdot 729 \cdot 30 - 2 \cdot 486 \cdot 324) \gamma^4 + (-2) \cdot 729 \cdot 28 \\ &\quad + 2 \cdot 486 \cdot 30 - 2 \cdot 270 \cdot 324) \gamma^5 + (324^2 + 2 \cdot 486 \cdot 28 + 2 \cdot 270 \cdot 30) \gamma^6 \\ &\quad + (2 \cdot 270 \cdot 28 - 2 \cdot 324 \cdot 30) \gamma^7 + (900 - 2 \cdot 324 \cdot 28) \gamma^8 + 60 \cdot 28\gamma^9 + 784\gamma^{10}. \end{aligned}$$

that is,

$$\begin{aligned} &324\gamma^2 (648 - 864\gamma - 324\gamma^2 + 816\gamma^3 - 146\gamma^4 - 168\gamma^5 + 65\gamma^6 - 12\gamma^7 + 4\gamma^8) \\ &\leq 729^2 - 2 \cdot 729 \cdot 486\gamma + (486^2 - 2 \cdot 729 \cdot 270) \gamma^2 + (2 \cdot 729 \cdot 324 + 2 \cdot 486 \cdot 270) \gamma^3 \\ &\quad + (270^2 - 2 \cdot 729 \cdot 30 - 2 \cdot 486 \cdot 324) \gamma^4 + (-2 \cdot 729 \cdot 28 \\ &\quad + 2 \cdot 486 \cdot 30 - 2 \cdot 270 \cdot 324) \gamma^5 + (324^2 + 2 \cdot 486 \cdot 28 + 2 \cdot 270 \cdot 30) \gamma^6 \\ &\quad + (2 \cdot 270 \cdot 28 - 2 \cdot 324 \cdot 30) \gamma^7 + (900 - 2 \cdot 324 \cdot 28) \gamma^8 + 60 \cdot 28\gamma^9 + 784\gamma^{10}. \end{aligned}$$

Thus the reformulated inequality is given by

$$\begin{aligned} &729^2 - 2 \cdot 729 \cdot 486\gamma + (486^2 - 2 \cdot 729 \cdot 270 - 324 \cdot 648) \gamma^2 \\ &\quad + (2 \cdot 729 \cdot 324 + 2 \cdot 486 \cdot 270 + 324 \cdot 864) \gamma^3 + (270^2 - 2 \cdot 729 \cdot 30 \\ &\quad - 2 \cdot 486 \cdot 324 + 324^2) \gamma^4 + (-2) \cdot 729 \cdot 28 + 2 \cdot 486 \cdot 30 - 2 \cdot 270 \cdot 324 \\ &\quad - 324 \cdot 816\gamma^5 + (324^2 + 2 \cdot 486 \cdot 28 + 2 \cdot 270 \cdot 30 + 324 \cdot 146) \gamma^6 \\ &\quad + (2 \cdot 270 \cdot 28 - 2 \cdot 324 \cdot 30 + 324 \cdot 168) \gamma^7 + (900 - 2 \cdot 324 \cdot 28 - 324 \cdot 65) \gamma^8 \\ &\quad + (1680 + 324 \cdot 12) \gamma^9 + (784 - 324 \cdot 4) \gamma^{10} \geq 0. \end{aligned} \quad (2.15)$$

Now the left-hand side of (2.14) is

$$\begin{aligned}
& 531441 - 708588\gamma - 367416\gamma^2 + 1014768\gamma^3 - 180792\gamma^4 - 451008\gamma^5 \\
& \quad + 195696\gamma^6 + 50112\gamma^7 - 38304\gamma^8 + 5568\gamma^9 - 512\gamma^{10} \\
& = 531441(1 - \gamma) - 171147\gamma(1 - \gamma) - 544563\gamma^2(1 - \gamma) + 470205\gamma^3(1 - \gamma) \\
& \quad + 289413\gamma^4(1 - \gamma) - 161595\gamma^5(1 - \gamma) + 34101\gamma^6(1 - \gamma) + 84213\gamma^7(1 - \gamma) \\
& \quad + 45909\gamma^8(1 - \gamma) + 51477\gamma^9(1 - \gamma) + 50965\gamma^{10} \\
& = (1 - \gamma)(531441 - 177147\gamma - 544563\gamma^2 + 470205\gamma^3 + 289413\gamma^4 - 16595\gamma^5 \\
& \quad + 34101\gamma^6 + 84213\gamma^7 + 45909\gamma^8 + 51477\gamma^9 + 50965\gamma^{10}) \\
& = (1 - \gamma) \left[(1 - \gamma)(531441 + 354294\gamma - 190269\gamma^2 + 161595\gamma^4) + 279936\gamma^3 \right. \\
& \quad \left. + 127818\gamma^4 + 34101\gamma^6 + 84213\gamma^7 + 45909\gamma^8 + 51477\gamma^9 \right] + 50965\gamma^{10} \\
& = (1 - \gamma) \left[(1 - \gamma) \{ 531441 + 164025\gamma + 190260\gamma(1 - \gamma) + 161595\gamma^4 \} \right. \\
& \quad \left. + 279936\gamma^3 + 127818\gamma^4 + 34101\gamma^6 + 84213\gamma^7 + 45909\gamma^9 + 51477\gamma^9 \right] + 50965\gamma^{10} \\
& > 0.
\end{aligned}$$

Hence (2.15) is true. This completes the proof of the main assertion of Theorem 3. The result can be shown to be sharp by setting $b_2 = b_3 = 1$, $\beta = 0$, $\alpha_1 = \frac{2 - 3\lambda}{6\lambda}$, and $\alpha_2 = 1 - \alpha_1^2$ in (1.12).

Proof of Theorem 4. Putting $\lambda = \frac{2}{3}$ in (1.12), we get

$$a_3 - \frac{2}{3}a_2^2 = \frac{1}{3} \left(b_3 - \frac{1}{2}b_2^2 \right) + \frac{2}{3}e^{-i\beta} \left[\alpha_2 + \left(1 - e^{-i\beta} \cos \beta \right) \alpha_1^2 \right] \cos \beta.$$

Hence

$$\left| a_3 - \frac{2}{3}a_2^2 \right| \leq \frac{1}{3} \left| b_3 - \frac{1}{2}b_2^2 \right| + \frac{2}{3} \left| \alpha_2 - \left(e^{-i\beta} \cos \beta - 1 \right) \alpha_1^2 \right| \cos \beta.$$

Using Theorem A and Theorem B, we get

$$\begin{aligned}
\left| a_3 - \frac{2}{3}a_2^2 \right| & \leq \frac{1}{3} \cdot \frac{1}{2} + \frac{2}{3} \left[1 + \left(\left| e^{-i\beta} \cos \beta - 1 \right| - 1 \right) |\alpha_1|^2 \right] \cos \beta \\
& = \frac{1}{6} + \frac{2}{3} \left[1 + \left(\sqrt{1 - \cos^2 \beta} - 1 \right) |\alpha_1|^2 \right] \cos \beta \\
& = \frac{1}{6} + \frac{2}{3} \cos \beta + \frac{2}{3} (|\sin \beta| - 1) |\alpha_1|^2 \cos \beta \\
& \leq \frac{5}{6} - \frac{2}{3} (1 - |\sin \beta|) |\alpha_1|^2 \cos \beta \\
& \leq \frac{5}{6},
\end{aligned}$$

which completes the proof of Theorem 4.

Proof of Theorem 5. Putting $\lambda = 1$ in (1.12), we get

$$a_3 - a_2^2 = \frac{1}{3} \left(b_3 - \frac{3}{4} b_2^2 \right) + \frac{2}{3} e^{-i\beta} \left[\alpha_2 + \left(1 - \frac{3}{2} e^{-i\beta} \cos \beta \right) \alpha_1^2 \right] \cos \beta - \frac{1}{3} \alpha_1 b_2 e^{-i\beta} \cos \beta.$$

Hence

$$|a_3 - a_2^2| \leq \frac{1}{3} |b_3 - b_2^2| + \frac{|b_2|^2}{12} + \frac{2}{3} \left| \alpha_2 - \left(\frac{3}{2} e^{-i\beta} \cos \beta - 1 \right) \alpha_1^2 \right| + \frac{|\alpha_1 b_2|}{3} \cos \beta. \quad (2.16)$$

Using Theorem B and a result of Trimble [18], we find from (2.16) that

$$\begin{aligned} |a_3 - a_2^2| &\leq \frac{1}{3} \left(\frac{1 - |b_2|^2}{3} \right) + \frac{|b_2|^2}{12} + \frac{2}{3} + \frac{2}{3} \left| \left(\frac{3}{2} \cos^2 \beta - 1 - \frac{3}{2} i \sin \beta \cos \beta \right) \alpha_1^2 + \frac{|\alpha_1 b_2|}{3} \right| \\ &= \frac{7}{9} + \frac{|b_2|^2}{18} - \frac{|b_2|^2}{12} + \frac{2}{3} \left(\sqrt{1 - \frac{3}{4} \cos^2 \beta} - 1 \right) |\alpha_1|^2 + \frac{|\alpha_1 b_2|}{3} \\ &= \frac{7}{9} + \frac{|b_2|^2}{18} - \frac{1}{3} \left[\frac{|b_2|^2}{4} - 2 \left(\sqrt{1 - \frac{3}{4} \cos^2 \beta} - 1 \right) |\alpha_1|^2 - |\alpha_1 b_2| \right]. \end{aligned} \quad (2.17)$$

The elementary inequality:

$$1 - \frac{3}{4} \cos^2 \beta \geq \frac{1}{4}$$

immediately yields

$$\sqrt{1 - \frac{3}{4} \cos^2 \beta} \geq \frac{1}{2}.$$

Thus we find from (2.17) that

$$\begin{aligned} |a_3 - a_2^2| &\leq \frac{7}{9} + \frac{1}{18} - \frac{1}{3} \left(\frac{|b_2|^4}{4} + |\alpha_1|^2 - |\alpha_1 b_2| \right) \\ &= \frac{5}{6} - \frac{1}{3} \left(\frac{|b_2|}{2} - |\alpha_1| \right)^2 \\ &\leq \frac{5}{6}. \end{aligned}$$

This completes the proof of Theorem 5.

Proof of Theorem 6. Observe that

$$a_3 - \lambda a_2^2 = (3\lambda - 2) (a_3 - a_2^2) + 3(1 - \lambda) \left(a_3 - \frac{2}{3} a_2^2 \right).$$

The main assertion of Theorem 6 follows from Theorem 4 and Theorem 5.

Equality can be shown to hold true by setting $b_2 = b_3 = 1$, $\alpha_2 = 1 - \alpha_1^2$, $\beta = 0$, and

$$\alpha_1 = \frac{(2 - 3\lambda) \pm i\sqrt{6\lambda - 4}}{6\lambda}$$

in (1.12).

Acknowledgements

The present investigation was supported, in part, by the *Natural Sciences and Engineering Research Council of Canada* under Grant OGP0007353.

REFERENCES

- [1] O.P. Ahuja and M.Jahangiri, Fekete-Szegő problem for a unified class of analytic functions, *Pan Amer. Math. J.* **7**(2)(1997), 67-78.
- [2] H.R. Abdel-Gawad and D.K. Thomas, A subclass of close-to-convex functions, *Publ. Inst. Math. (Beograd)(N.S.)* **49**(63)(1991), 61-66.
- [3] H.R. Abdel-Gawad and D.K. Thomas, The Fekete-Szegő problem for strongly close-to-convex functions, *Proc. Amer. Math. Soc.* **114**(1992), 345-349.
- [4] A. Chonweerayoot, D.K. Thomas, and W. Upakarnitikaset, On the Fekete-Szegő theorem for close-to-convex functions, *Publ. Inst. Math. (Beograd)(N.S.)* **52**(66)(1992), 18-26.
- [5] M. Fekete and G. Szegő, Eine Bemerkung über ungerade schlichte Funktionen, *J. London Math. Soc.* **8**(1933), 85-89.
- [6] Chun-Yi Gao, The Fekete-Szegő problem for a class of close-to-convex functions, *Chinese Ann. Math. Ser. A* **15**(1994), 650-656.
- [7] G.M. Goluzin, Some questions in the theory of univalent functions, *Trudy Mat. Inst. Steklova* **27**(1949), 1-112 (in Russian).
- [8] A.W. Goodman and E.B. Saff, On the definition of close-to-convex functions, *Internat. J. Math. and Math. Sci.* **1**(1978), 125-128.
- [9] J.A. Jenkins, On certain coefficients of univalent functions, in *Analytic Functions*, Princeton Univ. Press, Princeton, New Jersey, 1960, 159-194.
- [10] W. Kaplan, Close-to-convex schlicht functions, *Michigan Math. J.* **1**(1952), 169-185.
- [11] F.R. Keogh and E.P. Merkes, A coefficient inequality for certain classes of analytic functions, *Proc. Amer. Math. Soc.* **20**(1969), 8-12.
- [12] W. Koepf, On the Fekete-Szegő problem for close-to-convex functions. I and II, *Proc. Amer. Math. Soc.* **101**(1987), 89-95; *Arch. Math. (Basel)* **49**(1987), 420-433.
- [13] W. Ma and D. Minda, Uniformly convex functions. II, *Ann. Polon. Math.* **58**(1993), 275-285.
- [14] A. Pfluger, The Fekete-Szegő inequality by a variational method, *Ann. Acad. Sci. Fenn. Ser. AI* **10**(1984).
- [15] A.C. Shaeffer and D.C. Spencer, The coefficients of schlicht functions, *Duke Math. J.* **10**(1943), 611-635.
- [16] H. Silverman and D.N. Telage, Extreme points of subclasses of close-to-convex functions, *Proc. Amer. Math. Soc.* **74**(1979), 59-65.
- [17] D.K. Thomas, The Fekete-Szegő problem for strongly close-to-convex functions, in *Topics in Univalent Functions and Its Applications* (Kyoto, 1989), Sūrikaiseikikenkyūsho Kōkyūroku **714**(1990), 11-17.
- [18] S.Y. Trimble, A coefficient inequality for convex univalent functions, *Proc. Amer. Math. Soc.* **48**(1975), 266-267.