

ON NEW SUBCLASSES OF ANALYTIC
AND UNIVALENT FUNCTIONS*

by

SHIGEYOSHI OWA

DM-343-IR

JANUARY 1985

* This research was completed at the University of Victoria while the author was on study leave from Kinki University, Osaka, Japan.

1980 Mathematics Subject Classification. Primary 26A24, 30C45.

ABSTRACT

The object of the present paper is to prove several interesting results involving coefficient inequalities, distortion theorems, starlikeness and convexity for functions belonging to the subclasses $\mathcal{T}^*(\alpha, \lambda)$ and $\mathcal{L}(\alpha, \lambda)$ of analytic and univalent functions, which are introduced here in terms of fractional calculus. Some theorems for the classes $\mathcal{T}^*(\alpha, \lambda, k)$ and $\mathcal{L}(\alpha, \lambda, k)$ of functions with fixed finitely many coefficients are also proved, and the classes of univalent functions with quasiconformal extension are considered.

1. INTRODUCTION

Let \mathcal{A} denote the class of functions of the form

$$(1.1) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

which are analytic in the unit disk $\mathcal{U} = \{z: |z| < 1\}$. Further, let \mathcal{S} be the subclass of \mathcal{A} consisting of analytic and univalent functions in the unit disk \mathcal{U} . Then a function $f(z)$ of \mathcal{S} is said to be starlike of order α if and only if

$$(1.2) \quad \operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) > \alpha \quad (z \in \mathcal{U})$$

for some α ($0 \leq \alpha < 1$). We denote by $\mathcal{S}^*(\alpha)$ the class of all starlike functions of order α .

A function $f(z)$ belonging to the class \mathcal{S} is said to be convex of order α if and only if

$$(1.3) \quad \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > \alpha \quad (z \in \mathcal{U})$$

for some α ($0 \leq \alpha < 1$). We denote by $\mathcal{K}(\alpha)$ the class of all convex functions of order α . We note that $f(z) \in \mathcal{K}(\alpha)$ if and only if $zf'(z) \in \mathcal{S}^*(\alpha)$, and that

$$\mathcal{S}^*(\alpha) \subset \mathcal{S}^*(0) \equiv \mathcal{S}^*, \quad \mathcal{K}(\alpha) \subset \mathcal{K}(0) \equiv \mathcal{K}, \quad \text{and} \quad \mathcal{K}(\alpha) \subset \mathcal{S}^*(\alpha)$$

for $0 \leq \alpha < 1$.

The classes $\mathcal{S}^*(\alpha)$ and $\mathcal{K}(\alpha)$ were introduced by Robertson [10], and were studied subsequently by Schild [13], MacGregor [4], Pinchuk [9], and others.

Many essentially equivalent definitions of the fractional calculus (that is, fractional derivatives and fractional integrals) have been given in the literature (cf., e.g., [2, Chapter 13], [6], [7], [11], [12], [15, p. 28 et seq.], [17], and [18]). We find it to be convenient to recall here the following definitions which were used recently by Owa [8] (and by Srivastava and Owa [16]).

DEFINITION 1. The fractional integral of order λ is defined, for a function $f(z)$, by

$$(1.4) \quad D_z^{-\lambda} f(z) = \frac{1}{\Gamma(\lambda)} \int_0^z \frac{f(\zeta)}{(z-\zeta)^{1-\lambda}} d\zeta,$$

where $\lambda > 0$, $f(z)$ is an analytic function in a simply-connected region of the z -plane containing the origin, and the multiplicity of $(z-\zeta)^{\lambda-1}$ is

removed by requiring $\log(z-\zeta)$ to be real when $z - \zeta > 0$.

DEFINITION 2. The fractional derivative of order λ is defined, for a function $f(z)$, by

$$(1.5) \quad D_z^\lambda f(z) = \frac{1}{\Gamma(1-\lambda)} \frac{d}{dz} \int_0^z \frac{f(\zeta)}{(z-\zeta)^\lambda} d\zeta,$$

where $0 \leq \lambda < 1$, $f(z)$ is an analytic function in a simply-connected region of the z -plane containing the origin, and the multiplicity of $(z-\zeta)^{-\lambda}$ is removed as in Definition 1 above.

DEFINITION 3. Under the hypotheses of Definition 2, the fractional derivative of order $n + \lambda$ is defined by

$$(1.6) \quad D_z^{n+\lambda} f(z) = \frac{d^n}{dz^n} D_z^\lambda f(z),$$

where $0 \leq \lambda < 1$, $n \in \mathcal{N}_0 = \{0, 1, 2, \dots\}$.

Let $\mathcal{S}^*(\alpha, \lambda)$ denote the class of all functions $f(z)$ in \mathcal{S} satisfying the inequality

$$(1.7) \quad \operatorname{Re} \left(\frac{\Gamma(1-\lambda) z^{1+\lambda} D_z^{1+\lambda} f(z)}{f(z)} \right) > \alpha \quad (z \in \mathcal{U})$$

for $\lambda < 1$ and $0 \leq \alpha < 1$. Also let $\mathcal{H}(\alpha, \lambda)$ denote the class of all functions $f(z)$ in \mathcal{S} such that

$$(1.8) \quad \Gamma(1-\lambda) z^{1+\lambda} D_z^{1+\lambda} f(z) \in \mathcal{S}^*(\alpha, \lambda)$$

for $\lambda < 1$ and $0 \leq \alpha < 1$. Clearly,

$$\mathcal{S}^*(\alpha, 0) = \mathcal{S}^*(\alpha) \quad \text{and} \quad \mathcal{K}(\alpha, 0) = \mathcal{K}(\alpha)$$

where we have set $\lambda = 0$. Thus $\mathcal{S}^*(\alpha, \lambda)$ and $\mathcal{K}(\alpha, \lambda)$ are generalizations of the classes $\mathcal{S}^*(\alpha)$ and $\mathcal{K}(\alpha)$, respectively.

Let \mathcal{T} be the subclass of \mathcal{S} consisting of all functions of the form

$$(1.9) \quad f(z) = z - \sum_{n=2}^{\infty} a_n z^n \quad (a_n \geq 0).$$

Denote by $\mathcal{T}^*(\alpha, \lambda)$ and $\mathcal{C}(\alpha, \lambda)$ the classes obtained by taking intersections, respectively, of the classes $\mathcal{S}^*(\alpha, \lambda)$ and $\mathcal{K}(\alpha, \lambda)$ with \mathcal{T} , that is,

$$(1.10) \quad \mathcal{T}^*(\alpha, \lambda) = \mathcal{S}^*(\alpha, \lambda) \cap \mathcal{T}$$

and

$$(1.11) \quad \mathcal{C}(\alpha, \lambda) = \mathcal{K}(\alpha, \lambda) \cap \mathcal{T}.$$

The special cases of the classes $\mathcal{T}^*(\alpha, \lambda)$ and $\mathcal{C}(\alpha, \lambda)$, when $\lambda = 0$, were studied by Silverman [14]. Thus the classes $\mathcal{T}^*(\alpha, \lambda)$ and $\mathcal{C}(\alpha, \lambda)$ provide interesting generalizations of the classes considered by Silverman [14].

In this paper, we first prove several interesting results for functions belonging to the general classes $\mathcal{T}^*(\alpha, \lambda)$ and $\mathcal{C}(\alpha, \lambda)$. We then introduce the subclasses $\mathcal{T}^*(\alpha, \lambda, k)$ and $\mathcal{C}(\alpha, \lambda, k)$ of functions with fixed finitely many coefficients. And, finally, we consider the classes of univalent functions with quasiconformal extension.

2. COEFFICIENT INEQUALITIES

THEOREM 1. Let the function $f(z)$ be defined by (1.1). If

$$(2.1) \quad \sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} |a_n| \leq 1 - \alpha$$

for $\lambda < 1$ and $0 \leq \alpha < 1$, then $f(z) \in \mathcal{S}^*(\alpha, \lambda)$. The result (2.1) is sharp.

PROOF. We need only prove that (2.1) implies (1.7). In order to prove (1.7), it suffices to show that

$$(2.2) \quad \left| \frac{\Gamma(1-\lambda) z^{1+\lambda} D_z^{1+\lambda} f(z)}{f(z)} - 1 \right| < 1 - \alpha \quad (z \in \mathcal{U}).$$

Note that

$$(2.3) \quad D_z^{1+\lambda} f(z) = \frac{1}{\Gamma(1-\lambda)} z^{-\lambda} + \sum_{n=2}^{\infty} \frac{\Gamma(n+1)}{\Gamma(n-\lambda)} a_n z^{n-1-\lambda},$$

which readily yields

$$(2.4) \quad \left| \frac{\Gamma(1-\lambda) z^{1+\lambda} D_z^{1+\lambda} f(z)}{f(z)} - 1 \right| = \left| \frac{\sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - 1 \right\} a_n z^n}{z - \sum_{n=2}^{\infty} a_n z^n} \right|$$

$$\leq \frac{\sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - 1 \right\} a_n}{1 - \sum_{n=2}^{\infty} a_n}$$

$$\leq 1 - \alpha,$$

provided that

$$(2.5) \quad \sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - 1 \right\} a_n \leq (1-\alpha) \left(1 - \sum_{n=2}^{\infty} a_n \right).$$

We note that (2.5) is equivalent to (2.1). Further, the result (2.1) is sharp for the functions

$$(2.6) \quad f(z) = z + \left[(1-\alpha) / \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \right] z^n \quad (n \geq 2).$$

Thus we complete the proof of Theorem 1.

REMARK 1. Letting $\lambda = 0$ in Theorem 1, we obtain the corresponding result for $\mathcal{S}^*(\alpha)$ due to Silverman [14].

THEOREM 2. Let the function $f(z)$ be defined by (1.1). If

$$(2.7) \quad \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} |a_n| \leq 1 - \alpha$$

for $\lambda < 1$ and $0 \leq \alpha < 1$, then $f(z) \in \mathcal{K}(\alpha, \lambda)$. The result (2.7) is sharp.

PROOF. Note that $f(z) \in \mathcal{K}(\alpha, \lambda)$ if and only if

$$\Gamma(1-\lambda) z^{1+\lambda} D_z^{1+\lambda} f(z) \in \mathcal{S}^*(\alpha, \lambda).$$

Therefore, on replacing a_n by $[\Gamma(n+1)\Gamma(1-\lambda)/\Gamma(n-\lambda)]a_n$ in Theorem 1, we have Theorem 2. Further, the result (2.7) is sharp for the functions

$$(2.8) \quad f(z) = z + \left[(1-\alpha) / \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \right] z^n \quad (n \geq 2).$$

REMARK 2. Letting $\lambda = 0$ in Theorem 2, we have the corresponding result for $\mathcal{K}(\alpha)$ given by Silverman [14].

THEOREM 3. Let the function $f(z)$ be defined by (1.9). Then $f(z)$ belongs to the class $\mathcal{J}^*(\alpha, \lambda)$ if and only if

$$(2.9) \quad \sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} a_n \leq 1 - \alpha$$

for $\lambda < 1$ and $0 \leq \alpha < 1$. The result (2.9) is sharp.

PROOF. By means of Theorem 1, (2.9) implies that $f(z) \in \mathcal{J}^*(\alpha, \lambda)$.

Suppose that $f(z)$ is in the class $\mathcal{J}^*(\alpha, \lambda)$. Then

$$(2.10) \quad \operatorname{Re} \left[\frac{\Gamma(1-\lambda) z^{1+\lambda} D_z^{1+\lambda} f(z)}{f(z)} \right] \\ = \operatorname{Re} \left[\frac{1 - \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} a_n z^{n-1}}{1 - \sum_{n=2}^{\infty} a_n z^{n-1}} \right] \\ > \alpha$$

for $\lambda < 1$, $0 \leq \alpha < 1$, and $z \in \mathcal{U}$. Choose values of z on the real axis

so that $\Gamma(1-\lambda)z^{1+\lambda} D_z^{1+\lambda} f(z)/f(z)$ is real. Upon clearing the denominator in (2.10) and letting $z \rightarrow 1^-$, we obtain

$$(2.11) \quad 1 - \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} a_n \cong \alpha \left(1 - \sum_{n=2}^{\infty} a_n \right)$$

which implies (2.9). The result (2.9) is sharp for the functions

$$(2.12) \quad f(z) = z - \left[(1-\alpha) / \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \right] z^n \quad (n \cong 2).$$

This completes the proof of Theorem 3.

COROLLARY 1. Let the function $f(z)$ defined by (1.9) belong to the class $\mathcal{F}^*(\alpha, \lambda)$. Then

$$(2.13) \quad a_n \cong (1-\alpha) / \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \quad (n \cong 2).$$

The result (2.13) is sharp for the functions $f(z)$ given by (2.12).

Next we have

THEOREM 4. Let the function $f(z)$ be defined by (1.9). Then $f(z)$ belongs to the class $\mathcal{C}(\alpha, \lambda)$ if and only if

$$(2.14) \quad \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} a_n \cong 1 - \alpha$$

for $\lambda < 1$ and $0 \cong \alpha < 1$. The result (2.14) is sharp for the functions

$$(2.15) \quad f(z) = z - \left[(1-\alpha) / \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \right] z^n \quad (n \cong 2).$$

COROLLARY 2. Let the function $f(z)$ defined by (1.9) belong to the class $\mathcal{C}(\alpha, \lambda)$. Then

$$(2.16) \quad a_n \leq (1-\alpha) \left/ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \right\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \} \quad (n \geq 2).$$

The result (2.16) is sharp for the functions $f(z)$ given by (2.15).

3. DISTORTION THEOREMS

By virtue of the coefficient inequalities, we can prove the following distortion theorems for functions $f(z)$ belonging to the classes $\mathcal{F}^*(\alpha, \lambda)$ and $\mathcal{C}(\alpha, \lambda)$.

LEMMA 1. The class $\mathcal{F}^*(\alpha, \lambda)$ is closed under linear combinations.

PROOF. Let the functions

$$(3.1) \quad f_i(z) = z - \sum_{n=2}^{\infty} a_{n,i} z^n \quad (i = 1, 2; a_{n,i} \geq 0)$$

be in the class $\mathcal{F}^*(\alpha, \lambda)$. Then we need only prove that the function

$$\delta f_1(z) + (1-\delta)f_2(z) \quad (0 \leq \delta \leq 1)$$

is also in the class $\mathcal{F}^*(\alpha, \lambda)$. In fact, we have

$$(3.2) \quad \begin{aligned} & \delta f_1(z) + (1-\delta)f_2(z) \\ &= z - \sum_{n=2}^{\infty} \left(\delta a_{n,1} + (1-\delta)a_{n,2} \right) z^n. \end{aligned}$$

Hence, with the aid of Theorem 3, we have

$$\begin{aligned}
 (3.3) \quad & \sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \left\{ \delta a_{n,1} + (1-\delta)a_{n,2} \right\} \\
 &= \delta \sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} a_{n,1} \\
 &\quad + (1-\delta) \sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} a_{n,2} \\
 &\leq \delta(1-\alpha) + (1-\delta)(1-\alpha) \\
 &= 1 - \alpha,
 \end{aligned}$$

which completes the proof of Lemma 1.

By means of Lemma 1, we know that $\mathcal{F}^*(\alpha, \lambda)$ is convex, and hence further that $\mathcal{F}^*(\alpha, \lambda)$ has some extreme points.

LEMMA 2. Let

$$(3.4) \quad f_1(z) = z$$

and

$$(3.5) \quad f_n(z) = z - \frac{1 - \alpha}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha} z^n \quad (n \geq 2).$$

Then $f(z)$ is in the class $\mathcal{F}^*(\alpha, \lambda)$ if and only if it can be expressed in the form

$$(3.6) \quad f(z) = \sum_{n=1}^{\infty} \lambda_n f_n(z),$$

where $\lambda_n \geq 0$ ($n \geq 1$) and

$$(3.7) \quad \sum_{n=1}^{\infty} \lambda_n = 1.$$

PROOF. We assume that $f(z)$ has the form (3.6). Then

$$(3.8) \quad f(z) = z - \sum_{n=2}^{\infty} \frac{(1-\alpha)\lambda_n}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha} z^n.$$

Consequently, we obtain

$$(3.9) \quad \begin{aligned} \sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \cdot \frac{(1-\alpha)\lambda_n}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha} \\ = (1-\alpha) \sum_{n=2}^{\infty} \lambda_n \\ = (1-\alpha)(1-\lambda_1) \\ \leq 1 - \alpha, \end{aligned}$$

which implies that $f(z) \in \mathcal{F}^*(\alpha, \lambda)$.

For the converse, we assume that $f(z)$ is in the class $\mathcal{F}^*(\alpha, \lambda)$. Then, by setting

$$(3.10) \quad \lambda_n = \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} a_n / (1-\alpha) \quad (n \geq 2)$$

and

$$(3.11) \quad \lambda_1 = 1 - \sum_{n=2}^{\infty} \lambda_n,$$

we have (3.6).

In view of Lemma 2, we have

THEOREM 5. The extreme points of $\mathcal{F}^*(\alpha, \lambda)$ are the functions $f_n(z)$
($n \geq 1$) given by (3.4) and (3.5).

Similarly, we have

THEOREM 6. The extreme points of $\mathcal{C}(\alpha, \lambda)$ are the functions $f_n(z)$
($n \geq 1$) given by (3.4) and

$$(3.12) \quad f_n(z) = z - \frac{1 - \alpha}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\}} z^n \quad (n \geq 2).$$

THEOREM 7. Let the function $f(z)$ defined by (1.9) belong to the class
 $\mathcal{F}^*(\alpha, \lambda)$. Then, if $-1 \leq \lambda < 1$

$$(3.13) \quad |f(z)| \geq |z| - \frac{(1-\alpha)(1-\lambda)}{2 - \alpha(1-\lambda)} |z|^2$$

and

$$(3.14) \quad |f(z)| \leq |z| + \frac{(1-\alpha)(1-\lambda)}{2 - \alpha(1-\lambda)} |z|^2$$

for $z \in \mathcal{U}$. Furthermore, if $0 \leq \lambda < 1$,

$$(3.15) \quad |f'(z)| \geq 1 - \frac{2(1-\alpha)(1-\lambda)}{2 - \alpha(1-\lambda)} |z|$$

and

$$(3.16) \quad |f'(z)| \leq 1 + \frac{2(1-\alpha)(1-\lambda)}{2 - \alpha(1-\lambda)} |z|$$

for $z \in \mathcal{U}$. The bounds (3.13) to (3.16) are sharp.

PROOF. Note that the extremal function is one of the extreme points.

Therefore, we have

$$(3.17) \quad |f(z)| \cong |z| - \max_{n \geq 2} \left\{ \frac{1 - \alpha}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha} |z|^n \right\}$$

and

$$(3.18) \quad |f(z)| \leq |z| + \max_{n \geq 2} \left\{ \frac{1 - \alpha}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha} |z|^n \right\}.$$

Now define

$$(3.19) \quad \mathcal{E}(\lambda, \alpha, n, |z|) = \frac{\Gamma(n-\lambda) |z|^n}{\Gamma(n+1)\Gamma(1-\lambda) - \alpha \Gamma(n-\lambda)}$$

for $-1 \leq \lambda < 1$, $0 \leq \alpha < 1$, $n \geq 2$, and $z \in \mathcal{U}$. If

$$(3.20) \quad \mathcal{E}(\lambda, \alpha, n, |z|) \geq \mathcal{E}(\lambda, \alpha, n+1, |z|)$$

for $|z| \neq 0$, then we have the first half of the theorem.

Set

$$(3.21) \quad \mathcal{E}_1(\lambda, \alpha, n, |z|) = \Gamma(n+1)\Gamma(1-\lambda) \{ (n+1) - (n-\lambda)|z| \} \\ - \alpha \Gamma(n+1-\lambda) (1-|z|).$$

If $\mathcal{E}_1(\lambda, \alpha, n, |z|) \geq 0$, then we have (3.20). We note that $\mathcal{E}_1(\lambda, \alpha, n, |z|)$ is

a decreasing function of α . This implies that

$$\begin{aligned}
 (3.22) \quad \mathcal{E}_1(\lambda, \alpha, n, |z|) &\geq \mathcal{E}_1(\lambda, 1, n, |z|) \\
 &= \{ \Gamma(n+2)\Gamma(1-\lambda) - \Gamma(n+1-\lambda) \} \\
 &\quad - \{ (n-\lambda)\Gamma(n+1)\Gamma(1-\lambda) - \Gamma(n+1-\lambda) \} |z|.
 \end{aligned}$$

Since $\mathcal{E}_1(\lambda, 1, n, |z|)$ is a decreasing function of $|z|$, we also have

$$\begin{aligned}
 (3.23) \quad \mathcal{E}_1(\lambda, \alpha, n, |z|) &\geq \mathcal{E}_1(\lambda, 1, n, 1) \\
 &= (1+\lambda)\Gamma(n+1)\Gamma(1-\lambda) \\
 &\geq 0
 \end{aligned}$$

for $-1 \leq \lambda < 1$, and $n \geq 2$.

In order to establish the second half of Theorem 7, we note that

$$(3.24) \quad |f'(z)| \geq 1 - \max_{n \geq 2} \frac{n(1-\alpha)}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha} |z|^{n-1}$$

and

$$(3.25) \quad |f'(z)| \leq 1 + \max_{n \geq 2} \frac{n(1-\alpha)}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha} |z|^{n-1}.$$

Define

$$(3.26) \quad F(\lambda, \alpha, n, |z|) = \frac{n \Gamma(n-\lambda) |z|^{n-1}}{\Gamma(n+1)\Gamma(1-\lambda) - \alpha \Gamma(n-\lambda)}$$

and

$$(3.27) \quad F_1(\lambda, \alpha, n, |z|) = \Gamma(n+2)\Gamma(1-\lambda)\{n - (n-\lambda)|z|\} \\ - \alpha \Gamma(n+1-\lambda)\{n - (n-\lambda)|z|\}$$

for $0 \leq \lambda < 1$, $0 \leq \alpha < 1$, $n \geq 2$, and $z \in \mathcal{U}$. Since $F_1(\lambda, \alpha, n, |z|)$ is a decreasing function of α ,

$$(3.28) \quad F_1(\lambda, \alpha, n, |z|) \geq F_1(\lambda, 1, n, |z|) \\ = n\{\Gamma(n+2)\Gamma(1-\lambda) - \Gamma(n+1-\lambda)\} \\ - (n-\lambda)\{\Gamma(n+2)\Gamma(1-\lambda) - \Gamma(n+1-\lambda)\}|z|.$$

Further, since $F_1(\lambda, 1, n, |z|)$ is a decreasing function of $|z|$, we get

$$(3.29) \quad F_1(\lambda, \alpha, n, |z|) \geq F_1(\lambda, 1, n, 1) \\ = \lambda\{\Gamma(n+2)\Gamma(1-\lambda) - \Gamma(n+1-\lambda)\} \\ \geq 0$$

for $0 \leq \lambda < 1$ and $n \geq 2$. Thus we have (3.15) and (3.16).

Finally, by taking the function

$$(3.30) \quad f(z) = z - \frac{(1-\alpha)(1-\lambda)}{2 - \alpha(1-\lambda)} z^2,$$

we can show that all bounds given by Theorem 7 are sharp.

COROLLARY 3. Let the function $f(z)$ defined by (1.9) be in the class $\mathcal{F}(\alpha, \lambda)$ with $-1 \leq \lambda < 1$ and $0 \leq \alpha < 1$. Then the unit disk \mathcal{U} is mapped

onto a domain that contains the disk $|w| < (1+\lambda)/\{2 - \alpha(1-\lambda)\}$.

THEOREM 8. Let the function $f(z)$ defined by (1.9) be in the class $\mathcal{C}(\alpha, \lambda)$. Then, if $-1 \leq \lambda < 1$,

$$(3.31) \quad |f(z)| \geq |z| - \frac{(1-\alpha)(1-\lambda)^2}{2\{2 - \alpha(1-\lambda)\}} |z|^2$$

and

$$(3.32) \quad |f(z)| \leq |z| + \frac{(1-\alpha)(1-\lambda)^2}{2\{2 - \alpha(1-\lambda)\}} |z|^2$$

for $z \in \mathcal{U}$. Furthermore, if $0 \leq \lambda < 1$,

$$(3.33) \quad |f'(z)| \geq 1 - \frac{(1-\alpha)(1-\lambda)^2}{2 - \alpha(1-\lambda)} |z|$$

and

$$(3.34) \quad |f'(z)| \leq 1 + \frac{(1-\alpha)(1-\lambda)^2}{2 - \alpha(1-\lambda)} |z|$$

for $z \in \mathcal{U}$. The bounds (3.31) to (3.34) are sharp.

PROOF. By means of Theorem 6, we have

$$(3.35) \quad |f(z)| \geq |z| - \max_{n \geq 2} \left\{ \frac{1 - \alpha}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\}} \right\} |z|^n$$

and

$$(3.36) \quad |f(z)| \leq |z| + \max_{n \geq 2} \left\{ \frac{1 - \alpha}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\}} \right\} |z|^n.$$

Let

$$(3.37) \quad H(\lambda, \alpha, n, |z|) = \frac{\Gamma(n-\lambda)^2 |z|^n}{\Gamma(n+1)\Gamma(1-\lambda) \{ \Gamma(n+1)\Gamma(1-\lambda) - \Gamma(n-\lambda)\alpha \}}$$

and

$$(3.38) \quad \begin{aligned} H_1(\lambda, \alpha, n, |z|) &= \Gamma(n+1)\Gamma(1-\lambda) \{ (n+1)^2 - (n-\lambda)^2 |z| \} \\ &\quad - \alpha(n-\lambda)\Gamma(n-\lambda) \{ (n+1) - (n-\lambda) |z| \} \end{aligned}$$

for $-1 \leq \lambda < 1$, $0 \leq \alpha < 1$, $n \geq 2$, and $z \in \mathcal{U}$. Then, we know that

$H(\lambda, \alpha, n, |z|)$ is a decreasing function of n if

$$H_1(\lambda, \alpha, n, |z|) \geq 0.$$

In fact, since $H_1(\lambda, \alpha, n, |z|)$ is a decreasing function of α , and $H_1(\lambda, 1, n, |z|)$ is a decreasing function of $|z|$, we can prove that

$$(3.39) \quad \begin{aligned} H_1(\lambda, \alpha, n, |z|) &\geq H_1(\lambda, 1, n, |z|) \\ &\geq H_1(\lambda, 1, n, |z|) \\ &= (1+\lambda) \{ (2n+1-\lambda)\Gamma(n+1)\Gamma(1-\lambda) - \Gamma(n+1-\lambda) \} \\ &\geq 0 \end{aligned}$$

for $-1 \leq \lambda < 1$ and $n \geq 2$. Consequently, we have the first half of Theorem 8.

Next, we note that

$$(3.40) \quad |f'(z)| \geq 1 - \max_{n \geq 2} \frac{1 - \alpha}{\frac{\Gamma(n)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\}} |z|^{n-1}$$

and

$$(3.41) \quad |f'(z)| \leq 1 + \max_{n \geq 2} \frac{1 - \alpha}{\frac{\Gamma(n)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\}} |z|^{n-1}.$$

Define

$$(3.42) \quad G(\lambda, \alpha, n, |z|) = \frac{\Gamma(n-\lambda)^2 |z|^{n-1}}{\Gamma(n)\Gamma(1-\lambda) \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\}}$$

and

$$(3.43) \quad G_1(\lambda, \alpha, n, |z|) = \Gamma(n+1)\Gamma(1-\lambda) \{n(n+1) - (n-\lambda)^2 |z|\} \\ - \alpha \Gamma(n+1-\lambda) \{n - (n-\lambda) |z|\}$$

for $0 \leq \lambda < 1$, $0 \leq \alpha < 1$, $n \geq 2$, and $z \in \mathcal{U}$. Then it is sufficient to prove that

$$G_1(\lambda, \alpha, n, |z|) \geq 0.$$

Note that $G_1(\lambda, \alpha, n, |z|)$ is a decreasing function of α , and $G_1(\lambda, 1, n, |z|)$ is a decreasing function of $|z|$. Thus we have,

$$(3.44) \quad G_1(\lambda, \alpha, n, |z|) \geq G_1(\lambda, 1, n, |z|) \\ \geq G_1(\lambda, 1, n, 1) \\ = (n+2\lambda n-\lambda^2)\Gamma(n+1)\Gamma(1-\lambda) - \lambda \Gamma(n+1-\lambda) \\ \geq 0$$

for $0 \leq \lambda < 1$ and $n \geq 2$, which implies the second half of Theorem 8.

Finally, all bounds asserted by Theorem 8 are sharp for the function

$$(3.45) \quad f(z) = z - \frac{(1-\alpha)(1-\lambda)^2}{2\{2 - \alpha(1-\lambda)\}} z^2.$$

COROLLARY 4. Let the function $f(z)$ defined by (1.9) be in the class $\mathcal{C}(\alpha, \lambda)$ with $-1 \leq \lambda < 1$ and $0 \leq \alpha < 1$. Then the unit disk \mathcal{U} is mapped onto a domain that contains the disk $|w| < r_0$, where r_0 is given by

$$(3.46) \quad r_0 = 1 - \frac{(1-\alpha)(1-\lambda)^2}{2\{2 - \alpha(1-\lambda)\}}.$$

4. STARLIKENESS AND CONVEXITY

Silverman [14] proved the following lemmas.

LEMMA 3. Let the function $f(z)$ be defined by (1.9). Then $f(z)$ is starlike of order α if and only if

$$(4.1) \quad \sum_{n=2}^{\infty} (n-\alpha) a_n \leq 1 - \alpha$$

for $0 \leq \alpha < 1$.

LEMMA 4. Let the function $f(z)$ be defined by (1.9). Then $f(z)$ is convex of order α if and only if

$$(4.2) \quad \sum_{n=2}^{\infty} n(n-\alpha) a_n \leq 1 - \alpha$$

for $0 \leq \alpha < 1$.

By applying the above lemmas, we now prove

THEOREM 9. Let the function $f(z)$ defined by (1.9) be in the class $\mathcal{F}^*(\alpha, \lambda)$ with $0 \leq \lambda < 1$ and $0 \leq \alpha < 1$. Then $f(z)$ is starlike of order α .

PROOF. Note that

$$(4.3) \quad n \leq \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)}$$

for $0 \leq \lambda < 1$, $0 \leq \alpha < 1$, and $n \geq 2$. This shows that

$$(4.4) \quad \sum_{n=2}^{\infty} (n-\alpha) a_n \leq \sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} a_n$$

$$\leq 1 - \alpha,$$

and we complete the proof of Theorem 9 in view of (4.1).

Similarly, by using Lemma 4, we have

THEOREM 10. Let the function $f(z)$ defined by (1.9) be in the class $\mathcal{C}(\alpha, \lambda)$ with $0 \leq \lambda < 1$ and $0 \leq \alpha < 1$. Then $f(z)$ is convex of order α .

REMARK 3. For $\lambda = 0$, the classes $\mathcal{F}^*(\alpha, \lambda)$ and $\mathcal{C}(\alpha, \lambda)$ reduce to the classes $\mathcal{F}^*(\alpha)$ and $\mathcal{C}(\alpha)$, respectively, which were introduced by Silverman [14]. It follows that

$$(4.5) \quad \mathcal{F}^*(\alpha, 0) = \mathcal{F}^*(\alpha)$$

and

$$(4.6) \quad \mathcal{C}(\alpha, 0) = \mathcal{C}(\alpha).$$

Hence, by means of Theorem 9 and Theorem 10, we have

$$(4.7) \quad \mathcal{F}^*(\alpha, \lambda) \subset \mathcal{F}^*(\alpha, 0) \quad (0 \leq \lambda < 1)$$

and

$$(4.8) \quad \mathcal{C}(\alpha, \lambda) \subset \mathcal{C}(\alpha, 0) \quad (0 \leq \lambda < 1).$$

Since

$$(4.9) \quad \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \leq n - \alpha$$

and

$$(4.10) \quad \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \leq n(n-\alpha)$$

for $\lambda < 0$, $0 \leq \alpha < 1$, and $n \geq 2$, we also have

$$(4.11) \quad \mathcal{F}^*(\alpha, \lambda) \supset \mathcal{F}^*(\alpha, 0) \quad (\lambda < 0)$$

and

$$(4.12) \quad \mathcal{C}(\alpha, \lambda) \supset \mathcal{C}(\alpha, 0) \quad (\lambda < 0).$$

THEOREM 11. Let the function $f(z)$ defined by (1.9) be in the class $\mathcal{F}^*(\alpha, \lambda)$ with $\lambda < 1$ and $0 \leq \alpha < 1$. Then $f(z)$ is convex of order δ

($0 \leq \delta < 1$) in the disk $|z| < r_1$, where

$$(4.13) \quad r_1 = \inf_{n \geq 2} \left[\frac{(1-\delta) \{ \Gamma(n+1) \Gamma(1-\lambda) - \alpha \Gamma(n-\lambda) \}}{n(n-\delta)(1-\alpha) \Gamma(n-\lambda)} \right]^{1/(n-1)}.$$

PROOF. It is sufficient to show that

$$(4.14) \quad \left| \frac{zf''(z)}{f'(z)} \right| < 1 - \delta$$

for $|z| < r_1$. Note that

$$(4.15) \quad \left| \frac{zf''(z)}{f'(z)} \right| \leq \frac{\sum_{n=2}^{\infty} n(n-1)a_n |z|^{n-1}}{1 - \sum_{n=2}^{\infty} n a_n |z|^{n-1}} \leq 1 - \delta,$$

if and only if

$$(4.16) \quad \sum_{n=2}^{\infty} n(n-\delta)a_n |z|^{n-1} \leq 1 - \delta.$$

With the aid of Theorem 3, we need only find the values of $|z|$ for which

$$(4.17) \quad |z|^{n-1} \leq \frac{(1-\delta) \{ \Gamma(n+1) \Gamma(1-\lambda) - \alpha \Gamma(n-\lambda) \}}{n(n-\delta)(1-\alpha) \Gamma(n-\lambda)}, \quad (n \geq 2)$$

and the proof of Theorem 11 is thus completed.

5. CLASSES OF FUNCTIONS WITH FIXED FINITELY MANY COEFFICIENTS

In view of Theorem 3, we denote by $\mathcal{F}^*(\alpha, \lambda, k)$ the subclass of $\mathcal{F}^*(\alpha, \lambda)$ consisting of all functions of the form

$$(5.1) \quad f(z) = z - \sum_{i=2}^k A_i p_i z^i - \sum_{n=k+1}^{\infty} a_n z^n,$$

where

$$(5.2) \quad A_i = (1-\alpha) / \left\{ \frac{\Gamma(i+1)\Gamma(1-\lambda)}{\Gamma(i-\lambda)} - \alpha \right\},$$

$$0 \leq p_i \leq 1, \quad \text{and} \quad 0 \leq \sum_{i=2}^k p_i \leq 1.$$

Further, in view of Theorem 4, we denote by $\mathcal{C}(\alpha, \lambda, k)$ the subclass of $\mathcal{C}(\alpha, \lambda)$ consisting of all functions of the form

$$(5.3) \quad f(z) = z - \sum_{i=2}^k B_i p_i z^i - \sum_{n=k+1}^{\infty} a_n z^n,$$

where

$$(5.4) \quad B_i = (1-\alpha) / \frac{\Gamma(i+1)\Gamma(1-\lambda)}{\Gamma(i-\lambda)} \left\{ \frac{\Gamma(i+1)\Gamma(1-\lambda)}{\Gamma(i-\lambda)} - \alpha \right\},$$

$$0 \leq p_i \leq 1, \quad \text{and} \quad 0 \leq \sum_{i=2}^k p_i \leq 1.$$

Our first result on the classes of functions with fixed finitely many coefficients is contained in

THEOREM 12. Let the function $f(z)$ be defined by (5.1). Then $f(z)$ is in the class $\mathcal{F}^*(\alpha, \lambda, k)$ if and only if

$$(5.5) \quad \sum_{n=k+1}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} a_n \leq (1-\alpha) \left[1 - \sum_{i=2}^k p_i \right]$$

for $\lambda < 1$ and $0 \leq \alpha < 1$. The result (5.5) is sharp.

PROOF. Putting

$$(5.6) \quad a_i = A_i p_i \quad (i = 2, 3, \dots, k)$$

in Theorem 3, we have

$$(5.7) \quad (1-\alpha) \sum_{i=2}^k p_i + \sum_{n=k+1}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} a_n \leq 1 - \alpha$$

which implies (5.5). Furthermore, the result is sharp for the functions

$$(5.8) \quad f(z) = z - \sum_{i=2}^k A_i p_i z^i \\ - \left[(1-\alpha) \left(1 - \sum_{i=2}^k p_i \right) / \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \right] z^n \quad (n \geq k+1).$$

COROLLARY 5. Let the function $f(z)$ defined by (5.1) be in the class $\mathcal{F}^*(\alpha, \lambda, k)$. Then

$$(5.9) \quad a_n \leq (1-\alpha) \left[1 - \sum_{i=2}^k p_i \right] / \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \quad (n \geq k+1).$$

The result (5.9) is sharp for functions $f(z)$ of the form (5.8).

Similarly, we prove

THEOREM 13. Let the function $f(z)$ be defined by (5.3). Then $f(z)$ is
in the class $\mathcal{C}(\alpha, \lambda, k)$ if and only if

$$(5.10) \quad \sum_{n=k+1}^{\infty} \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} a_n \leq (1-\alpha) \left[1 - \sum_{i=2}^k p_i \right]$$

for $\lambda < 1$ and $0 \leq \alpha < 1$. The result (5.10) is sharp for the functions

$$(5.11) \quad f(z) = z - \sum_{i=2}^k B_i p_i z^i$$

$$- \left[(1-\alpha) \left(1 - \sum_{i=2}^k p_i \right) \right] / \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} z^n$$

($n \geq k + 1$).

COROLLARY 6. Let the function $f(z)$ defined by (5.3) be in the class
 $\mathcal{C}(\alpha, \lambda, k)$. Then

$$(5.12) \quad a_n \leq (1-\alpha) \left(1 - \sum_{i=2}^k p_i \right) / \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \quad (n \geq k + 1).$$

The result (5.12) is sharp for functions $f(z)$ of the form (5.11).

We find the extreme points of $\mathcal{T}^*(\alpha, \lambda, k)$ and $\mathcal{C}(\alpha, \lambda, k)$ by using the same technique as in Theorem 5.

THEOREM 14. The extreme points of $\mathcal{T}^*(\alpha, \lambda, k)$ are

$$(5.13) \quad f_k(z) = z - \sum_{i=2}^k A_i p_i z^i$$

and

$$(5.14) \quad f_n(z) = z - \sum_{i=2}^k A_i p_i z^i$$

$$- \left[(1-\alpha) \left(1 - \sum_{i=2}^k p_i \right) / \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \right] z^n \quad (n \geq k + 1),$$

where A_i is given by (5.2).

THEOREM 15. The extreme points of $\mathcal{C}(\alpha, \lambda, k)$ are

$$(5.15) \quad f_k(z) = z - \sum_{i=2}^k B_i p_i z^i$$

and

$$(5.16) \quad f_n(z) = z - \sum_{i=2}^k B_i p_i z^i$$

$$- \left[(1-\alpha) \left(1 - \sum_{i=2}^k p_i \right) / \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} \right] z^n \quad (n \geq k + 1),$$

where B_i is given by (5.4).

Next we determine the radius of convexity for functions belonging to the class $\mathcal{F}^*(\alpha, \lambda, k)$. Our result is given by

THEOREM 16. Let the function $f(z)$ defined by (5.1) be in the class $\mathcal{F}^*(\alpha, \lambda, k)$. Then $f(z)$ is convex in the disk $|z| < r_2$, where r_2 is the largest value for which

$$(5.17) \quad \sum_{i=2}^k i^2 A_i p_i r^{i-1} + \frac{n^2(1-\alpha) \left(1 - \sum_{i=2}^k p_i\right) \Gamma(n-\lambda)}{\Gamma(n+1)\Gamma(1-\lambda) - \alpha \Gamma(n-\lambda)} r^{n-1} \leq 1$$

for $n \geq k + 1$. The result (5.17) is sharp.

PROOF. It suffices to prove that

$$(5.18) \quad \left| \frac{zf''(z)}{f'(z)} \right| \leq \frac{\sum_{i=2}^k i(i-1)A_i p_i r^{i-1} + \sum_{n=k+1}^{\infty} n(n-1)a_n r^{n-1}}{1 - \sum_{i=2}^k i A_i p_i r^{i-1} - \sum_{n=k+1}^{\infty} n a_n r^{n-1}} \leq 1$$

for $|z| \leq r$. Note that (5.18) is true if and only if

$$(5.19) \quad \sum_{i=2}^k i^2 A_i p_i r^{i-1} + \sum_{n=k+1}^{\infty} n^2 a_n r^{n-1} \leq 1.$$

By virtue of Theorem 12, we can write

$$(5.20) \quad a_n = \frac{(1-\alpha) \left(1 - \sum_{i=2}^k p_i\right)}{\frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha} \lambda_n \quad (n \geq k + 1),$$

where $\lambda_n \geq 0$ ($n \geq k + 1$) and

$$(5.21) \quad \sum_{n=k+1}^{\infty} \lambda_n \leq 1.$$

For each fixed r , we choose the integer $n_0 = n_0(r)$ for which

$$n^2 r^{n-1} / \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\}$$

is a maximum. Then we have

$$(5.22) \quad \sum_{n=k+1}^{\infty} n^2 a_n r^{n-1} \leq \frac{n_0^2(1-\alpha) \left(1 - \sum_{i=2}^k p_i \right)}{\frac{\Gamma(n_0+1)\Gamma(1-\lambda)}{\Gamma(n_0-\lambda)} - \alpha} r^{n_0-1}.$$

Hence the function $f(z)$ is convex in $|z| \leq r_2$ if

$$(5.23) \quad \sum_{i=2}^k i^2 A_i p_i r^{i-1} + \frac{n_0^2(1-\alpha) \left(1 - \sum_{i=2}^k p_i \right)}{\frac{\Gamma(n_0+1)\Gamma(1-\lambda)}{\Gamma(n_0-\lambda)} - \alpha} r^{n_0-1} \leq 1.$$

We find the value r_0 and the corresponding $n_0(r_0)$ so that

$$(5.24) \quad \sum_{i=2}^k i^2 A_i p_i r_0^{i-1} + \frac{n_0^2(1-\alpha) \left(1 - \sum_{i=2}^k p_i \right)}{\frac{\Gamma(n_0+1)\Gamma(1-\lambda)}{\Gamma(n_0-\lambda)} - \alpha} r_0^{n_0-1} = 1.$$

Then this value r_0 is the radius of convexity of functions $f(z)$ in the class $\mathcal{F}^*(\alpha, \lambda, k)$. Furthermore, we know that the result of the theorem is sharp for the functions $f(z)$ given by (5.8).

6. QUASICONFORMAL EXTENSION

Let f be a sense-preserving homeomorphism of a domain \mathcal{D} . Then f is said to be a $\left(\frac{1+k}{1-k}\right)$ -quasiconformal mapping of \mathcal{D} if it is absolutely continuous on almost all lines parallel to the coordinate axes and

$$(6.1) \quad \left| \frac{\partial}{\partial \bar{z}} f(z) \right| \leq k \left| \frac{\partial}{\partial z} f(z) \right| \quad (z \in \mathcal{D})$$

for $0 \leq k < 1$. We say that a function $f(z)$ is in the class \mathcal{S}_k if it is in the class \mathcal{S} and has a $\left(\frac{1+k}{1-k}\right)$ -quasiconformal extension defined on the extended complex plane (cf. [5]).

Let $\tilde{\mathcal{S}}^*(\alpha, \lambda)$ denote the subclass of \mathcal{S} whose members satisfy the coefficient inequality (2.1) for $\lambda < 1$ and $0 \leq \alpha < 1$. Then it follows that

$$(6.2) \quad \mathcal{F}^*(\alpha, \lambda) \subset \tilde{\mathcal{S}}^*(\alpha, \lambda) \subset \mathcal{S}^*(\alpha, \lambda).$$

Also let $\tilde{\mathcal{H}}(\alpha, \lambda)$ denote the subclass of \mathcal{S} whose members satisfy the coefficient inequality (2.7) for $\lambda < 1$ and $0 \leq \alpha < 1$. Then we have

$$(6.3) \quad \mathcal{C}(\alpha, \lambda) \subset \tilde{\mathcal{H}}(\alpha, \lambda) \subset \mathcal{H}(\alpha, \lambda).$$

We shall apply the following lemma due to Fait, Krzyz and Zygmunt [3] in order to prove Theorem 17 below.

LEMMA 5. If the function $f(z)$ defined by (1.1) is in the class \mathcal{A} and satisfies

$$(6.4) \quad \sum_{n=2}^{\infty} n |a_n| \leq k,$$

then $f(z) \in \mathcal{S}_k$.

THEOREM 17. Let the function $f(z)$ defined by (1.1) be in the class $\tilde{\mathcal{S}}^*(\alpha, \lambda)$ with $0 < \lambda < 1$ and $0 \leq \alpha < 1$. Then $f(z) \in \mathcal{S}_k$, where

$$(6.5) \quad k = \frac{2(1-\alpha)(1-\lambda)}{2 - \alpha(1-\lambda)}.$$

PROOF. In view of (2.1) and Lemma 5, we need to prove that (2.1) implies (6.4). Define $L(\alpha, \lambda, n)$ by

$$(6.6) \quad L(\alpha, \lambda, n) = \frac{1}{1 - \alpha} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} - \frac{n}{k}$$

for $0 < \lambda < 1$, $0 \leq \alpha < 1$, and $n \geq 2$. Then it suffices to show that $L(\alpha, \lambda, n) \geq 0$ for $0 < \lambda < 1$, $0 \leq \alpha < 1$, and $n \geq 2$. Note that $L(\alpha, \lambda, n) \geq 0$ if

$$(6.7) \quad k \geq \frac{n(1-\alpha)\Gamma(n-\lambda)}{\Gamma(n+1)\Gamma(1-\lambda) - \alpha\Gamma(n-\lambda)}$$

for $0 < \lambda < 1$, $0 \leq \alpha < 1$, and $n \geq 2$. Since

$$\frac{n(1-\alpha)\Gamma(n-\lambda)}{\Gamma(n+1)\Gamma(1-\lambda) - \alpha\Gamma(n-\lambda)}$$

is a decreasing function of n , and

$$(6.8) \quad 0 \leq \frac{2(1-\alpha)(1-\lambda)}{2 - \alpha(1-\lambda)} < 1$$

for $0 < \lambda < 1$ and $0 \leq \alpha < 1$, we conclude that $f(z) \in \mathcal{S}_k$ for k given by (6.5).

COROLLARY 7. Let the function $f(z)$ defined by (1.9) be in the class $\mathcal{F}^*(\alpha, \lambda)$ with $0 < \lambda < 1$ and $0 \leq \alpha < 1$. Then $f(z) \in \mathcal{S}_k$ for k given by (6.5).

Recently, Brown [1] established the following results.

LEMMA 6. If the function $f(z)$ defined by (1.1) is in the class \mathcal{A} and
satisfies

$$(6.9) \quad \sum_{n=2}^{\infty} \left(n - \frac{1-k}{1+k} \right) |a_n| \leq \frac{2k}{1+k},$$

then $f(z) \in \mathcal{S}^* \cap \mathcal{S}_k$.

LEMMA 7. If the function $f(z)$ defined by (1.1) is in the class \mathcal{A} and
satisfies

$$(6.10) \quad \sum_{n=2}^{\infty} n \left(n - \frac{1-k}{1+k} \right) |a_n| \leq \frac{2k}{1+k},$$

then $f(z) \in \mathcal{H} \cap \mathcal{S}_k$.

THEOREM 18. Let the function $f(z)$ defined by (1.1) be in the class
 $\tilde{\mathcal{S}}^*(\alpha, \lambda)$ with $0 \leq \lambda < 1$ and $0 < \alpha < 1$. Then $f(z) \in \mathcal{S}^*(\alpha) \cap \mathcal{S}_k$, where
 $k = (1-\alpha)/(1+\alpha)$.

PROOF. Since

$$(6.11) \quad n \leq \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)}$$

for $0 \leq \lambda < 1$ and $n \geq 2$, we have

$$(6.12) \quad \sum_{n=2}^{\infty} (n-\alpha) |a_n| \leq \sum_{n=2}^{\infty} \left\{ \frac{\Gamma(n+1)\Gamma(1-\lambda)}{\Gamma(n-\lambda)} - \alpha \right\} |a_n|$$

$$\leq 1 - \alpha,$$

which implies that $f(z) \in \mathcal{S}^*(\alpha)$, by means of [14].

Further, replacing α by $(1-k)/(1+k)$ in (6.12), we get (6.9). Hence $f(z)$ is also in the class $\mathcal{S}^* \cap \mathcal{S}_k$ for $k = (1-\alpha)/(1+\alpha)$. We note that $\mathcal{S}^*(\alpha) \subset \mathcal{S}^*$ for $0 \leq \alpha < 1$. Therefore, $f(z) \in \mathcal{S}^*(\alpha) \cap \mathcal{S}_k$.

COROLLARY 8. Let the function $f(z)$ defined by (1.9) be in the class $\mathcal{F}^*(\alpha, \lambda)$ with $0 \leq \lambda < 1$ and $0 < \alpha < 1$. Then $f(z) \in \mathcal{F}^*(\alpha, 0) \cap \mathcal{S}_k$, where $k = (1-\alpha)/(1+\alpha)$.

By using Lemma 7, we prove

THEOREM 19. Let the function $f(z)$ defined by (1.1) be in the class $\mathcal{H}(\alpha, \lambda)$ with $0 \leq \lambda < 1$ and $0 < \alpha < 1$. Then $f(z) \in \mathcal{H}(\alpha) \cap \mathcal{S}_k$, where $k = (1-\alpha)/(1+\alpha)$.

COROLLARY 9. Let the function $f(z)$ defined by (1.9) be in the class $\mathcal{C}(\alpha, \lambda)$ with $0 \leq \lambda < 1$ and $0 < \alpha < 1$. Then $f(z) \in \mathcal{C}(\alpha, 0) \cap \mathcal{S}_k$, where $k = (1-\alpha)/(1+\alpha)$.

ACKNOWLEDGEMENTS

The author's thanks are due to Professor H.M. Srivastava for his kind encouragement and guidance. The author also wishes to thank the referee for his valuable suggestions.

REFERENCES

- [1] J.E. Brown, Quasiconformal extensions for some geometric subclasses of univalent functions, Internat. J. Math. and Math. Sci. 7(1984), 187-195.
- [2] A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, Tables of Integral Transforms, Vol. II, McGraw-Hill Book Co., New York, London and Toronto, 1954.
- [3] M. Fait, J. Krzyz and J. Zygmunt, Explicit quasiconformal extensions for some classes of univalent functions, Comment. Math. Helv. 51(1976), 279-285.
- [4] T.H. MacGregor, The radius of convexity for starlike functions of order $\frac{1}{2}$, Proc. Amer. Math. Soc. 14(1963), 71-76.
- [5] O. Lehto and K.I. Virtanen, Quasiconformal Mappings in the Plane, Springer-Verlag, Berlin, Heidelberg and New York, 1973.
- [6] K. Nishimoto, Fractional derivative and integral. Part I, J. College Engrg. Nihon Univ. Ser. B 17(1976), 11-19.
- [7] T.J. Osler, Leibniz rule for fractional derivatives generalized and an application to infinite series, SIAM J. Appl. Math. 18(1970), 658-674.
- [8] S. Owa, On the distortion theorems. I, Kyungpook Math. J. 18(1978), 53-59.
- [9] B. Pinchuk, On starlike and convex functions of order α , Duke Math. J. 35(1968), 721-734.
- [10] M.S. Robertson, On the theory of univalent functions, Ann. of Math. 37(1936), 374-408.
- [11] B. Ross, A brief history and exposition of the fundamental theory of fractional calculus, in Fractional Calculus and Its Applications (B. Ross, Ed.), pp. 1-36, Springer-Verlag, Berlin, Heidelberg and New York, 1975.
- [12] M. Saigo, A remark on integral operators involving the Gauss hypergeometric functions, Math. Rep. College General Ed. Kyushu Univ. 11(1978), 135-143.
- [13] A. Schild, On starlike functions of order α , Amer. J. Math. 87(1965), 65-70.
- [14] H. Silverman, Univalent functions with negative coefficients, Proc. Amer. Math. Soc. 51(1975), 109-116.
- [15] H.M. Srivastava and R.G. Buschman, Convolution Integral Equations with Special Function Kernels, John Wiley and Sons, New York, London, Sydney and Toronto, 1977.

- [16] H.M. Srivastava and S. Owa, An application of the fractional derivative, Math. Japon. 29(1984), 383-389.
- [17] H.M. Srivastava, S. Owa and K. Nishimoto, Some fractional differintegral equations, J. Math. Anal. Appl. 106(1985), to appear.
- [18] H.M. Srivastava, S. Owa and K. Nishimoto, A note on a certain class of fractional differintegral equations, J. College Engrg. Nihon Univ. Ser. B 25(1984), 69-73.

Department of Mathematics
Kinki University
Higashi-Osaka, Osaka 577
Japan