



mathematics



Article

Laguerre-Type Bernoulli and Euler Numbers and Related Fractional Polynomials

Paolo Emilio Ricci, Rekha Srivastava and Diego Caratelli

Special Issue

Advances in Higher-Order Linear Equations and Linear Polynomial Differential Operators

Edited by

Prof. Dr. Rekha Srivastava



<https://doi.org/10.3390/math12030381>

Article

Laguerre-Type Bernoulli and Euler Numbers and Related Fractional Polynomials

Paolo Emilio Ricci ^{1,*}, Rekha Srivastava ^{2,†} and Diego Caratelli ^{3,†}

¹ Mathematics Section, International Telematic University UniNettuno, Corso Vittorio Emanuele II 39, 00186 Rome, Italy

² Department of Mathematics and Statistics, University of Victoria, Victoria, BC V8W 3R4, Canada; rekhas@math.uvic.ca

³ Department of Electrical Engineering, Eindhoven University of Technology, 5600 MB Eindhoven, The Netherlands; d.caratelli@tue.nl

* Correspondence: paoloemilio.ricci@uninettunouniversity.net or paoloemilioricci@gmail.com

† These authors contributed equally to this work.

Abstract: We extended the classical Bernoulli and Euler numbers and polynomials to introduce the Laguerre-type Bernoulli and Euler numbers and related fractional polynomials. The case of fractional Bernoulli and Euler polynomials and numbers has already been considered in a previous paper of which this article is a further generalization. Furthermore, we exploited the Laguerre-type fractional exponentials to define a generalized form of the classical Laplace transform. We show some examples of these generalized mathematical entities, which were derived using the computer algebra system Mathematica© (latest v. 14.0).

Keywords: Bernoulli numbers and polynomials; Euler numbers and polynomials; Laguerre-type exponential functions; generating functions; generalized Laplace transform

MSC: 11B83; 33C99; 33C10; 34A30; 44A10



Citation: Ricci, P.E.; Srivastava, R.; Caratelli, D. Laguerre-Type Bernoulli and Euler Numbers and Related Fractional Polynomials. *Mathematics* **2024**, *12*, 381. <https://doi.org/10.3390/math12030381>

Academic Editor: Li Guo

Received: 18 December 2023

Revised: 18 January 2024

Accepted: 22 January 2024

Published: 24 January 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The purpose of this paper is to introduce a different version of the Bernoulli and Euler numbers and polynomials, related to the Laguerre-type exponentials.

The consideration of generalized forms of polynomials and classical numbers has received much attention in recent years, as can be seen in the [1–5]. This has often been achieved by exploiting the generating functions [6–8].

One recent field has involved the extension of such entities to the case of fractional numbers or polynomials in relation to the fractional derivative and its applications (see [9] and the references therein).

In a recent article [10], using the fractional exponential function and the generating function of the classical Bernoulli and Euler numbers and polynomials, we have introduced the fractional index case of these well-known entities. Their definition is shortly reported in Section 3.

It is worth noting that a similar construction can be achieved by exploiting different exponential functions. Examples of such exponentials are shown in Section 2, where the definition of *true* exponentials with respect to suitable differential operators is highlighted.

The Laguerre-type exponentials

$$e_1(x) := \sum_{k=0}^{\infty} \frac{x^k}{(k!)^2}, \quad e_n(x) := \sum_{k=0}^{\infty} \frac{x^k}{(k!)^{n+1}},$$

where n is a positive integer, are recalled in Section 4. These functions have been considered in several articles (see, e.g., [11,12]), where it has been shown that these functions are related

to a differential isomorphism acting onto the space \mathcal{A}_x of the analytic functions, of the real (or complex) variable x , preserving differential identities at a higher level of differentiation. This allows us to introduce a symmetry, into the space \mathcal{A}_x , in which to a classical special function, it is possible to associate the Laguerre-type one, and since this can be repeated for each level of Laguerre-type exponentials, the construction can be iterated as many times as we need.

Since in the above-cited article [10] we have considered the fractional exponential function, we first extend to the fractional case the Laguerre-type exponentials by considering the expansions

$$e_{\alpha,1}(x) := \sum_{k=0}^{\infty} \frac{x^{k\alpha}}{[\Gamma(k\alpha + 1)]^2}, \quad e_{\alpha,n}(x) := \sum_{k=0}^{\infty} \frac{x^{k\alpha}}{[\Gamma(k\alpha + 1)]^{n+1}}.$$

Then, the construction of the Laguerre-type Bernoulli and Euler numbers and polynomials, starting from their generating functions in which the classical exponential is replaced by its Laguerre-type version, can be obtained in both cases, namely exploiting the Laguerre-type exponentials or their fractional Laguerre-type versions.

This is examined in Section 5, where we limit ourselves to consider only the first-order Laguerre-type derivative since the used technique is always the same, and only different numbers or functions are obtained. Furthermore, as the numbers and functions of the fractional case tend to the corresponding ordinary ones in the limit $\alpha \rightarrow 1$, we will consider tables of the fractional case for special values of the parameter α , including $\alpha = 1$.

Lastly, in Sections 5 and 6, exploiting the Laguerre-type exponentials and their reciprocals, obtained using an extension of the Blissard problem, a generalized form of the Laplace transform is shown.

Some examples of the newly introduced entities are shown in Appendix A (for the Bernoulli case) and Appendix B (for the Euler case) at the end of this article, obtained by the third author using the computer algebra system Mathematica©.

2. Exponential Functions

It is quite obvious to point out that the definition of the exponential function e^x is strictly connected with the derivative operator $D_x = d/dx$ as it is an eigenfunction of this operator, satisfying

$$D_x e^{\alpha x} = \alpha e^{\alpha x}, \quad \forall \alpha \in \mathbf{C}. \tag{1}$$

According to this property, and using the McLaurin expansion, the Euler constant follows as the sum of the series: $e = \sum_{k=0}^{\infty} \frac{1}{k!}$.

Then, using an operator \mathcal{D}_x , different from D_x , it is possible to introduce the definition.

Definition 1. $\varphi(x)$ is called a true exponential function if the following eigenvalue property is satisfied:

$$\mathcal{D}_x \varphi(\alpha x) = \alpha \varphi(\alpha x), \quad \forall \alpha \in \mathbf{C}. \tag{2}$$

In the forthcoming subsections, we show some possible operators and true exponential functions of this type.

2.1. The Fractional Exponential Function

An example of such operators is given, for any real number $\alpha > 0$, by the fractional derivative D_x^α , defined by the Euler equation, falling as a special case, in the definition of a fractional derivative introduced by Caputo:

$$D_x^\alpha x^n = \begin{cases} \frac{\Gamma(n+1)}{\Gamma(n+1-\alpha)} x^{n-\alpha}, & \text{if } n > [\alpha] - 1, \\ 0, & \text{if } n = 0, 1, \dots, [\alpha] - 1, \end{cases} \tag{3}$$

where $n \geq 0$ and $[\alpha]$ denote the ceiling function (the smallest integer greater than or equal to α).

If c is a constant, then $D_x^\alpha c = 0$.

We recall that the Caputo derivative is defined as follows [13]:

$$D_{a+}^\alpha f(x) = \begin{cases} \frac{1}{\Gamma(m-\alpha)} \int_a^x \frac{f^{(m)}(\tau)}{(x-\tau)^{\alpha-m+1}} d\tau, & \text{where } m = [\alpha], \text{ if } \alpha \notin \mathbb{N} \\ f^{(\alpha)}(x), & \text{if } \alpha \in \mathbb{N}, \end{cases}$$

and reduces to the above Equation (3) when $a = 0$ and $f(x) = x^n$.

The fractional exponential function (depending on the parameter α) defined by

$$\text{Exp}_\alpha(t) = 1 + \frac{t^\alpha}{\Gamma(\alpha+1)} + \frac{t^{2\alpha}}{\Gamma(2\alpha+1)} + \dots + \frac{t^{n\alpha}}{\Gamma(n\alpha+1)} + \dots, \tag{4}$$

is an eigenfunction of the operator D_x^α since

$$D_x^\alpha \text{Exp}_\alpha(xt) = t^\alpha \text{Exp}_\alpha(xt). \tag{5}$$

For any $\alpha, 0 < \alpha < 1$, an analog of the Euler constant for the fractional exponential function is

$$\mathcal{E}_\alpha = \sum_{n=0}^\infty \frac{1}{\Gamma(n\alpha+1)}. \tag{6}$$

Remark 1. Note that the pseudo-exponential functions introduced in [14], and those connected to the Mittag-Leffler function [15], do not satisfy the eigenvalue property of the classical exponential, so they cannot be considered as true exponentials in the sense we consider here.

In [10], we have exploited the fractional exponential function (4) in order to introduce fractional index versions of fractional Bernoulli and Euler numbers and polynomials.

Remark 2. Note that what we call fractional Bernoulli and Euler polynomials are actually functions and not polynomials in the strict sense, but since they are combinations of monomials with fractional powers, it seems more appropriate for us to retain the name polynomials, as shorthand for fractional power polynomials. This terminology is used in what follows.

2.2. Fractional Index Bernoulli Numbers and Polynomials

We introduced the following definitions:

1. For the fractional index numbers $B_{\alpha,k}$, where α is a fractional number,

$$\frac{x^\alpha}{\text{Exp}_\alpha(x) - 1} = \sum_{k=0}^\infty B_{\alpha,k} \frac{x^{k\alpha}}{\Gamma(k\alpha+1)}; \tag{7}$$

2. For the fractional index polynomials $B_{\alpha,k}(z)$,

$$\frac{x^\alpha \text{Exp}_\alpha(xz)}{\text{Exp}_\alpha(x) - 1} = \sum_{k=0}^\infty B_{\alpha,k}(z) \frac{x^{k\alpha}}{\Gamma(k\alpha+1)}. \tag{8}$$

Then, we have found the result (see [10]).

Theorem 1. *The Bernoulli numbers with fractional indices $B_{\alpha,k}$ can be sequentially computed by solving the triangular system*

$$B_{\alpha,0} = \Gamma(\alpha + 1),$$

$$\sum_{k=1}^n \frac{B_{\alpha,(n-k)}}{\Gamma(k\alpha + 1) \Gamma((n-k)\alpha + 1)} = 0, \quad \forall n \geq 1. \tag{9}$$

2.3. Fractional Index Bernoulli Polynomials

Starting from (8), we have found the following (see [10]).

Theorem 2. *The fractional index Bernoulli polynomials $B_{\alpha,n}(z)$ can be sequentially constructed by solving the triangular system*

$$B_{\alpha,0}(z) = \Gamma(\alpha + 1),$$

$$\sum_{k=0}^n \frac{B_{\alpha,(n-k)}(z) \Gamma(n\alpha + 1)}{\Gamma((k+1)\alpha + 1) \Gamma((n-k)\alpha + 1)} = z^{n\alpha}, \quad \forall n \geq 1. \tag{10}$$

2.4. Fractional Index Euler Numbers and Polynomials

In a similar way, we have the definitions:

3. For the fractional index Euler numbers $E_{\alpha,k}$,

$$\frac{2 \text{Exp}_{\alpha}(x)}{\text{Exp}_{\alpha}(2x) + 1} = \sum_{k=0}^{\infty} E_{\alpha,k} \frac{x^{k\alpha}}{\Gamma(k\alpha + 1)}; \tag{11}$$

4. For the fractional index Euler polynomials $E_{\alpha,k}(z)$,

$$\frac{2 \text{Exp}_{\alpha}(xz)}{\text{Exp}_{\alpha}(2x) + 1} = \sum_{k=0}^{\infty} E_{\alpha,k}(z) \frac{x^{k\alpha}}{\Gamma(k\alpha + 1)}. \tag{12}$$

In [10], we have proven the results.

Theorem 3. *The Euler numbers with fractional indices $E_{\alpha,k}$ can be sequentially computed by solving the triangular system*

$$E_{\alpha,0} = 1,$$

$$E_{\alpha,n} + \sum_{k=0}^n \frac{2^{k\alpha} \Gamma(n\alpha + 1) E_{\alpha,n-k}}{\Gamma(k\alpha + 1) \Gamma((n-k)\alpha + 1)} = 2. \tag{13}$$

Theorem 4. *The fractional index Euler polynomials $E_{\alpha,n}(z)$ can be sequentially constructed by solving the triangular system*

$$E_{\alpha,0}(z) = 1,$$

$$\frac{1}{2} \left[E_{\alpha,n}(z) + \sum_{k=0}^n \frac{2^{(n-k)\alpha} \Gamma(n\alpha + 1) E_{\alpha,k}(z)}{\Gamma((n-k)\alpha + 1) \Gamma(k\alpha + 1)} \right] = z^{n\alpha}. \tag{14}$$

3. The Laguerre-Type Exponential Functions

Other examples of differential operators that gave rise to exponential functions are given by the Laguerre-type derivatives recalled in [11] and were studied with some applications in [12].

Starting from the first-order case, we recall that the operator $D_L := D_x x D_x = D_x + x D_x^2$ introduces a linear differential isomorphism acting on the space of the analytic functions of the variable x , which creates a parallel structure within this space. This allows us to easily derive differentiation properties.

The relevant exponential function is given by

$$e_1(x) := \sum_{k=0}^{\infty} \frac{x^k}{(k!)^2} = C_0(-x), \tag{15}$$

where $C_0(x)$ is the Tricomi function of order zero, satisfying the eigenvalue property

$$D_L e_1(ax) = a e_1(ax), \quad \forall a \in \mathbf{C}. \tag{16}$$

Iterating the Laguerre derivative, an endless cycle of construction at higher levels of differentiation occurs, showcasing a great cycle that sometimes appears within mathematical structures.

The n th-order Laguerre-type exponential is defined as

$$e_n(x) := \sum_{k=0}^{\infty} \frac{x^k}{(k!)^{n+1}}. \tag{17}$$

It is an eigenfunction of the operator

$$D_{nL} := D x \cdots D x D x D = D(xD + x^2 D^2 + \cdots + x^n D^n) = S(n+1, 1)D + S(n+1, 2)x D^2 + \cdots + S(n+1, n+1)x^n D^{n+1}, \tag{18}$$

where $S(n, k)$ denotes Stirling numbers of the second kind.

For every integer n , the function (17) is an exponential function since it satisfies the eigenvalue property

$$D_{nL} e_n(ax) = a e_n(ax). \tag{19}$$

The corresponding Euler constants are given by the sum of the series

$$e_n := \sum_{k=0}^{\infty} \frac{1}{(k!)^{n+1}}. \tag{20}$$

Further information can be found in the article cited above [12].

Remark 3. For completeness, we recall that the operators $D_L = D x D$ and their iterates as $D_{nL} = D x D x D x \cdots D x D$ can be considered as particular cases of the hyper-Bessel differential operators when $\alpha_0 = \alpha_1 = \cdots = \alpha_n = 1$ (the special case considered in operational calculus by Ditkin and Prudnikov [16]). In general, the Bessel-type differential operators of arbitrary order n were introduced by Dimovski in 1966 [17] and were later called, by Kiryakova, hyper-Bessel operators because they are closely related to their eigenfunctions, called hyper-Bessel by Delerue [18] in 1953. These operators were studied in 1994 by Kiryakova in her book ([19], Ch. 3).

The Fractional Laguerre-Type Exponential Functions

Combining the results in the preceding sections, we can also consider the fractional Laguerre-type exponential function

$$e_{\alpha,1}(x) := \sum_{k=0}^{\infty} \frac{x^{k\alpha}}{[\Gamma(k\alpha + 1)]^2}, \tag{21}$$

which is an eigenfunction of the operator $D_L^\alpha := D_x^\alpha x^\alpha D_x^\alpha$ since

$$D_x^\alpha x^\alpha D_x^\alpha e_{\alpha,1}(tx) = t^\alpha e_{\alpha,1}(tx), \tag{22}$$

and in general,

$$e_{\alpha,n}(x) := \sum_{k=0}^{\infty} \frac{x^{k\alpha}}{[\Gamma(k\alpha + 1)]^{n+1}}, \tag{23}$$

for which the operator $D_{nL}^\alpha := D_x^\alpha x^\alpha \cdots D_x^\alpha x^\alpha D_x^\alpha x^\alpha D_x^\alpha$ ($n + 1$ fractional derivatives) is such that

$$D_{nL}^\alpha e_{\alpha,n}(tx) = t^\alpha e_{\alpha,n}(tx). \tag{24}$$

In what follows, we exploit the first-order Laguerre-type exponential function (21) in order to introduce the Laguerre-type Bernoulli and Euler numbers and polynomials. Of course, these new entities could be considered even in the general case of the n th-order Laguerre-type exponentials (23), but this will not be considered here since the technique is the same and the relevant formulas are much more complicated.

4. First-Order Laguerre-Type Bernoulli and Euler Numbers and Polynomials

Since we considered only the first-order case, we will call these numbers simply Laguerre-type Bernoulli and Euler numbers (and the following Laguerre-type Bernoulli and Euler polynomials).

Using their generating functions, we now introduce the following definitions.

The Laguerre-type Bernoulli numbers and polynomials

1. For the ordinary Laguerre-type Bernoulli numbers ${}_L B_k$,

$$\frac{x}{e_1(x) - 1} = \sum_{k=0}^{\infty} {}_L B_k \frac{x^k}{[k!]^2}; \tag{25}$$

2. For the ordinary Laguerre-type Bernoulli polynomials ${}_L B_k(z)$,

$$\frac{x e_1(xz)}{e_1(x) - 1} = \sum_{k=0}^{\infty} {}_L B_k(z) \frac{x^k}{[k!]^2}. \tag{26}$$

The fractional Laguerre-type Bernoulli numbers and polynomials

3. For the fractional Laguerre-type Bernoulli numbers ${}_L B_{\alpha,k}$, where α is a fractional number,

$$\frac{x^\alpha}{e_{\alpha,1}(x) - 1} = \sum_{k=0}^{\infty} {}_L B_{\alpha,k} \frac{x^{k\alpha}}{[\Gamma(k\alpha + 1)]^2}; \tag{27}$$

4. For the fractional index Laguerre-type Bernoulli polynomials ${}_L B_{\alpha,k}(z)$,

$$\frac{x^\alpha e_{\alpha,1}(xz)}{e_{\alpha,1}(x) - 1} = \sum_{k=0}^{\infty} {}_L B_{\alpha,k}(z) \frac{x^{k\alpha}}{[\Gamma(k\alpha + 1)]^2}. \tag{28}$$

The Laguerre-type Euler numbers and polynomials

5. For the ordinary Laguerre-type Euler numbers ${}_L E_k$,

$$\frac{2 e_1(x)}{e_1(2x) + 1} = \sum_{k=0}^{\infty} {}_L E_k \frac{x^k}{[k!]^2}; \tag{29}$$

6. For the ordinary Laguerre-type Euler polynomials ${}_L E_k(z)$,

$$\frac{2 e_1(x z)}{e_1(2 x)+1}=\sum_{k=0}^{\infty} {}_L E_k(z) \frac{x^k}{[k!]^2} . \tag{30}$$

The fractional Laguerre-type Euler numbers and polynomials

7. For the fractional Laguerre-type Euler numbers ${}_L E_{\alpha, k}$, where α is a fractional number

$$\frac{2 e_{\alpha, 1}(x)}{e_{\alpha, 1}(2 x)+1}=\sum_{k=0}^{\infty} {}_L E_{\alpha, k} \frac{x^{k \alpha}}{[\Gamma(k \alpha+1)]^2} ; \tag{31}$$

8. For the fractional index Laguerre-type Euler polynomials ${}_L E_{\alpha, k}(z)$,

$$\frac{2 e_{\alpha, 1}(x z)}{e_{\alpha, 1}(2 x)+1}=\sum_{k=0}^{\infty} {}_L E_{\alpha, k}(z) \frac{x^{k \alpha}}{[\Gamma(k \alpha+1)]^2} . \tag{32}$$

5. The Reciprocal of the Fractional Laguerre-Type Exponential Function

We consider the function $e_{\alpha, 1}(a ; t)$, where the symbol $a \equiv\left\{a_n\right\}$ denotes the sequence of coefficients, according to the position

$$e_{\alpha, 1}(a ; t)=1+a_1 \frac{t^{\alpha}}{[\Gamma(\alpha+1)]^2}+a_2 \frac{t^{2 \alpha}}{[\Gamma(2 \alpha+1)]^2}+\cdots+a_n \frac{t^{n \alpha}}{[\Gamma(n \alpha+1)]^2}+\cdots . \tag{33}$$

The equation

$$\frac{1}{e_{\alpha, 1}(a ; t)}=b_0+b_1 \frac{t^{\alpha}}{[\Gamma(\alpha+1)]^2}+b_2 \frac{t^{2 \alpha}}{[\Gamma(2 \alpha+1)]^2}+\cdots+b_n \frac{t^{n \alpha}}{[\Gamma(n \alpha+1)]^2}+\cdots \tag{34}$$

in terms of the unknown sequence $b \equiv\left\{b_n\right\}$ can be solved using Bell’s polynomials.

General Result

In the literature, there exists the following general result [20].

Consider the sequences $a :=\left\{a_k\right\}=\left(a_0, a_1, a_2, a_3, \dots\right)$, and $b :=\left\{b_k\right\}=\left(b_0, b_1, b_2, b_3, \dots\right)$. Using the umbral formalism (that is, letting $a_k \equiv a^k$ and $b_k \equiv b^k$), the solution of the equation

$$\frac{1}{\sum_{n=0}^{\infty} \frac{a^n t^{n \alpha}}{[\Gamma(n \alpha+1)]^2}}=\sum_{n=0}^{\infty} \frac{b^n t^{n \alpha}}{[\Gamma(n \alpha+1)]^2}, \quad \text { i.e. } \quad e_{\alpha, 1}(a ; t) e_{\alpha, 1}(b ; t)=1 ,$$

according to the Faà di Bruno formula, is given by

$$b_n=\frac{[\Gamma(n \alpha+1)]^2}{n!} \sum_{k=0}^n(-1)^k k! a_0^{-(k+1)} \cdot B_{n, k}\left(\frac{1! a_1}{[\Gamma(\alpha+1)]^2}, \frac{2! a_2}{[\Gamma(2 \alpha+1)]^2}, \dots, \frac{(n-k+1)! a_{n-k+1}}{[\Gamma((n-k+1) \alpha+1)]^2}\right), \quad(\forall n \geq 0),$$

where $B_{n, k}$ are partial Bell polynomials [20,21].

In our case, we have $a_0=1$, and we have to consider the reciprocal of Equation (33), i.e.,

$$\frac{1}{1+a_1 \frac{t^{\alpha}}{[\Gamma(\alpha+1)]^2}+a_2 \frac{t^{2 \alpha}}{[\Gamma(2 \alpha+1)]^2}+a_3 \frac{t^{3 \alpha}}{[\Gamma(3 \alpha+1)]^2}+\dots} \quad(t \geq 0) .$$

Then, according to the above general result, we find

$$\frac{1}{\sum_{n=0}^{\infty} \frac{a_n t^{n\alpha}}{[\Gamma(n\alpha + 1)]^2}} = \sum_{n=0}^{\infty} \frac{b_n t^{n\alpha}}{[\Gamma(n\alpha + 1)]^2} = \sum_{n=0}^{\infty} \frac{t^{n\alpha}}{n!} \sum_{k=0}^n (-1)^k k! B_{n,k} \left(\frac{1! a_1}{[\Gamma(\alpha + 1)]^2}, \frac{2! a_2}{[\Gamma(2\alpha + 1)]^2}, \dots, \frac{(n - k + 1)! a_{n-k+1}}{[\Gamma((n - k + 1)\alpha + 1)]^2} \right). \tag{35}$$

Therefore, for $\alpha = 1/2$ and $a_n = 1/(n + 1)$, the first few values of the b_n coefficients are found to be

$$\begin{aligned} b_0 &= 1 \\ b_1 &= -\frac{1}{2}, \\ b_2 &= -\frac{1}{3} + \frac{4}{\pi^2}, \\ b_3 &= \frac{1}{2} - \frac{9}{2\pi^2}, \\ b_4 &= \frac{11}{45} + \frac{64}{\pi^4} - \frac{80}{9\pi^2}, \\ b_5 &= -\frac{49}{64} - \frac{225}{2\pi^4} + \frac{75}{4\pi^2}, \\ b_6 &= -\frac{29}{105} + \frac{2304}{\pi^6} - \frac{448}{\pi^4} + \frac{5476}{225\pi^2}. \end{aligned}$$

6. Laguerre-Type Fractional-Order Laplace Transforms

Using the above definition of the reciprocal of the fractional Laguerre-type exponential function, we can introduce a fractional-order Laguerre-type Laplace transform by setting

$$\begin{aligned} {}_{L_1}\mathcal{L}_\alpha(f) &:= \int_0^\infty f(t) [e_{\alpha,1}(st)]^{-1} dt = {}_{L_1}\mathcal{F}_\alpha(s) = \\ &\int_0^\infty f(t) \left[\sum_{n=0}^{\infty} \frac{(st)^{n\alpha}}{n!} \sum_{k=0}^n (-1)^k k! B_{n,k} \left(\frac{1!}{[\Gamma(\alpha + 1)]^2}, \dots, \frac{(n - k + 1)!}{[\Gamma((n - k + 1)\alpha + 1)]^2} \right) \right] dt. \end{aligned} \tag{36}$$

In what follows, we make a comparison among the Laguerre-type Laplace transform of the assigned functions and the fractional order Laguerre-type Laplace transforms of order $\alpha = 1/2$ and $\alpha = 3/2$.

As shown in the obtained results, in all cases, the graphs of the modulus and argument of the Laguerre-type Laplace transform lie between the corresponding graphs of the two considered fractional order Laguerre-type Laplace transforms. This provides graphical evidence of the monotonicity property satisfied by the fractional order Laguerre-type Laplace transforms.

Lastly, we write for completeness the general form of the Laguerre-type fractional-order Laplace transforms, using the m -th order Laguerre-type exponentials, which is written as

$$\begin{aligned} {}_{L_m}\mathcal{L}_\alpha(f) &:= \int_0^\infty f(t) [e_{\alpha,m}(st)]^{-1} dt = {}_{L_m}\mathcal{F}_\alpha(s) = \\ &\int_0^\infty f(t) \left[\sum_{n=0}^{\infty} \frac{(st)^{n\alpha}}{n!} \sum_{k=0}^n (-1)^k k! B_{n,k} \left(\frac{1!}{[\Gamma(\alpha + 1)]^{m+1}}, \dots, \frac{(n - k + 1)!}{[\Gamma((n - k + 1)\alpha + 1)]^{m+1}} \right) \right] dt. \end{aligned} \tag{37}$$

7. Numerical Examples

7.1. Example 1

Consider the fractional Laguerre-type Laplace transforms ${}_L\mathcal{F}_{1/2}, {}_L\mathcal{F}_{3/2}$ of the Bessel function $J_0(2\sqrt{t})$ compared with the Laguerre-type LT ${}_L\mathcal{F} = \mathcal{F}_1$ of the same function. The case of the modulus, assuming $s = \sigma + 5i$, is depicted in Figure 1, and the case of the argument, is shown in Figure 2. The case of the modulus, assuming $s = 5 + i\omega$, is shown in Figure 3, and the case of the argument is shown in Figure 4.




Figure 1. Comparing the fractional Laguerre-type LTs ${}_L\mathcal{F}_{1/2}, {}_L\mathcal{F}_1, {}_L\mathcal{F}_{3/2}$ of the function $J_0(2\sqrt{t})$ —the case of the modulus, assuming $s = \sigma + 5i$.




Figure 2. Comparing the fractional Laguerre-type LTs ${}_L\mathcal{F}_{1/2}, {}_L\mathcal{F}_1, {}_L\mathcal{F}_{3/2}$ of the function $J_0(2\sqrt{t})$ —the case of the argument, assuming $s = \sigma + 5i$.




Figure 3. Comparing the fractional Laguerre-type LTs ${}_L\mathcal{F}_{1/2}, {}_L\mathcal{F}_1, {}_L\mathcal{F}_{3/2}$ of the function $J_0(2\sqrt{t})$ —the case of the modulus, assuming $s = 5 + i\omega$.




Figure 4. Comparing the fractional Laguerre-type LTs ${}_L\mathcal{F}_{1/2}, {}_L\mathcal{F}_1, {}_L\mathcal{F}_{3/2}$ of the function $J_0(2\sqrt{t})$ —the case of the argument, assuming $s = 5 + i\omega$.

7.2. Example 2

Consider the fractional Laguerre-type Laplace transforms ${}_L\mathcal{F}_{1/2}, {}_L\mathcal{F}_{3/2}$ of the function $\exp(-t^2)$ compared with the Laguerre-type LT $F = {}_L\mathcal{F}_1$ of the same function. The case of the modulus, assuming $s = \sigma + i$, is depicted in Figure 5, and the case of the argument is shown in Figure 6. The case of the modulus, assuming $s = 1 + i\omega$, is shown in Figure 7, and the case of the argument is shown in Figure 8.




Figure 5. Comparing the fractional Laguerre-type LTs ${}_L\mathcal{F}_{1/2}, {}_L\mathcal{F}_1, {}_L\mathcal{F}_{3/2}$ of the function $\exp(-t^2)$ —the case of the modulus, assuming $s = \sigma + i$.




Figure 6. Comparing the fractional Laguerre-type LTs ${}_L\mathcal{F}_{1/2}, {}_L\mathcal{F}_1, {}_L\mathcal{F}_{3/2}$ of the function $\exp(-t^2)$ —the case of the argument, assuming $s = \sigma + i$.




Figure 7. Comparing the fractional Laguerre-type LTs ${}_L\mathcal{F}_{1/2}$, ${}_L\mathcal{F}_1$, ${}_L\mathcal{F}_{3/2}$ of the function $\exp(-t^2)$ —the case of the modulus, assuming $s = 1 + i\omega$.




Figure 8. Comparing the fractional Laguerre-type LTs ${}_L\mathcal{F}_{1/2}$, ${}_L\mathcal{F}_1$, ${}_L\mathcal{F}_{3/2}$ of the function $\exp(-t^2)$ —the case of the argument, assuming $s = 1 + i\omega$.

8. Conclusions

We have shown that using the Laguerre-type exponentials and their fractional versions, it is possible to define the Laguerre-type fractional forms of the classical Bernoulli and Euler numbers and polynomials. The reciprocal of these Laguerre-type exponentials can also be used in order to generalize the ordinary Laplace transform.

All these extensions show further applications of the Laguerre-type derivatives, which introduce, within the space of analytic functions, a differential isomorphism producing the possibility of finding new identities. This means that all differential equations can be reproduced at different levels of differentiation, maintaining their formal structure, a sort of invariance of differential identities with respect to their Laguerre-type versions. Some applications of this technique have been shown in Ref. [12] and will be further exploited in subsequent articles.

In this paper, we have shown numerical values of the newly considered entities and examples of the relevant Laguerre-type Laplace transforms computed by using the computer algebra system Mathematica©.

Author Contributions: Conceptualization, P.E.R., R.S. and D.C.; methodology, P.E.R., R.S. and D.C.; software, D.C.; validation, P.E.R., R.S. and D.C.; investigation, P.E.R., R.S. and D.C.; writing—original draft, P.E.R.; supervision, P.E.R. All authors have read and agreed to the published version of this manuscript.

Funding: This research received no external funding.

Data Availability Statement: Data is contained within the article.

Acknowledgments: Thanks to the anonymous referee for helpful comments.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A. Examples of Laguerre-Type Fractional First-Order Bernoulli Numbers and Fractional Polynomials

The first few values of the Laguerre-type fractional first-order Bernoulli numbers for $\alpha = 1/2$ are reported in Figure A1.

$$\begin{aligned}
 {}_L B_{\frac{1}{2},0}^1 &= \frac{\pi}{4} \\
 {}_L B_{\frac{1}{2},1}^1 &= -\frac{\pi^3}{64} \\
 {}_L B_{\frac{1}{2},2}^1 &= -\frac{\pi}{9} + \frac{\pi^3}{64} \\
 {}_L B_{\frac{1}{2},3}^1 &= -\frac{\pi^3(-92 + 9\pi^2)}{4096} \\
 {}_L B_{\frac{1}{2},4}^1 &= \frac{256\pi}{2025} - \frac{5\pi^3}{96} + \frac{\pi^5}{256} \\
 {}_L B_{\frac{1}{2},5}^1 &= -\frac{\pi^3(44224 + 75\pi^2(-148 + 9\pi^2))}{786432} \\
 {}_L B_{\frac{1}{2},6}^1 &= -\frac{27136\pi}{99225} + \frac{4571\pi^3}{19200} - \frac{11\pi^5}{256} + \frac{9\pi^7}{4096} \\
 {}_L B_{\frac{1}{2},7}^1 &= -\frac{\pi^3(-305461696 + 441\pi^2(325744 + 675\pi^2(-68 + 3\pi^2)))}{1358954496} \\
 {}_L B_{\frac{1}{2},8}^1 &= \frac{21889024\pi}{22325625} - \frac{972703\pi^3}{635040} + \frac{941\pi^5}{1920} - \frac{29\pi^7}{512} + \frac{9\pi^9}{4096} \\
 {}_L B_{\frac{1}{2},9}^1 &= -\frac{\pi^3(320770368512 + 225\pi^2(-1089244928 + 1323\pi^2(199072 + 75\pi^2(-260 + 9\pi^2))))}{241591910400} \\
 {}_L B_{\frac{1}{2},10}^1 &= -\frac{5185601536\pi}{972504225} + \frac{511345159\pi^3}{38102400} - \frac{3383785\pi^5}{508032} + \frac{47647\pi^7}{36864} - \frac{225\pi^9}{2048} + \frac{225\pi^{11}}{65536}
 \end{aligned}$$

Figure A1. Laguerre-type Bernoulli numbers ${}_L B_{1/2,k}$ for $0 \leq k \leq 10$.

The Laguerre-type fractional first-order Bernoulli numbers for $\alpha = 0.2, 0.4, 0.6, 0.8$ and $0 \leq k \leq 10$ are shown in Figure A2.

${}_L B_{0.2,0} \approx 0.843034$	${}_L B_{0.4,0} \approx 0.787237$	${}_L B_{0.6,0} \approx 0.79837$	${}_L B_{0.8,0} \approx 0.867476$
${}_L B_{0.2,1} \approx -0.761079$	${}_L B_{0.4,1} \approx -0.562418$	${}_L B_{0.6,1} \approx -0.419184$	${}_L B_{0.8,1} \approx -0.319395$
${}_L B_{0.2,2} \approx 0.0602827$	${}_L B_{0.4,2} \approx 0.119564$	${}_L B_{0.6,2} \approx 0.143879$	${}_L B_{0.8,2} \approx 0.146344$
${}_L B_{0.2,3} \approx 0.0415215$	${}_L B_{0.4,3} \approx 0.0424666$	${}_L B_{0.6,3} \approx -0.00145996$	${}_L B_{0.8,3} \approx -0.0618663$
${}_L B_{0.2,4} \approx 0.0261019$	${}_L B_{0.4,4} \approx -0.00180359$	${}_L B_{0.6,4} \approx -0.034349$	${}_L B_{0.8,4} \approx -0.00193884$
${}_L B_{0.2,5} \approx 0.0140814$	${}_L B_{0.4,5} \approx -0.0163987$	${}_L B_{0.6,5} \approx 0.00457052$	${}_L B_{0.8,5} \approx 0.0648545$
${}_L B_{0.2,6} \approx 0.00531646$	${}_L B_{0.4,6} \approx -0.0102322$	${}_L B_{0.6,6} \approx 0.0309876$	${}_L B_{0.8,6} \approx -0.10911$
${}_L B_{0.2,7} \approx -0.000469608$	${}_L B_{0.4,7} \approx 0.00381348$	${}_L B_{0.6,7} \approx -0.0123209$	${}_L B_{0.8,7} \approx 0.0206612$
${}_L B_{0.2,8} \approx -0.0036569$	${}_L B_{0.4,8} \approx 0.012409$	${}_L B_{0.6,8} \approx -0.0576747$	${}_L B_{0.8,8} \approx 0.537441$
${}_L B_{0.2,9} \approx -0.00471035$	${}_L B_{0.4,9} \approx 0.00732876$	${}_L B_{0.6,9} \approx 0.0477632$	${}_L B_{0.8,9} \approx -1.96528$
${}_L B_{0.2,10} \approx -0.00416126$	${}_L B_{0.4,10} \approx -0.0087206$	${}_L B_{0.6,10} \approx 0.181364$	${}_L B_{0.8,10} \approx 1.56549$

Figure A2. Sequences of Laguerre-type Bernoulli numbers ${}_L B_{\alpha,k}$ with fractional indices $\alpha = 0.2, 0.4, 0.6, 0.8$ for $0 \leq k \leq 10$.

A set of the Laguerre-type fractional Bernoulli numbers ${}_L B_{\alpha,k}$ is shown in Figure A3.

${}_L B_{\alpha,k}$	k=0	k=1	k=2	k=3	k=4
$\alpha=0.2$	0.843034...	-0.761079...	0.060283...	0.041522...	0.026102...
$\alpha=0.4$	0.787237...	-0.562418...	0.119564...	0.042467...	-0.001804...
$\alpha=0.6$	0.798370...	-0.419184...	0.143879...	-0.001460...	-0.034349...
$\alpha=0.8$	0.867476...	-0.319395...	0.146344...	-0.061866...	-0.001939...
$\alpha=1.0$	1.000000...	-0.250000...	0.138889...	-0.125000...	0.154444...

Figure A3. Table of the fractional index Laguerre-type Bernoulli numbers ${}_L B_{\alpha,k}$.

The first few values of the Laguerre-type fractional Bernoulli fractional polynomials ${}_L B_{1/2,k}(z)$ are reported in Figure A4.

$$\begin{aligned}
 {}_L B_{\frac{1}{2},0}(z) &= \frac{\pi}{4} \\
 {}_L B_{\frac{1}{2},1}(z) &= -\frac{1}{64} \pi (\pi^2 - 16 \sqrt{z}) \\
 {}_L B_{\frac{1}{2},2}(z) &= \frac{\pi^3}{64} + \frac{1}{36} \pi (-4 - 9 \sqrt{z} + 9z) \\
 {}_L B_{\frac{1}{2},3}(z) &= \frac{-9 \pi^5 + 4 \pi^3 (23 + 36 \sqrt{z} - 36z) + 1024 \pi (-1 + z) \sqrt{z}}{4096} \\
 {}_L B_{\frac{1}{2},4}(z) &= \frac{\pi^5}{256} + \frac{1}{96} \pi^3 (-5 - 6 \sqrt{z} + 6z) + \pi \left(\frac{256}{2025} + \frac{1}{36} (23 \sqrt{z} - 16z - 16z^{3/2} + 9z^2) \right) \\
 {}_L B_{\frac{1}{2},5}(z) &= \frac{1}{2359296} \left(-2025 \pi^7 + 65536 \pi (-1 + z) \sqrt{z} (-16 + 9z) - \right. \\
 &\quad \left. 900 \pi^5 (-37 - 36 \sqrt{z} + 36z) - 192 \pi^3 (691 + 75 (30 \sqrt{z} - 23z - 16z^{3/2} + 9z^2)) \right) \\
 {}_L B_{\frac{1}{2},6}(z) &= \frac{9 \pi^7}{4096} + \frac{1}{256} \pi^5 (-11 - 9 \sqrt{z} + 9z) + \pi \left(-\frac{27136}{99225} - \frac{691 \sqrt{z}}{300} + \frac{256z}{225} + \frac{23z^{3/2}}{9} - z^2 - \frac{16z^{5/2}}{25} + \frac{z^3}{4} \right) + \\
 &\quad \frac{\pi^3 (4571 + 300 (37 \sqrt{z} - 30z - 16z^{3/2} + 9z^2))}{19200} \\
 {}_L B_{\frac{1}{2},7}(z) &= \frac{1}{1358954496} \left(-893025 \pi^9 - 1190700 \pi^7 (-17 - 12 \sqrt{z} + 12z) + 4194304 \pi (-1 + z) \sqrt{z} (424 + 9z (-40 + 9z)) - \right. \\
 &\quad \left. 7056 \pi^5 (20359 + 900 (44 \sqrt{z} - 37z - 16z^{3/2} + 9z^2)) - \right. \\
 &\quad \left. 64 \pi^3 (-4772839 + 5292 (-1 + \sqrt{z}) \sqrt{z} (4571 + 1807 \sqrt{z} - 2193z - 468z^{3/2} + 300z^2)) \right)
 \end{aligned}$$

Figure A4. Fractional index Laguerre-type Bernoulli polynomials ${}_L B_{1/2,k}(z)$.

Appendix B. Examples of Laguerre-Type Fractional First-Order Euler Numbers and Fractional Polynomials

The first few values of the Laguerre-type fractional first-order Euler numbers for $\alpha = 1/2$ are reported in Figure A5.

$$\begin{aligned}
 {}_L E_{\frac{1}{2},0} &= 1 \\
 {}_L E_{\frac{1}{2},1} &= 1 - \frac{1}{\sqrt{2}} \\
 {}_L E_{\frac{1}{2},2} &= \frac{8 - 8\sqrt{2}}{\pi^2} \\
 {}_L E_{\frac{1}{2},3} &= \frac{1}{8} \left(-10 + \sqrt{2} \right) - \frac{9 \left(-2 + \sqrt{2} \right)}{\pi^2} \\
 {}_L E_{\frac{1}{2},4} &= -1 - \frac{256 \left(-1 + \sqrt{2} \right)}{\pi^4} + \frac{64 \left(-1 + 3\sqrt{2} \right)}{9\pi^2} \\
 {}_L E_{\frac{1}{2},5} &= \frac{1}{128} \left(228 + 319\sqrt{2} \right) - \frac{450 \left(-2 + \sqrt{2} \right)}{\pi^4} + \frac{25 \left(-14 + 3\sqrt{2} \right)}{4\pi^2} \\
 {}_L E_{\frac{1}{2},6} &= 6 - \frac{18432 \left(-1 + \sqrt{2} \right)}{\pi^6} + \frac{1024 \left(-1 + 2\sqrt{2} \right)}{\pi^4} - \frac{8 \left(3617 + 680\sqrt{2} \right)}{225\pi^2} \\
 {}_L E_{\frac{1}{2},7} &= \frac{1}{512} \left(4040 - 12779\sqrt{2} \right) + \frac{49 \left(-518400 \left(-2 + \sqrt{2} \right) + 7200 \left(-18 + 5\sqrt{2} \right) \pi^2 + 7 \left(124 + 609\sqrt{2} \right) \pi^4 \right)}{576\pi^6} \\
 {}_L E_{\frac{1}{2},8} &= -31 + \frac{1}{11025\pi^8} 512 \left(-50803200 \left(-1 + \sqrt{2} \right) + \right. \\
 &\quad \left. 1411200 \left(-3 + 5\sqrt{2} \right) \pi^2 - 294 \left(1603 + 96\sqrt{2} \right) \pi^4 + \left(43271 - 18365\sqrt{2} \right) \pi^6 \right)
 \end{aligned}$$

Figure A5. Laguerre-type Euler numbers ${}_L E_{1/2,k}$ for $0 \leq k \leq 8$.

The Laguerre-type fractional first-order Euler numbers for $\alpha = 0.2, 0.4, 0.6, 0.8$ and $0 \leq k \leq 10$ are shown in Figure A6.

${}_L E_{0.2,0} \approx 1.$	${}_L E_{0.4,0} \approx 1.$	${}_L E_{0.6,0} \approx 1.$	${}_L E_{0.8,0} \approx 1.$
${}_L E_{0.2,1} \approx 0.425651$	${}_L E_{0.4,1} \approx 0.340246$	${}_L E_{0.6,1} \approx 0.242142$	${}_L E_{0.8,1} \approx 0.129449$
${}_L E_{0.2,2} \approx 0.0694483$	${}_L E_{0.4,2} \approx -0.184762$	${}_L E_{0.6,2} \approx -0.498206$	${}_L E_{0.8,2} \approx -0.821788$
${}_L E_{0.2,3} \approx -0.143665$	${}_L E_{0.4,3} \approx -0.458549$	${}_L E_{0.6,3} \approx -0.452782$	${}_L E_{0.8,3} \approx 0.963643$
${}_L E_{0.2,4} \approx -0.244102$	${}_L E_{0.4,4} \approx -0.267725$	${}_L E_{0.6,4} \approx 1.50158$	${}_L E_{0.8,4} \approx 5.1334$
${}_L E_{0.2,5} \approx -0.247917$	${}_L E_{0.4,5} \approx 0.479429$	${}_L E_{0.6,5} \approx 2.62412$	${}_L E_{0.8,5} \approx -40.429$
${}_L E_{0.2,6} \approx -0.170415$	${}_L E_{0.4,6} \approx 1.29637$	${}_L E_{0.6,6} \approx -12.4234$	${}_L E_{0.8,6} \approx 13.4821$
${}_L E_{0.2,7} \approx -0.033849$	${}_L E_{0.4,7} \approx 0.596329$	${}_L E_{0.6,7} \approx -34.3072$	${}_L E_{0.8,7} \approx 2578.13$
${}_L E_{0.2,8} \approx 0.128122$	${}_L E_{0.4,8} \approx -3.68634$	${}_L E_{0.6,8} \approx 208.346$	${}_L E_{0.8,8} \approx -22813.4$
${}_L E_{0.2,9} \approx 0.269354$	${}_L E_{0.4,9} \approx -9.43384$	${}_L E_{0.6,9} \approx 826.97$	${}_L E_{0.8,9} \approx -168892.$
${}_L E_{0.2,10} \approx 0.334973$	${}_L E_{0.4,10} \approx -0.00135782$	${}_L E_{0.6,10} \approx -6021.41$	${}_L E_{0.8,10} \approx 6.98957 \times 10^6$

Figure A6. Sequences of Laguerre-type Euler numbers ${}_L E_{\alpha,k}$ with fractional indices $\alpha = 0.2, 0.4, 0.6, 0.8$ for $0 \leq k \leq 10$.

A set of the Laguerre-type fractional Euler numbers ${}_L E_{\alpha,k}$ is shown in Figure A7.

${}_L E_{\alpha,k}$	k=0	k=1	k=2	k=3	k=4
$\alpha=0.2$	1.000000...	0.425651...	0.069448...	-0.143665...	-0.244102...
$\alpha=0.4$	1.000000...	0.340246...	-0.184762...	-0.458549...	-0.267725...
$\alpha=0.6$	1.000000...	0.242142...	-0.498206...	-0.452782...	1.501580...
$\alpha=0.8$	1.000000...	0.129449...	-0.821788...	0.963643...	5.133400...
$\alpha=1.0$	1.000000...	0.000000...	-1.000000...	6.000000...	-31.000000...

Figure A7. Table of the fractional index Laguerre-type Euler numbers ${}_L E_{\alpha,k}$.

The first few values of the Laguerre-type fractional Euler fractional polynomials ${}_L E_{1/2,k}(z)$ are reported in Figure A8.

$$\begin{aligned}
 {}_L E_{\frac{1}{2},0}(z) &= 1 \\
 {}_L E_{\frac{1}{2},1}(z) &= -\frac{1}{\sqrt{2}} + \sqrt{z} \\
 {}_L E_{\frac{1}{2},2}(z) &= -1 + \frac{8 - 8\sqrt{2}\sqrt{z}}{\pi^2} + z \\
 {}_L E_{\frac{1}{2},3}(z) &= \frac{1}{4} (5\sqrt{2} - 9\sqrt{z}) - \frac{9(\sqrt{2} - 2\sqrt{z})}{\pi^2} - \frac{9z}{4\sqrt{2}} + z^{3/2} \\
 {}_L E_{\frac{1}{2},4}(z) &= 2 - \frac{256(-1 + \sqrt{2}\sqrt{z})}{\pi^4} + (-4 + z)z - \frac{32(11 - 10\sqrt{2}\sqrt{z} - 9z + 4\sqrt{2}z^{3/2})}{9\pi^2} \\
 {}_L E_{\frac{1}{2},5}(z) &= -\frac{450(\sqrt{2} - 2\sqrt{z})}{\pi^4} - \frac{25(-12\sqrt{2} + 22\sqrt{z} + 9\sqrt{2}z - 8z^{3/2})}{4\pi^2} + \\
 &\quad \frac{1}{128} (-456\sqrt{2} + 900\sqrt{z} + 1000\sqrt{2}z - 800z^{3/2} - 225\sqrt{2}z^2 + 128z^{5/2}) \\
 {}_L E_{\frac{1}{2},6}(z) &= \\
 &\quad -4 - \frac{18432(-1 + \sqrt{2}\sqrt{z})}{\pi^6} + (-6 + z)(-3 + z)z - \frac{256(13 - 12\sqrt{2}\sqrt{z} - 9z + 4\sqrt{2}z^{3/2})}{\pi^4} \\
 &\quad \frac{8(-4258 + 4104\sqrt{2}\sqrt{z} + 9900z - 4000\sqrt{2}z^{3/2} - 2025z^2 + 576\sqrt{2}z^{5/2})}{225\pi^2}
 \end{aligned}$$

Figure A8. Fractional index Laguerre-type Euler polynomials ${}_L E_{1/2,k}(z)$.

References

1. Srivastava, H.M. An introductory overview of Bessel polynomials, the generalized Bessel polynomials and the q-Bessel polynomials. *Symmetry* **2023**, *15*, 822. [\[CrossRef\]](#)
2. Srivastava, H.M.; Masjed-Jamei, M.; Beyki, M.R. Some new generalizations and applications of the Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi polynomials. *Rocky Mountain J. Math.* **2019**, *49*, 681–697. [\[CrossRef\]](#)
3. Srivastava, H.M.; Kurt, B.; Kurt, V. Identities and relations involving the modified degenerate Hermite-based Apostol-Bernoulli and Apostol-Euler polynomials. *Rev. Real Acad. Cienc. Exactas Fís. Natur. Ser. A Mat. (RACSAM)* **2019**, *113*, 1299–1313. [\[CrossRef\]](#)
4. He, Y.; Araci, S.; Srivastava, H.M.; Abdel-Aty, M. Higher-order convolutions for Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi polynomials. *Mathematics* **2018**, *6*, 329. [\[CrossRef\]](#)
5. Boutiche, M.A.; Rahmani, M.; Srivastava, H.M. Explicit formulas associated with some families of generalized Bernoulli and Euler polynomials. *Mediterr. J. Math.* **2017**, *14*, 89. [\[CrossRef\]](#)
6. Srivastava, H.M. Some families of generating functions associated with orthogonal polynomials and other higher transcendental functions. *Mathematics* **2022**, *10*, 3730. [\[CrossRef\]](#)
7. Srivastava, H.M. Some families of generating functions and associated hypergeometric transformations. *J. Nonlinear Convex Anal.* **2023**, *24*, 2329–2349.
8. Srivastava, H.M. A survey of some recent developments on higher transcendental functions of analytic number theory and applied mathematics. *Symmetry* **2021**, *13*, 2294. [\[CrossRef\]](#)
9. Kilbas, A.A.; Srivastava, H.M.; Trujillo, J.J. *Theory and Applications of Fractional Differential Equations, North-Holland Mathematical Studies*; Elsevier (North-Holland) Science Publishers: Amsterdam, The Netherlands; London, UK; New York, NY, USA, 2006; Volume 204.
10. Caratelli, D.; Natalini, P.; Ricci, P.E. Fractional Bernoulli and Euler Numbers and related fractional polynomials. A Symmetry in Number Theory. *Symmetry* **2023**, *15*, 1900. [\[CrossRef\]](#)
11. Dattoli, G.; Ricci, P.E. Laguerre-type exponentials, and the relevant L-circular and L-hyperbolic functions. *Georgian Math. J.* **2003**, *10*, 481–494. [\[CrossRef\]](#)
12. Ricci, P.E.; Tavkhelidze, I. An introduction to operational techniques and special polynomials. *J. Math. Sci.* **2009**, *157*, 161–189; Translated from: *Contemp. Math. Its Appl.-(Sovremennaya Matematika i ee Prilozheniya)* **2009**, *51*, 1512–1712. (In Russian) [\[CrossRef\]](#)
13. Beghin, L.; Caputo, M. Commutative and associative properties of the Caputo fractional derivative and its generalizing convolution operator. *Commun. Nonlinear Sci. Numer. Simul.* **2020**, *89*, 105338. [\[CrossRef\]](#)
14. Mishra, A. (Birla Institute of Technology & Science, Pilani, Jhunjhunu, Rajasthan, India). Private communication, 2022.
15. Gorenflo, R.; Kilbas, A.A.; Mainardi, F.; Rogosin, S.V. *Mittag-Leffler Functions, Related Topics and Applications*; Springer: New York, NY, USA, 2014; 443p, ISBN 978-3-662-43929-6
16. Ditkin, A.P.; Prudnikov, V.A. *Integral Transforms and Operational Calculus*; Pergamon Press: Oxford, UK, 1965.
17. Dimovski, I. Operational calculus for a class of differential operators. *C. R. Acad. Bulg. Sci.* **1966**, *19*, 1111–1114.
18. Delerue, P. Sur le calcul symbolique à n variables et fonctions hyper-besséliennes (II). *Ann. Soc. Sci. Bruxelles* **1953**, *3*, 229–274.
19. Kiryakova, V. *Generalized Fractional Calculus and Applications*; Pitman Res. Notes in Math. Ser.; Longman: Harlow, UK, 1994; Volume 301.
20. Comtet, L. *Advanced Combinatorics: The Art of Finite and Infinite Expansions*; Reidel Publishing Company: Dordrecht, The Netherlands, 1974.
21. Riordan, J. *An Introduction to Combinatorial Analysis*; J Wiley & Sons: Chichester, UK, 1958.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.