

**A SUBCLASS OF ANALYTIC FUNCTIONS  
DEFINED BY USING CERTAIN  
OPERATORS OF  
FRACTIONAL CALCULUS**

**O. Altıntaş, Hüseyin Irmak and H.M. Srivastava**

**DMS-690-IR**

**November 1994**

**A SUBCLASS OF ANALYTIC FUNCTIONS DEFINED BY  
USING CERTAIN OPERATORS OF FRACTIONAL CALCULUS**

**Osman Altıntaş**

Department of Mathematics  
Faculty of Science  
Hacettepe University  
06532 Beytepe - Ankara  
Turkey

**Hüseyin Irmak**

Department of Science Education  
Faculty of Education  
Hacettepe University  
06532 Beytepe - Ankara  
Turkey

**and**

**H.M. Srivastava**

Department of Mathematics and Statistics  
University of Victoria  
Victoria, British Columbia V8W 3P4  
Canada

---

1991 *Mathematics Subject Classification*. Primary 26A33, 30C45; Secondary 26A24.

*Key words and phrases*. Analytic functions, fractional calculus, coefficient bounds, distortion theorems, close-to-convex functions, starlike functions.

## Abstract

Making use of certain operators of fractional calculus, we introduce a new class  $\mathbb{F}_\delta(n, \lambda, \alpha)$  of functions which are analytic in the open unit disk  $\mathcal{U}$  and obtain a necessary and sufficient condition for a function to be in the class  $\mathbb{F}_\delta(n, \lambda, \alpha)$ . We also determine the radii of close-to-convexity, starlikeness, and convexity. Finally, an application involving fractional calculus of functions in the class  $\mathbb{F}_\delta(n, \lambda, \alpha)$  is considered.

### 1. Introduction and Definitions

Let  $\mathbb{F}(n)$  denote the class of functions  $f(z)$  of the form:

$$f(z) = z - \sum_{k=n+1}^{\infty} a_k z^k \quad (a_k \geq 0; \quad n \in \mathbb{N} := \{1, 2, 3, \dots\}), \quad (1.1)$$

which are analytic in the *open* unit disk

$$\mathcal{U} = \{z : z \in \mathbb{C} \quad \text{and} \quad |z| < 1\}.$$

Let  $\mathbb{F}_\delta(n, \lambda, \alpha)$  be the subclass of  $\mathbb{F}(n)$  consisting of functions which also satisfy the inequality:

$$\Re \left\{ \Gamma(2 - \delta) z^{\delta-1} \left[ (1 - \lambda) D_z^\delta f(z) + \lambda z D_z^{1+\delta} f(z) \right] \right\} > \alpha \quad (\delta + \alpha < 1) \quad (1.2)$$

for some  $\delta$  ( $0 \leq \delta < 1$ ),  $\lambda$  ( $0 \leq \lambda \leq 1$ ), and  $\alpha$  ( $0 \leq \alpha < 1$ ), and for all  $z \in \mathcal{U}$ . Here, and *throughout this paper*,  $D_z^\delta$  denotes an operator of fractional calculus, which is defined as follows (*cf.*, *e.g.*, [3] and [4]):

**Definition 1.** The *fractional integral of order*  $\mu$  is defined by

$$D_z^{-\mu} f(z) = \frac{1}{\Gamma(\mu)} \int_0^z \frac{f(\zeta)}{(z - \zeta)^{1-\mu}} d\zeta \quad (\mu > 0), \quad (1.3)$$

where  $f(z)$  is an analytic function in a simply-connected region of the  $z$ -plane containing the origin, and the multiplicity of  $(z - \zeta)^{\mu-1}$  is removed by requiring  $\log(z - \zeta)$  to be real when  $z - \zeta > 0$ .

**Definition 2.** The *fractional derivative of order  $\mu$*  is defined by

$$D_z^\mu f(z) = \frac{1}{\Gamma(1-\mu)} \frac{d}{dz} \int_0^z \frac{f(\zeta)}{(z-\zeta)^\mu} d\zeta \quad (0 \leq \mu < 1), \quad (1.4)$$

where  $f(z)$  is constrained, and the multiplicity of  $(z-\zeta)^{-\mu}$  is removed, as in Definition 1.

**Definition 3.** Under the hypotheses of Definition 1, the *fractional derivative of order  $k + \mu$*  is defined by

$$D_z^{k+\mu} f(z) = \frac{d^k}{dz^k} D_z^\mu f(z) \quad (0 \leq \mu < 1; \quad k \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}). \quad (1.5)$$

The object of the present paper is to investigate various interesting properties of functions belonging to the class  $\mathbb{F}_\delta(n, \lambda, \alpha)$ . We remark in passing that

$$\mathbb{F}_0(1, \lambda, \alpha) = \mathbb{F}_\lambda(\alpha) \quad (0 \leq \lambda \leq 1; \quad 0 \leq \alpha < 1), \quad (1.6)$$

where the class  $\mathbb{F}_\lambda(\alpha)$  was studied recently by Bhoosnurmath and Swamy [1].

## 2. A Theorem on Coefficient Bounds

**Theorem 1.** A function  $f(z) \in \mathbb{F}(n)$  is in the class  $\mathbb{F}_\delta(n, \lambda, \alpha)$  if and only if

$$\sum_{k=n+1}^{\infty} \frac{[1 + \lambda(k-1-\delta)]\Gamma(k+1)}{\Gamma(k+1-\delta)} a_k \leq 1 - \lambda\delta - \alpha \quad (\delta + \alpha < 1). \quad (2.1)$$

*The result is sharp.*

**Proof.** Suppose that  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ . Then we find from Definitions 1 and 3, and the inequality (1.2), that

$$\Re \left\{ 1 - \lambda\delta - \sum_{k=n+1}^{\infty} \frac{[1 + \lambda(k-1-\delta)]\Gamma(k+1)}{\Gamma(k+1-\delta)} a_k z^{k-1} \right\} > \alpha \quad (z \in \mathcal{U}).$$

If we choose  $z$  to be real and let  $z \rightarrow 1-$ , we get

$$1 - \lambda\delta - \sum_{k=n+1}^{\infty} \frac{[1 + \lambda(k-1-\delta)]\Gamma(k+1)}{\Gamma(k+1-\delta)} a_k \geq \alpha$$

$$(\delta + \alpha < 1; \quad 0 \leq \alpha < 1; \quad 0 \leq \delta < 1),$$

which is equivalent to the assertion (2.1) of Theorem 1.

Conversely, let us suppose that the inequality (2.1) holds true. Then we have

$$\begin{aligned}
& |\Gamma(2 - \delta) z^{\delta-1} [(1 - \lambda) D_z^\delta f(z) + z \lambda D_z^{1+\delta} f(z)] - 1 + \gamma \delta| \\
&= \left| - \sum_{k=n+1}^{\infty} \frac{[1 + \lambda(k - 1 - \delta)] \Gamma(k + 1)}{\Gamma(k + 1 - \delta)} a_k z^{k-1} \right| \\
&\leq \sum_{k=n+1}^{\infty} \frac{[1 + \lambda(k - 1 - \delta)] \Gamma(k + 1)}{\Gamma(k + 1 - \delta)} a_k |z|^{k-1} \\
&\leq 1 - \lambda \delta - \alpha \quad (z \in \mathcal{U}; \quad \delta + \alpha < 1; \quad 0 \leq \alpha < 1; \quad 0 \leq \delta < 1),
\end{aligned}$$

which implies that  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ .

Finally, we note that the assertion (2.1) of Theorem 1 is sharp, the extremal function being

$$f(z) = z - \frac{(1 - \lambda \delta - \alpha) \Gamma(n + 2 - \delta)}{[1 + \lambda(n - \delta)] \Gamma(n + 2)} z^{n+1} \quad (n \in \mathbb{N}). \quad (2.2)$$

**Corollary 1.** *If  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ , then*

$$a_{n+1} \leq \frac{(1 - \lambda \delta - \alpha) \Gamma(n + 2 - \delta)}{[1 + \lambda(n - \delta)] \Gamma(n + 2)} \quad (n \in \mathbb{N}). \quad (2.3)$$

**Corollary 2.** *A function  $f(z) \in \mathbb{F}(n)$  is in the class  $\mathbb{F}_0(n, \lambda, \alpha)$  if and only if*

$$\sum_{k=n+1}^{\infty} [1 + \lambda(k - 1)] a_k \leq 1 - \alpha \quad (0 \leq \lambda \leq 1; \quad 0 \leq \alpha < 1). \quad (2.4)$$

**Corollary 3** (cf. Bhoosnurmah and Swamy [1, p. 90, Theorem 1]). *A function  $f(z) \in \mathbb{F}(1)$  is in the class  $\mathbb{F}_0(1, \lambda, \alpha)$  if and only if*

$$\sum_{k=2}^{\infty} [1 + \lambda(k - 1)] a_k \leq 1 - \alpha \quad (0 \leq \lambda \leq 1; \quad 0 \leq \alpha < 1). \quad (2.5)$$

**Corollary 4.** *If  $f(z) \in \mathbb{F}_0(n, 1, \alpha)$ , then  $\Re\{f(z)\} > \alpha$  for all  $z \in \mathcal{U}$ .*

**Proof.** Since  $f(z) \in \mathbb{F}_0(n, 1, \alpha)$ , we have (cf. Silverman [5])

$$\sum_{k=n+1}^{\infty} k a_k \leq 1 - \alpha \quad (0 \leq \alpha < 1). \quad (2.6)$$

The result now follows from Theorem 1.

**Corollary 5.** *If  $f(z) \in \mathbb{F}_0(n, 0, \alpha)$ , then*

$$\Re \left\{ \frac{f(z)}{z} \right\} > \frac{1}{n+1} \quad (n \in \mathbb{N}).$$

**Proof.** Since  $f(z) \in \mathbb{F}_0(n, 0, \alpha)$ , we have

$$(n+1) \sum_{k=n+1}^{\infty} a_k \leq \sum_{k=n+1}^{\infty} k a_k \leq 1 - \alpha \quad (0 \leq \alpha < 1; \quad n \in \mathbb{N}), \quad (2.7)$$

by applying the known inequality (2.6). Therefore, we obtain

$$\sum_{k=n+1}^{\infty} a_k \leq \frac{1 - \alpha}{n+1} \quad (n \in \mathbb{N}). \quad (2.8)$$

**Corollary 6** (cf. Bhoosnurmath and Swamy [1, p. 91, Corollary 1.2]). *If  $f(z) \in \mathbb{F}_0(1, 0, \alpha)$ , then*

$$\Re \left\{ \frac{f(z)}{z} \right\} > \frac{1}{2}$$

for all  $z \in \mathcal{U}$ .

**Theorem 2.** *Let the function  $f(z)$  defined by (1.1) and the function  $g(z)$  defined by*

$$g(z) = z - \sum_{k=n+1}^{\infty} b_k z^k \quad (b_k \geq 0; \quad n \in \mathbb{N}) \quad (2.9)$$

be in the same class  $\mathbb{F}_\delta(n, \lambda, \alpha)$ . Then the function  $h(z)$  defined by

$$h(z) = (1 - \beta) f(z) + \beta g(z) = z - \sum_{k=n+1}^{\infty} c_k z^k$$

$$(c_k := (1 - \beta) a_k + \beta b_k \geq 0; \quad 0 \leq \beta \leq 1; \quad n \in \mathbb{N})$$

is also in the class  $\mathbb{F}_\delta(n, \lambda, \alpha)$ .

**Proof.** By the hypotheses of Theorem 2, we find from (2.1) that

$$\begin{aligned} & \sum_{k=n+1}^{\infty} \frac{[1 + \lambda(k-1-\delta)] \Gamma(k+1)}{\Gamma(k+1-\delta)} c_k \\ &= (1 - \beta) \sum_{k=n+1}^{\infty} \frac{[1 + \lambda(k-1-\delta)] \Gamma(k+1)}{\Gamma(k+1-\delta)} a_k \\ & \quad + \beta \sum_{k=n+1}^{\infty} \frac{[1 + \lambda(k-1-\delta)] \Gamma(k+1)}{\Gamma(k+1-\delta)} b_k \\ & \leq (1 - \beta)(1 - \lambda\delta - \alpha) + \beta(1 - \lambda\delta - \alpha) = 1 - \delta\lambda - \alpha, \end{aligned}$$

which completes the proof of Theorem 2.

### 3. Distortion Theorems Involving Operators of Fractional Calculus

**Theorem 3.** *If  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ , then*

$$|D_z^{-\mu} f(z)| \leq \frac{|z|^{1+\mu}}{\Gamma(2+\mu)} \left( 1 + \frac{(1-\lambda\delta-\alpha)\Gamma(2+\mu)\Gamma(n+2-\delta)}{[1+\lambda(n-\delta)]\Gamma(n+2+\mu)} |z| \right) \quad (3.1)$$

and

$$|D_z^{-\mu} f(z)| \geq \frac{|z|^{1+\mu}}{\Gamma(2+\mu)} \left( 1 - \frac{(1-\lambda\delta-\alpha)\Gamma(2+\mu)\Gamma(n+2-\delta)}{[1+\lambda(n-\delta)]\Gamma(n+2+\mu)} |z| \right) \quad (3.2)$$

for  $\mu > 0$  and  $n \in \mathbb{N}$ , and for all  $z \in \mathcal{U}$ .

**Proof.** Suppose that  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ . Then we find from (2.1) that

$$\begin{aligned} & \frac{[1+\lambda(n-\delta)]\Gamma(n+2)}{\Gamma(n+2-\delta)} \sum_{k=n+1}^{\infty} a_k \\ & \leq \sum_{k=n+1}^{\infty} \frac{[1+\lambda(k-1-\delta)]\Gamma(k+1)}{\Gamma(k+1-\delta)} a_k, \end{aligned} \quad (3.3)$$

which evidently yields

$$\sum_{k=n+1}^{\infty} a_k \leq \frac{(1-\lambda\delta-\alpha)\Gamma(n+2-\delta)}{[1+\lambda(n-\delta)]\Gamma(n+2)} \quad (n \in \mathbb{N}). \quad (3.4)$$

Making use of (3.4) and Definition 1, we have

$$\begin{aligned} D_z^{-\mu} f(z) &= \frac{z^{1+\mu}}{\Gamma(2+\mu)} \left( 1 - \sum_{k=n+1}^{\infty} \frac{\Gamma(k+1)\Gamma(2+\mu)}{\Gamma(k+1+\mu)} a_k z^{k-1} \right) \\ &= \frac{z^{1+\mu}}{\Gamma(2+\mu)} \left( 1 - \sum_{k=n+1}^{\infty} \Theta(k) a_k z^{k-1} \right), \end{aligned} \quad (3.5)$$

where, for convenience,

$$\Theta(k) = \frac{\Gamma(k+1)\Gamma(2+\mu)}{\Gamma(k+1+\mu)} \quad (\mu > 0; \quad k \geq n+1; \quad n \in \mathbb{N}).$$

Clearly, the function  $\Theta(k)$  is decreasing in  $k$ , and we have

$$0 < \Theta(k) \leq \Theta(n+1) = \frac{\Gamma(n+2)\Gamma(2+\mu)}{\Gamma(n+2+\mu)}. \quad (3.6)$$

Thus we find from (3.4), (3.5), and (3.6) that

$$\begin{aligned} |D_z^{-\mu} f(z)| &\leq \frac{|z|^{1+\mu}}{\Gamma(2+\mu)} \left( 1 + |z| \Theta(n+1) \sum_{k=n+1}^{\infty} a_k \right) \\ &\leq \frac{|z|^{1+\mu}}{\Gamma(2+\mu)} \left( 1 + \frac{(1-\lambda\delta-\alpha)\Gamma(2+\mu)\Gamma(n+2-\delta)}{[1+\lambda(n-\delta)]\Gamma(n+2+\mu)} |z| \right), \end{aligned}$$

which is precisely the assertion (3.1), and that

$$\begin{aligned} |D_z^{-\mu} f(z)| &\geq \frac{|z|^{1+\mu}}{\Gamma(2+\mu)} \left( 1 - |z| \Theta(n+1) \sum_{k=n+1}^{\infty} a_k \right) \\ &\geq \frac{|z|^{1+\mu}}{\Gamma(2+\mu)} \left( 1 - \frac{(1-\lambda\delta-\alpha)\Gamma(2+\mu)\Gamma(n+2-\delta)}{[1+\lambda(n-\delta)]\Gamma(n+2+\mu)} |z| \right), \end{aligned}$$

which is the same as the assertion (3.2).

**Theorem 4.** *If  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ , then*

$$|D_z^\mu f(z)| \leq \frac{|z|^{1-\mu}}{\Gamma(2-\mu)} \left( 1 + \frac{(1-\lambda\delta-\alpha)\Gamma(2-\mu)\Gamma(n+2-\delta)}{[1+\lambda(n-\delta)]\Gamma(n+2-\mu)} |z| \right) \quad (3.7)$$

and

$$|D_z^\mu f(z)| \geq \frac{|z|^{1-\mu}}{\Gamma(2-\mu)} \left( 1 - \frac{(1-\lambda\delta-\alpha)\Gamma(2-\mu)\Gamma(n+2-\delta)}{[1+\lambda(n-\delta)]\Gamma(n+2-\mu)} |z| \right) \quad (3.8)$$

for  $0 \leq \mu < 1$  and  $n \in \mathbb{N}$ , and for all  $z \in \mathcal{U}$ .

**Proof.** Suppose that  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ . Then we find from (2.1) that

$$\begin{aligned} &\frac{[1+\lambda(n-\delta)]\Gamma(n+1)}{\Gamma(n+2-\delta)} \sum_{k=n+1}^{\infty} k a_k \\ &\leq \sum_{k=n+1}^{\infty} \frac{[1+\lambda(k-1-\delta)]\Gamma(k+1)}{\Gamma(k+1-\delta)} a_k, \end{aligned} \quad (3.9)$$

which evidently yields

$$\sum_{k=n+1}^{\infty} k a_k \leq \frac{(1-\lambda\delta-\alpha)\Gamma(n+2-\delta)}{[1+\lambda(n-\delta)]\Gamma(n+1)} \quad (3.10)$$

$$(0 \leq \lambda \leq 1; \quad 0 \leq \delta < 1; \quad n \in \mathbb{N}).$$

Now, making use of (3.10) and Definition 2, we have

$$\begin{aligned} D_z^\mu f(z) &= \frac{z^{1-\mu}}{\Gamma(2-\mu)} \left( 1 - \sum_{k=n+1}^{\infty} \frac{\Gamma(k+1)\Gamma(2-\mu)}{\Gamma(k+1-\mu)} a_k z^{k-1} \right) \\ &= \frac{z^{1-\mu}}{\Gamma(2-\mu)} \left( 1 - \sum_{k=n+1}^{\infty} \Phi(k) k a_k z^{k-1} \right), \end{aligned} \quad (3.11)$$

where, for convenience,

$$\Phi(k) = \frac{\Gamma(k)\Gamma(2-\mu)}{\Gamma(k+1-\mu)} \quad (0 \leq \mu < 1; \quad k \geq n+1; \quad n \in \mathbb{N}).$$

Since the function  $\Phi(k)$  is decreasing in  $k$ , we also have

$$0 < \Phi(k) \leq \Phi(n+1) = \frac{\Gamma(n+1)\Gamma(2-\mu)}{\Gamma(n+2-\mu)}. \quad (3.12)$$

Thus we find from (3.10), (3.11), and (3.12) that

$$\begin{aligned} |D_z^\mu f(z)| &\leq \frac{|z|^{1-\mu}}{\Gamma(2-\mu)} \left( 1 + |z| \Phi(n+1) \sum_{k=n+1}^{\infty} k a_k \right) \\ &\leq \frac{|z|^{1-\mu}}{\Gamma(2-\mu)} \left( 1 + \frac{(1-\lambda\delta-\alpha)\Gamma(2-\mu)\Gamma(n+2-\delta)}{[1+\lambda(n-\delta)]\Gamma(n+2-\mu)} |z| \right), \end{aligned}$$

which is precisely the assertion (3.7), and that

$$\begin{aligned} |D_z^\mu f(z)| &\geq \frac{|z|^{1-\mu}}{\Gamma(2-\mu)} \left( 1 - |z| \Phi(n+1) \sum_{k=n+1}^{\infty} k a_k \right) \\ &\geq \frac{|z|^{1-\mu}}{\Gamma(2-\mu)} \left( 1 - \frac{(1-\lambda\delta-\alpha)\Gamma(2-\mu)\Gamma(n+2-\delta)}{[1+\lambda(n-\delta)]\Gamma(n+1-\mu)} |z| \right), \end{aligned}$$

which is the same as the assertion (3.8).

**Theorem 5.** *If  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ , then*

$$|D_z^{1+\delta} f(z)| \leq \frac{|z|^{-\delta}}{\Gamma(1-\delta)} \left( 1 + \frac{(1-\lambda\delta-\alpha)(n+1-\delta)\Gamma(1-\delta)}{1+\lambda(n-\delta)} |z| \right) \quad (3.13)$$

and

$$|D_z^{1+\delta} f(z)| \geq \frac{|z|^{-\delta}}{\Gamma(1-\delta)} \left( 1 - \frac{(1-\lambda\delta-\alpha)(n+1-\delta)\Gamma(2-\delta)}{1+\lambda(n-\delta)} |z| \right) \quad (3.14)$$

for  $0 \leq \delta < 1$  and  $n \in \mathbb{N}$ , and for all  $z \in \mathcal{U}$ .

**Proof.** Suppose that  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ . Then we find from (2.1) that

$$\sum_{k=n+1}^{\infty} \frac{\Gamma(k+1)\Gamma(1-\delta)}{\Gamma(k-\delta)} a_k \leq \frac{(1-\lambda\delta-\alpha)(n+1-\delta)\Gamma(1-\delta)}{1+\lambda(n-\delta)} \quad (3.15)$$

$$(0 \leq \lambda \leq 1; \quad 0 \leq \delta < 1; \quad n \in \mathbb{N}).$$

On the other hand, by applying Definition 3 (with  $k = 1$  and  $\mu = \delta$ ) we obtain

$$D_z^{1+\delta} f(z) = \frac{z^{-\delta}}{\Gamma(1-\delta)} \left( 1 - \sum_{k=n+1}^{\infty} \frac{\Gamma(k+1)\Gamma(1-\delta)}{\Gamma(k-\delta)} a_k z^{k-1} \right). \quad (3.16)$$

Thus, by combining (3.15) and (3.16), we immediately get the assertions (3.13) and (3.14) of Theorem 5.

Setting  $\delta = \mu = 0$  in Theorem 4, we have

**Corollary 7.** *If  $f(z) \in \mathbb{F}_0(n, \lambda, \alpha)$ , then*

$$\left| |z| - \frac{1-\alpha}{1+\lambda n} |z|^2 \right| \leq |f(z)| \leq |z| + \frac{1-\alpha}{1+\lambda n} |z|^2 \quad (3.17)$$

for all  $z \in \mathcal{U}$  and  $n \in \mathbb{N}$ .

For  $\delta = 0$ , Theorem 5 yields

**Corollary 8.** *If  $f(z) \in \mathbb{F}_0(n, \lambda, \alpha)$ , then*

$$1 - \frac{(1-\alpha)(n+1)}{1+\lambda n} |z| \leq |f'(z)| \leq 1 + \frac{(1-\alpha)(n+1)}{1+\lambda n} |z| \quad (3.18)$$

for all  $z \in \mathcal{U}$  and  $n \in \mathbb{N}$ .

Next, setting  $\delta = \mu = 0$  and  $n = 1$  in Theorem 4 (or, simply,  $n = 1$  in Corollary 7), we have

**Corollary 9** (cf. Bhoosnurmath and Swamy [1, p. 91, Theorem 2]). *If  $f(z) \in \mathbb{F}_0(1, \lambda, \alpha)$ , then*

$$\left| |z| - \frac{1-\alpha}{1+\lambda} |z|^2 \right| \leq |f(z)| \leq |z| + \frac{1-\alpha}{1+\lambda} |z|^2 \quad (3.19)$$

for all  $z \in \mathcal{U}$ .

If we set  $\delta = 0$  and  $n = 1$  in Theorem 5 (or, alternatively, if we just let  $n = 1$  in Corollary 8), we obtain

**Corollary 10** (cf. Bhoosnurmath and Swamy [1, p. 92, Theorem 3]). *If  $f(z) \in \mathbb{F}_0(1, \lambda, \alpha)$ , then*

$$1 - \frac{2(1-\alpha)}{1+\lambda} |z| \leq |f'(z)| \leq 1 + \frac{2(1-\alpha)}{1+\lambda} |z| \quad (3.20)$$

for all  $z \in \mathcal{U}$ .

Numerous further consequences of Theorems 3, 4, and 5 (and of Corollaries 7, 8, 9, and 10) can indeed be deduced by specializing the various parameters involved.

#### 4. Radii of Close-to-Convexity, Starlikeness, and Convexity

A function  $f(z) \in \mathbb{F}(n)$  is said to be *close-to-convex of order  $\beta$*  if it satisfies the inequality (cf. [2] and [6]):

$$\Re \{f'(z)\} > \beta \quad (4.1)$$

for some  $\beta$  ( $0 \leq \beta < 1$ ) and for all  $z \in \mathcal{U}$ . On the other hand, a function  $f(z) \in \mathbb{F}(n)$  is said to be *starlike of order  $\beta$*  if it satisfies the inequality (cf. [2] and [6]):

$$\Re \left\{ \frac{zf'(z)}{f(z)} \right\} > \beta \quad (4.2)$$

for some  $\beta$  ( $0 \leq \beta < 1$ ) and for all  $z \in \mathcal{U}$ . Furthermore, a function  $f(z) \in \mathbb{F}(n)$  is said to be *convex of order  $\beta$*  if and only if  $zf'(z)$  is starlike of order  $\beta$ , that is, if it satisfies the inequality (cf. [2] and [6]):

$$\Re \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \beta \quad (4.3)$$

for some  $\beta$  ( $0 \leq \beta < 1$ ) and for all  $z \in \mathcal{U}$ .

**Theorem 6.** *If  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ , then  $f(z)$  is close-to-convex of order  $\beta$  in  $|z| < r_1(\alpha, \lambda, \delta, \beta)$ , where*

$$r_1(\alpha, \lambda, \delta, \beta) = \inf_k \left[ \frac{(1-\beta)\Gamma(k)[1+\lambda(k-1-\delta)]}{(1-\lambda\delta-\alpha)\Gamma(k+1-\delta)} \right]^{1/(k-1)}$$

$$(k \geq n+1; \quad n \in \mathbb{N}).$$

**Proof.** It is sufficient to show that  $|f'(z) - 1| < 1 - \beta$ . Indeed we have

$$|f'(z) - 1| \leq \sum_{k=n+1}^{\infty} k a_k |z|^{k-1} \leq 1 - \beta \quad (4.4)$$

and

$$\sum_{k=n+1}^{\infty} \frac{\Gamma(k+1)[1 + \lambda(k - \delta - 1)]}{\Gamma(k+1 - \delta)} a_k \leq 1 - \lambda\delta - \alpha. \quad (4.5)$$

Hence (4.4) is true if

$$\frac{k|z|^{k-1}}{1 - \beta} \leq \frac{\Gamma(k+1)[1 + \lambda(k - \delta - 1)]}{(1 - \lambda\delta - \alpha)\Gamma(k+1 - \delta)} \quad (k \geq n+1; \quad n \in \mathbb{N}). \quad (4.6)$$

Solving (4.6) for  $|z|$ , we obtain

$$|z| \leq \left[ \frac{(1 - \beta)\Gamma(k)[1 + \lambda(k - 1 - \delta)]}{(1 - \lambda\delta - \alpha)\Gamma(k+1 - \delta)} \right]^{1/(k-1)} \quad (k \geq n+1; \quad n \in \mathbb{N}),$$

which obviously proves Theorem 6.

**Theorem 7.** *If  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ , then  $f(z)$  is starlike of order  $\beta$  in*

$$|z| < r_2(\alpha, \lambda, \delta, \beta),$$

where

$$r_2(\alpha, \lambda, \delta, \beta) = \inf_k \left[ \frac{(1 - \beta)\Gamma(k+1)[1 + \lambda(k - 1 - \delta)]}{(k - \beta)(1 - \lambda\delta - \alpha)\Gamma(k+1 - \delta)} \right]^{1/(k-1)} \\ (k \geq n+1; \quad n \in \mathbb{N}).$$

**Proof.** We must show that

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| \leq 1 - \beta \quad \text{for } |z| < r_2(\alpha, \lambda, \delta, \beta).$$

In fact, we have

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \frac{\sum_{k=n+1}^{\infty} (k-1) a_k |z|^{k-1}}{1 - \sum_{k=n+1}^{\infty} a_k |z|^{k-1}} \leq 1 - \beta, \quad (4.7)$$

if

$$\frac{(k - \beta)|z|^{k-1}}{1 - \beta} \leq \frac{\Gamma(k+1)[1 + \lambda(k - \delta - 1)]}{(1 - \lambda\delta - \alpha)\Gamma(k+1 - \delta)} \quad (k \geq n+1; \quad n \in \mathbb{N}), \quad (4.8)$$

which evidently proves Theorem 7.

**Corollary 11.** *If  $f(z) \in \mathbb{F}_\delta(n, \lambda, \alpha)$ , then  $f(z)$  is convex of order  $\beta$  in*

$$|z| < r_3(\alpha, \lambda, \delta, \beta),$$

where

$$r_3(\alpha, \lambda, \delta, \beta) = \inf_k \left[ \frac{(1 - \beta)\Gamma(k)[1 + \lambda(k - 1 - \delta)]}{(k - \beta)(1 - \lambda\delta - \alpha)\Gamma(k + 1 - \delta)} \right]^{1/(k-1)}$$

$$(k \geq n + 1; \quad n \in \mathbb{N}).$$

**Corollary 12.** *If  $f(z) \in \mathbb{F}_0(1, \lambda, \alpha)$ , then  $f(z)$  is close-to-convex of order  $\beta$  in  $|z| < r_4(\alpha, \lambda, \beta)$ , where*

$$r_4(\alpha, \lambda, \beta) = \inf_k \left[ \frac{(1 - \beta)[1 + \lambda(k - 1)]}{k(1 - \alpha)} \right]^{1/(k-1)} \quad (k \in \mathbb{N} \setminus \{1\}).$$

**Corollary 13.** *If  $f(z) \in \mathbb{F}_0(1, \lambda, \alpha)$ , then  $f(z)$  is starlike of order  $\beta$  in  $|z| < r_5(\alpha, \lambda, \beta)$ , where*

$$r_5(\alpha, \lambda, \beta) = \inf_k \left[ \frac{(1 - \beta)[1 + \lambda(k - 1)]}{(k - \beta)(1 - \alpha)} \right]^{1/(k-1)} \quad (k \in \mathbb{N} \setminus \{1\}).$$

**Corollary 14.** *If  $f(z) \in \mathbb{F}_0(1, \lambda, \alpha)$ , then  $f(z)$  is convex of order  $\beta$  in  $|z| < r_6(\alpha, \lambda, \beta)$ , where*

$$r_6(\alpha, \lambda, \beta) = \inf_k \left[ \frac{(1 - \beta)[1 + \lambda(k - 1)]}{k(k - \beta)(1 - \alpha)} \right]^{1/(k-1)} \quad (k \in \mathbb{N} \setminus \{1\}).$$

In their *special* cases when  $\beta = 0$ , Corollaries 12, 13, and 14 were proved earlier by Bhoosnurmath and Swamy [1, pp. 93-94, Theorems 5 and 6].

### Acknowledgments

The present investigation was supported, in part, by the Natural Sciences and Engineering Research Council of Canada under Grant OGP0007353.

## References

- [1] S.S. Bhoosnurmath and S.R. Swamy, Certain classes of analytic functions with negative coefficients, *Indian J. Math.* **27**(1985), 89-98.
- [2] P.L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften **259**, Springer-Verlag, New York, Berlin, Heidelberg, and Tokyo, 1983.
- [3] S. Owa, On the distortion theorems. I, *Kyungpook Math. J.* **18**(1978), 53-59.
- [4] S. Owa and H.M. Srivastava, Univalent and starlike generalized hypergeometric functions, *Canad. J. Math.* **39**(1987), 1057-1077.
- [5] H. Silverman, Univalent functions with negative coefficients, *Proc. Amer. Math. Soc.* **51**(1975), 109-116.
- [6] H.M. Srivastava and S. Owa (Editors), *Current Topics in Analytic Function Theory*, World Scientific Publishing Company, Singapore, New Jersey, London, and Hong Kong, 1992.