

**CERTAIN RESULTS ASSOCIATED WITH THE
GENERALIZED RIEMANN ZETA FUNCTIONS**

R.K. RAINA & H.M. SRIVASTAVA

DMS-675-IR

August 1994

CERTAIN RESULTS ASSOCIATED WITH THE GENERALIZED RIEMANN ZETA FUNCTIONS

R.K. Raina and H.M. Srivastava

Abstract

In this paper we derive new (linear and bilateral) generating functions involving the Riemann zeta function (or the generalized zeta function). A connection with the Lambert transform is given, and a generalized Lambert transform is introduced. An inversion formula of this transform is obtained and its relationship with the generalized zeta function is also depicted.

1. Introduction

The generalized (Hurwitz's) zeta function is defined by [1, p. 24]:

$$\zeta(s, a) = \sum_{n=0}^{\infty} (a+n)^{-s} \quad (1.1)$$

$$(\Re(s) > 1; \quad a \neq 0, -1, -2, \dots),$$

so that, evidently,

$$\zeta(s, 1) = \sum_{n=1}^{\infty} n^{-s} = \zeta(s), \quad (1.2)$$

where $\zeta(s)$ is the Riemann zeta function. The function $\Phi(z, s, a)$ extends (1.1) further, and is defined by [1, p. 27, Equation (1)]

$$\Phi(z, s, a) = \sum_{n=0}^{\infty} (a+n)^{-s} z^n \quad (\Re(a) > 0; \quad |z| < 1). \quad (1.3)$$

1991 *Mathematics Subject Classification*. Primary 11M06, 11M35; Secondary 33C20.

Key words and phrases. Riemann zeta function, generating function, Hurwitz zeta function, Lambert transform, Laguerre polynomials, Gauss hypergeometric function, Mellin transform, generalized hypergeometric function, Mellin inversion theorem.

Equivalently, the function $\Phi(z, s, a)$ has the integral representation:

$$\Phi(z, s, a) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{-at} (1 - ze^{-t})^{-1} dt, \quad (1.4)$$

provided that $\Re(a) > 0$ (and either $|z| \leq 1$, $z \neq 1$, and $\Re(s) > 0$, or $z = 1$ and $\Re(s) > 1$).

Our object in the present paper is to obtain certain types of linear as well as bilateral generating functions involving the function $\Phi(z, s, a)$ defined by (1.3). A generalized Lambert transform is introduced and its inversion is obtained, and further expressions in terms of the function $\Phi(z, s, a)$ are derived from the Lambert transform of a general system of polynomials.

2. Generating Functions

From the definition (1.3), and the binomial expansion

$$\sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} z^n = (1 - z)^{-\lambda} \quad (|z| < 1), \quad (2.1)$$

we readily have the generating function:

$$\sum_{n=0}^{\infty} (\lambda)_n \Phi(x, \lambda + n, a) \frac{t^n}{n!} = \Phi(x, \lambda, a - t) \quad (|t| < |a|; \lambda \neq 1), \quad (2.2)$$

where, and in what follows, $(\lambda)_n = \Gamma(\lambda + n)/\Gamma(\lambda)$.

The generating function (2.2) can be extended further. Indeed, in terms of Gauss's hypergeometric function [1, p. 56], it follows from (1.3) that

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{(\lambda)_n (\mu)_n}{(\nu)_n} \Phi(x, \lambda + \mu - \nu + n, a) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} x^n (n + a)^{\nu - \lambda - \mu} {}_2F_1 \left[\begin{matrix} \lambda, \mu; \\ \nu; \end{matrix} \frac{t}{n + a} \right], \end{aligned} \quad (2.3)$$

provided that $|t/a| < 1$ and $\Re(\lambda + \mu) > \Re(\nu) > 0$. More generally, (2.3) may be extended in the form:

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{\prod_{j=1}^p (a_j)_n}{\prod_{j=1}^q (b_j)_n} \Phi(x, \omega + n, a) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} x^n (n + a)^{-\omega} {}_pF_q \left[\begin{matrix} (a_p); \\ (b_q); \end{matrix} \frac{t}{n + a} \right], \end{aligned} \quad (2.4)$$

provided that $p \leq q$, $|t| < |a|$, and $\Re(\omega) > 0$. Here (a_p) abbreviates the array of p parameters

$$a_1, \dots, a_p,$$

with similar interpretations for (b_q) *et cetera*, ${}_pF_q$ being a generalized hypergeometric function with p numerator and q denominator parameters [1, Chapter 4].

Next we consider a polynomial set $\{S_n^{(m)}(z)\}_{n=0}^\infty$ defined by (*cf.* [3, p. 1, Eq. (1)])

$$S_n^{(m)}(z) = \sum_{j=0}^{\lfloor n/m \rfloor} \frac{(-n)_{mj}}{j!} C_j z^j \quad (m \in \mathbb{N}; n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}), \quad (2.5)$$

where, and in what follows, $\{C_n\}_{n=0}^\infty$ is a suitably bounded sequence of complex numbers. Then, by simple series rearrangement technique and the application of (2.2), we arrive at the following bilateral generating function:

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} \Phi(x, \lambda + n, a) S_n^{(m)}(y) t^n \\ = \sum_{n=0}^{\infty} \frac{(\lambda)_{mn}}{n!} C_n \Phi(x, \lambda + mn, a - t) \{y(-t)^m\}^n, \end{aligned} \quad (2.6)$$

provided that $|x| \leq 1$, $|t| < |a|$, and $\lambda \neq 1$.

Example. By specializing the sequence $\{C_n\}$ as follows:

$$C_n = \frac{\prod_{j=1}^p (a_j)_n}{m^{mn} \prod_{j=1}^q (b_j)_n} \quad (m \in \mathbb{N}; n \in \mathbb{N}_0), \quad (2.7)$$

we find from (2.6) that

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} \Phi(x, \lambda + n, a) {}_{m+p}F_q \left[\begin{matrix} \Delta(m; -n), (a_p); \\ (b_q); \end{matrix} y \right] t^n \\ = \sum_{n=0}^{\infty} \frac{(\lambda)_{mn}}{n!} \frac{\prod_{j=1}^p (a_j)_n}{\prod_{j=1}^q (b_j)_n} \Phi(x, \lambda + mn, a - t) \left\{ y \left(-\frac{t}{m} \right)^m \right\}^n, \end{aligned} \quad (2.8)$$

where $\Delta(m; \lambda)$ denotes the array of m parameters

$$\frac{\lambda}{m}, \frac{\lambda+1}{m}, \dots, \frac{\lambda+m-1}{m} \quad (m \in \mathbb{N}).$$

In particular, in terms of the classical Laguerre polynomials $L_n^{(\alpha)}(z)$, for $p = 0$, $q = m = 1$, and $b_1 = 1 + \alpha$, (2.8) yields

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{(\lambda)_n}{(1+\alpha)_n} \Phi(x, \lambda+n, a) L_n^{(\alpha)}(y) t^n \\ &= \sum_{n=0}^{\infty} \frac{(\lambda)_n}{(1+\alpha)_n} \Phi(x, \lambda+n, a-t) \frac{(-yt)^n}{n!}. \end{aligned} \quad (2.9)$$

Our formula (2.2) includes some recent results due to Srivastava ([4, p. 48, Equation (2.2)]; see also [5]) and Miller [2, p. 99, Equation (3.20)].

3. The Lambert Transform

Let $f(t)$ ($t \geq 0$) be a continuous function; then the Lambert transform of $f(t)$ is defined by

$$\bar{f}(s) = LM\{f(t)\} = \int_0^{\infty} \frac{st}{e^{st}-1} f(t) dt, \quad (3.1)$$

and it converges for all s such that $\Re(s) > 0$. It follows easily from (3.1) and (1.1) that

$$LM\{t^{\alpha-1} e^{-\nu st}\} = \frac{\Gamma(\alpha+1)}{s^\alpha} \zeta(\alpha+1, \nu+1) \quad (3.2)$$

$$(\Re(\alpha) > 0; \quad \nu \neq -1, -2, -3, \dots; \quad s \neq 0).$$

Also, in view of (3.1) and (2.5), we have

$$\begin{aligned} & LM\left\{t^{\alpha-1} e^{-\nu st} S_m^{(q)}(xt)\right\} \\ &= \sum_{j=0}^{[m/q]} \frac{(-m)_{qj}}{j!} \frac{\Gamma(\alpha+j+1)}{s^{\alpha+j}} \zeta(\alpha+j+1, \nu+1) C_j x^j. \end{aligned} \quad (3.3)$$

A special case of (3.3) of interest is the one which occurs when

$$C_j = \frac{1}{(\alpha+1)_j} \quad (j \in \mathbb{N}_0)$$

and $q = 1$. Thus, in terms of the Laguerre polynomials, (3.3) gives

$$\begin{aligned} LM \left\{ t^{\alpha-1} e^{-\nu st} L_m^{(\alpha)}(xt) \right\} \\ = \frac{\Gamma(m + \alpha + 1)}{m! s^\alpha} \sum_{j=0}^m \frac{(-m)_j}{j!} \zeta(\alpha + j + 1, \nu + 1) \left(\frac{x}{s}\right)^j, \end{aligned} \quad (3.4)$$

where $s \neq 0$, $m \in \mathbb{N}_0$, $\Re(\alpha) > 0$, and $|x/s| < 1$.

4. The Generalized Lambert Transform

We consider the following generalization of the Lambert transform (3.1):

$$F(s) = GLM\{f(t)\} = \int_0^\infty \frac{st}{e^{st} - x} f(t) dt, \quad (4.1)$$

provided that $\Re(s) > 0$, $|x| \leq 1$, $f(t) \in \Omega$, and $\Re(\gamma) > -2$, where Ω denotes the class of functions $f(t)$ which are continuous for $t > 0$ and satisfy the order estimates:

$$f(t) = \begin{cases} O(t^\gamma) & (t \rightarrow 0+) \\ O(t^\delta) & (t \rightarrow \infty). \end{cases} \quad (4.2)$$

Obviously, for $x = 1$, (4.1) reduces to the Lambert transform (3.1). If $f(t) = t^{\alpha-1} e^{-\nu st}$, then we find from (1.4) and (4.1) that

$$GLM \left\{ t^{\alpha-1} e^{-\nu st} \right\} = \frac{\Gamma(\alpha + 1)}{s^\alpha} \Phi(x, \alpha + 1, \nu + 1). \quad (4.3)$$

Inversion of the Generalized Lambert Transform (4.1). On applying the Mellin transform [6, p. 46], (4.1) yields

$$\begin{aligned} \phi(k) &= \int_0^\infty s^{-k-1} F(s) ds \\ &= \int_0^\infty s^{-k-1} \left(\int_0^\infty \frac{st}{e^{st} - x} f(t) dt \right) ds \\ &= \int_0^\infty f(t) \left(\int_0^\infty \frac{st}{e^{st} - x} \cdot s^{-k-1} ds \right) dt \\ &= \int_0^\infty f(t) \cdot \frac{\Gamma(1-k)}{t^{-k}} \Phi(x, 1-k, 1) dt, \end{aligned} \quad (4.4)$$

provided that $f \in \Omega$ and $|x| \leq 1$. Now, by the Mellin inversion theorem, we get the following inversion formula for the generalized Lambert transform (4.1):

$$\begin{aligned} & \frac{1}{2} \{f(t+0) + f(t-0)\} \\ &= \frac{1}{2\pi i} \int_{\tau-i\infty}^{\tau+i\infty} \{\Gamma(1-\zeta) \Phi(x, 1-\zeta, 1)\}^{-1} t^{-\zeta} \phi(\zeta) d\zeta \quad (\tau > 0), \end{aligned} \tag{4.5}$$

provided that $t^{\zeta-1} f(t) \in L(0, \infty)$ and $f(t)$ is of bounded variation in the neighbourhood of the point t , $\phi(\zeta)$ being given by (4.4).

Acknowledgements

The present investigation was supported, in part, by the Natural Sciences and Engineering Research Council of Canada under Grant OGP0007353. The first-named author wishes to thank the University Grants Commission of India for providing financial support for this project.

References

1. A. Erdélyi, W. Magnus, F. Oberhettinger, and F.G. Tricomi, *Higher Transcendental Functions*, Vol. 1, McGraw-Hill, New York, Toronto, and London, 1953.
2. E.L. Miller, Summability of power series using the Lambert transform, *Univ. Nac. Tucumán Rev. Ser. A* **28**(1978), 89-106.
3. H.M. Srivastava, A contour integral involving Fox's H -function, *Indian J. Math.* **14**(1972), 1-6.
4. H.M. Srivastava, Some infinite series associated with the Riemann zeta function, *Yokohama Math. J.* **35**(1987), 47-50.
5. H.M. Srivastava, Sums of certain series of the Riemann zeta function, *J. Math. Anal. Appl.* **134**(1988), 129-140.

6. E.C. Titchmarch, *Introduction to the Theory of Fourier Integrals*, Clarendon Press, Oxford, 1948.

R.K. Raina:

Department of Mathematics
College of Technology and Agricultural
Engineering
(Rajasthan Agricultural University)
Udaipur 313001, Rajasthan
India

H.M. Srivastava:

Department of Mathematics and Statistics
University of Victoria
Victoria, British Columbia V8W 3P4
Canada

