
Faculty of Science

Faculty Publications

Prioritizing conservation actions for Pacific salmon in Canada

Jessica C. Walsh, Katrina Connors, Eric Hertz, Laura Kehoe, Tara G. Martin, Brendan Connors, Michael J. Bradford, Cameron Freshwater, Alejandro Frid, Jessica Halverson, Jonathan W. Moore, Michael H. H. Price, & John D. Reynolds

April 2020

© 2020 Jessica C. Walsh et al. This is an open access article distributed under the terms of the Creative Commons Attribution License. <https://creativecommons.org/licenses/by-nc/4.0/>

This article was originally published at:
<https://doi.org/10.1111/1365-2664.13646>

Citation for this paper:

Walsh, J. C., Connors, K., Hertz, E., Kehoe, L., Martin, T. G., Connors, B., Bradford, M. J., Freshwater, C., Frid, A., Halverson, J., Moore, J. W., Price, M. H. H., & Reynolds, J. D. (2020). Prioritizing conservation actions for Pacific salmon in Canada. *Journal of Applied Ecology*, 2020, 1-12. <https://doi.org/10.1111/1365-2664.13646>.



Prioritizing conservation actions for Pacific salmon in Canada

Jessica C. Walsh¹ | Katrina Connors^{2,3} | Eric Hertz² | Laura Kehoe^{4,5,6} |
Tara G. Martin⁶ | Brendan Connors⁷ | Michael J. Bradford⁸ | Cameron Freshwater⁹ |
Alejandro Frid^{10,11} | Jessica Halverson¹ | Jonathan W. Moore¹ | Michael H. H. Price¹ |
John D. Reynolds¹

¹Department of Biological Sciences, Earth to Ocean Research Group, Simon Fraser University, Burnaby, BC, Canada; ²Pacific Salmon Foundation – Salmon Watersheds Program, Vancouver, BC, Canada; ³School of Environment and Sustainability, Royal Roads University, Victoria, BC, Canada; ⁴Department of Biology, University of Victoria, Victoria, BC, Canada; ⁵The Nature Conservancy, London, UK; ⁶Department of Forest and Conservation Sciences, University of British Columbia, Vancouver, BC, Canada; ⁷Institute of Ocean Sciences, Fisheries and Oceans Canada, Sidney, BC, Canada; ⁸West Vancouver Laboratory, Fisheries and Oceans Canada, West Vancouver, BC, Canada; ⁹Pacific Biological Station, Fisheries and Oceans Canada, Nanaimo, BC, Canada; ¹⁰Central Coast Indigenous Resource Alliance, Campbell River, BC, Canada and ¹¹School of Environmental Studies, University of Victoria, Victoria, BC, Canada

Correspondence

Jessica C. Walsh
Email: jessica.walsh@monash.edu

Present address

Jessica C. Walsh, School of Biological Sciences, Monash University, Clayton, 3800, Vic., Australia and
School of Earth and Environmental Sciences, University of Queensland, St Lucia, 4072, QLD, Australia

Funding information

Central Coast Indigenous Resource Alliance; Pacific Salmon Foundation; Tom Buell Endowment Fund; Mitacs; Environment and Climate Change Canada; NSERC

Handling Editor: Vitor Paiva

Abstract

1. Current investment in conservation is insufficient to adequately protect and recover all ecosystems and species. The challenge of allocating limited funds is acute for Pacific salmon *Oncorhynchus* spp. in Canada, which lack a strategic approach to ensure that resources are spent on actions most likely to cost-effectively recover diminished populations.
2. We applied the Priority Threat Management framework to prioritize strategies most likely to maximize the number of thriving Pacific salmon populations on the Central Coast of British Columbia, Canada. These included 79 genetically, ecologically and spatially distinct population groups called conservation units (CUs) for five salmon species. This region has high salmon biodiversity and spans the territories of four First Nations: the Heiltsuk, Nuxalk, Kitasoo/Xai'xais and Wuikinuxv.
3. Using structured expert elicitation of Indigenous and other experts, we quantified the estimated benefits, costs and feasibility of implementing 10 strategies. Under a business-as-usual scenario (i.e. no additional investments in salmon conservation or management), experts predicted that only one in four CUs would have >50% chance of achieving a thriving status within 20 years. Limiting future industrial development in salmon habitats, which was predicted to safeguard CUs from future declines, was identified as the most cost-effective strategy. Investment in three strategies: (a) removal of artificial barriers to fish migration, (b) watershed protection and (c) stream restoration—at 11.3M CAD per year—was predicted to result in nearly half (34 of 79) of the CUs having a >60% chance of meeting the conservation objective.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2020 The Authors. *Journal of Applied Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society

4. If all conservation strategies were implemented, experts estimated a >50% probability of achieving a thriving status for 78 of 79 CUs, at an annual cost of 17.3M CAD. However, even with the implementation of all strategies, most sockeye salmon CUs were unlikely to achieve higher probability targets of reaching the objective.
5. *Policy implications.* We illustrate how Priority Threat Management can incorporate the perspectives and expertise of Indigenous peoples and other experts to prioritize conservation strategies based on their cost, benefit and feasibility. Implementation of this framework can help safeguard and recover Pacific salmon in Canada, and could also be used to prioritize actions for other conservation issues globally.

KEYWORDS

conservation planning, decision-support tool, Indigenous knowledge, Pacific salmon, Priority Threat Management, recovery planning, stream restoration, Wild Salmon Policy

1 | INTRODUCTION

There is an urgent need for strategic planning and prioritization of conservation actions to ensure the persistence and recovery of species and ecosystems, many of which are in decline globally (Martin et al., 2018). However, available resources are inadequate to manage all threats, there is uncertainty around how best to abate these threats, and decision-makers often have limited time and information for prioritizing recovery actions (Martin, Burgman, et al., 2012; Martin, Nally, et al., 2012). Pacific salmon (*Oncorhynchus* spp.) exemplify a group of species that could benefit from strategic planning and prioritization because of their important economic, ecological and cultural role throughout their range (Mantua et al., 2009). Many populations are diminished or have declined dramatically in recent decades (Governor's Salmon Recovery Office, 2018; Gustafson et al., 2007; Malick & Cox, 2016; Price, English, Rosenberger, MacDuffee, & Reynolds, 2017). This has led to curtailed fisheries (Ogden et al., 2014; Walters, English, Korman, & Hilborn, 2019), loss of livelihoods, erosion of the cultural identities of Indigenous communities (Garibaldi & Turner, 2004) and potential adverse impacts on coastal ecosystems (Levi et al., 2012). National policies, such as Canada's Wild Salmon Policy (WSP; Fisheries and Oceans Canada, 2005, 2018), recognize the need for a strategic prioritization of management actions (Nelitz, Murray, & Wieckowski, 2008), however, there currently is no strategic framework in Canada for determining how and where to invest limited resources across multiple threats to maximize the probability of recovering salmon populations.

Deciding where and how best to invest in recovery efforts for salmon is a challenge. Factors regulating populations often are complex and poorly understood, but can include stressors such as: overfishing, habitat loss and degradation, barriers to migration, disease, predation, poor survival and climate change (Hoekstra, Bartz, Ruckelshaus, Moslemi, & Harms, 2007; Ruckelshaus, Levin, Johnson, & Kareiva, 2002; Schoen et al., 2017). Furthermore, Pacific salmon have a high degree of local adaptation, and policies in the United States and Canada mandate conservation of these evolutionarily distinct units (Waples, 1991). However, existing budgets cannot

manage all threats and conserve all salmon populations (Gardner & Pinfold, 2011). There have been many efforts to prioritize potential recovery options for Pacific salmon, such as stream restoration activities (Beechie, Pess, Roni, & Giannico, 2008; Roni et al., 2002), and recovery plans for at-risk populations (Good, Beechie, McElhany, McClure, & Ruckelshaus, 2007; Kareiva, Marvier, & McClure, 2000). However, it remains difficult to evaluate the costs and benefits of actions across multiple co-occurring populations and species to ensure that returns on investment are maximized.

Two aspects of the current approach to salmon conservation and recovery planning are problematic. First, the cost-effectiveness of management actions across populations often is not quantified. For example, billions of dollars have been spent on conservation and restoration efforts for Pacific salmon over the last several decades, including enhancement via hatcheries and restoration of spawning and rearing habitats; yet, the effects on populations often are mixed, negligible or unclear (Barnas, Katz, Hamm, Dias, & Jordan, 2015; Bernhardt et al., 2005). Careful comparison of the cost-effectiveness across management strategies for multiple species and populations could help to maximize salmon recovery potential and protection. Second, resources typically are allocated to the most threatened populations or species (e.g. Cultus Sockeye Recovery Team, 2009; Sakinaw Lake Sockeye Recovery Team, 2005), or those of high public interest (e.g. Chinook salmon). Despite the legal and social reasons for prioritizing these diminished populations, they may have the lowest probabilities of recovery and may require the most expensive solutions compared to less threatened populations with higher recovery potential. If the goal is to increase the total number of healthy populations and their benefits to society, then focusing on the most threatened ones may be suboptimal.

Decision-support tools that explicitly incorporate the cumulative benefits of implementing actions across all species of interest, and the costs and feasibility of these actions, can help identify the most cost-effective management intervention and maximize the benefits to ecosystems and society (Carwardine et al., 2008; Evans et al., 2015). In addition, analyses of the cost-effectiveness that

account for complementarity across species can achieve similar conservation outcomes as simple ranking methods but at much lower cost (Chadés et al., 2015). The Priority Threat Management (PTM) framework (Carwardine et al., 2012, 2019; Chadés et al., 2015) accounts for the costs, benefits, feasibility and complementarity of actions. This framework uses structured expert elicitation to evaluate the cost-effectiveness of strategies that mitigate threats to biodiversity, and identify which strategies would achieve the greatest conservation outcomes across multiple taxa for any given budget. The PTM framework has been applied extensively in Australia (e.g. Carwardine et al., 2012; Chadés et al., 2015) and Canada (Kehoe et al., 2020; Martin et al., 2018). Additional benefits of the PTM framework include greater clarity of objectives, incorporation of Indigenous values and priorities, stakeholder engagement, data centralization and an emphasis on developing baselines necessary to evaluate management performance (Carwardine et al., 2019). PTM uses scientific information, local and traditional knowledge and expert elicitation to identify where and how to restore imperilled populations. Thus, there is an opportunity to apply PTM to guide systematic planning efforts for Pacific salmon.

We applied the PTM framework to a genus of exceptional economic and cultural importance: Pacific salmon and, in particular, salmon of the Central Coast of British Columbia (BC), Canada. Past PTM exercises have focused on species of conservation concern as their unit of interest. Instead, we focused on populations within and

across species, given the importance of genetically distinct salmon populations, and included all levels of threat status (rather than considering only threatened populations). Our project was a collaboration among academics, NGO staff, federal scientists, the Heiltsuk, Nuxalk, Kitasoo/Xai'xais and Wuikinuxv Nations—who have lived in the region for at least 14,000 years (McLaren et al., 2018, 2020) and have relied on salmon for at least the past 7,500 years (Campbell & Butler, 2010; Cannon & Yang, 2006), and the Central Coast Indigenous Resource Alliance (CCIRA—the non-profit organization these Nations created for technical support in marine resource management). Our case study, therefore, incorporates the expertise and perspectives of Indigenous people, which are essential from both ecological and social standpoints (Ban et al., 2018), into the development and evaluation of conservation strategies for the cost-effective recovery of Pacific salmon.

2 | MATERIALS AND METHODS

2.1 | Study area

We focused on the Central Coast of BC, including the traditional territories of the four First Nations involved (Figure 1; Figures S1 and S2; Marine Planning Partnership Initiative, 2015; Walsh et al., 2020). This region is in temperate rainforest, with relatively low levels of




FIGURE 1 Conservation units (CUs) for Chinook and chum salmon within the Central Coast of British Columbia, Canada. The study region is outlined in green, showing the combined traditional use territories of the Heiltsuk, Kitasoo/Xai'xais, Nuxalk and Wuikinuxv Nations (total area = 55,266 km²). Due to space limitations, maps of coho, pink, river-type sockeye and lake-type sockeye salmon CUs are in Figures S1 and S2

industrial development, supporting hundreds of wild salmon spawning locations that can be delineated into 79 geographically, ecologically and genetically unique populations of five species: chum *O. keta*, coho *O. kisutch*, Chinook *O. tshawytscha*, pink *O. gorbuscha* and sockeye *O. nerka* salmon. These groups are called conservation units (CUs) under Canada's WSP (Fisheries and Oceans Canada, 2005). Four CUs in this region are in the red status zone (at or below their lower biological benchmarks; Holt, Cass, Holtby, & Riddell, 2009) and 22 CUs are in the amber zone (between the lower and upper biological benchmarks), indicating the need for increased management intervention (Table 1, Connors et al., 2018, Pacific Salmon Explorer, <http://www.salmonexplorer.ca/>). Twelve CUs are in the green status zone, which suggests relatively high recent spawning abundances compared to biological benchmarks (Table 1). However, it should be noted that over half of the CUs on the Central Coast have insufficient data to assess their biological status (Table 1).

2.2 | Priority Threat Management framework

The PTM framework (Carwardine et al., 2012, 2019) is an eight-step process that quantifies the cost-effectiveness of management strategies for meeting a stated biodiversity objective, by identifying strategies that abate threats to the biodiversity features of interest (in this case salmon CUs), and estimating the costs, feasibility and benefits of each strategy (Figure 2). We performed the first three steps of the PTM process in May 2018 during a day-long workshop in Vancouver, BC, with six First Nation representatives and staff from CCIRA ($n = 3$), Pacific Salmon Foundation ($n = 4$), Fisheries and Oceans Canada ($n = 1$) and the research team ($n = 5$), followed by several conference calls and emails with other resource managers. During a 2-day workshop in June 2018, a panel of 19 experts

quantified the costs, feasibility and benefits of the previously identified management strategies (steps 4 & 5). These people had expertise in salmon threats, the feasibility and costs of conservation actions in the Central Coast and/or the ecological response of salmon to each action. Experts included the Heiltsuk, Nuxalk, Kitasoo/Xai'xais and Wuikinuxv Nations' natural resource managers ($n = 6$), academics ($n = 3$) and resource managers and scientists from government ($n = 7$) and non-governmental organizations ($n = 3$). Our research was approved by the Simon Fraser University Office of Research Ethics (permit 2018s0206). The major steps that we followed are briefly described below, with further details in Appendix S1.

2.2.1 | Define objective, scope and timeframe

The objective was to maximize the number of thriving Pacific salmon CUs on BC's Central Coast over the next 20 years. We defined a *thriving* CU as fulfilling its ecological function and role, and providing livelihood opportunities for present and future generations. In the context of the WSP, the objective is analogous to maximizing the number of CUs in the green status zone, which reduces the need for conservation intervention and allows for fishing opportunities, including for First Nation Food, Social, and Ceremonial (FSC) purposes, and commercial and recreational sectors (Fisheries and Oceans Canada, 2005). This objective was defined by the First Nations and experts during the workshops.


2.2.2 | Identify salmon CUs to conserve

The 'biodiversity features' we included in our analysis were all 79 salmon CUs that overlap with our study area. Including CUs with green, amber, red and data-deficient (DD) status ensures that conservation

CU groups	Green	Amber	Red	Data deficient (DD)	Total CUs/group
Chinook salmon		5	1		6
Chum salmon		5	2	2	9
Coho salmon	1	5			6
Pink salmon	3	2			5
Coastal and inland lake-type sockeye salmon—green status	8				8
Coastal lake-type sockeye—amber, red or DD status		4		31	35
Inland lake-type sockeye salmon—amber, red or DD status		1		5	6
South Atnarko lake-type sockeye salmon CU			1		1
River-type sockeye salmon				3	3
All	12	22	4	41	79

TABLE 1 Pacific salmon species and the biological status of their conservation units (CUs) on British Columbia's Central Coast (Figure 1), which were grouped into nine CU groups for the analysis. Sockeye CUs were divided into CU groups based on their ecotype, distribution and biological status (Connors et al., 2018). Status presented here is based on historic spawner abundance benchmarks

FIGURE 2 Eight steps of the Priority Threat Management framework. Figure adapted from Carwardine et al. (2019)



strategies (Step 3) are designed to promote recovery of amber and red CUs while avoiding future declines in green status CUs. We grouped the 79 CUs into 'CU groups' by species (Table 1). Due to the large number of sockeye salmon CUs, the sockeye CUs were further divided into five groups based on their life-history ecotype (river-type or lake-type), geographic location (inland or coastal) and current level of threat (red/amber/DD status or green status). We assumed that each CU group experienced similar threats and would respond similarly to recovery actions within the region (Table 1). Grouping CUs made the exercise feasible by reducing the workload of experts when estimating the benefits of strategies (Step 5).

2.2.3 | Identify threats, strategies and actions















We identified threats based on previous summaries of human and environmental pressures in the region (Connors et al., 2018), and a literature review of broader threats to salmon, which were refined by experts at workshops, conference calls and meetings. These threats included overfishing, habitat loss and degradation due to logging, anthropogenic barriers to freshwater migration, future industrial development in salmon habitats, salmon aquaculture, hatcheries (due to competition and genetic introgression) and predation by marine mammals and other predators. Mitigating the effects of climate change in marine and freshwater environments was considered beyond the scope of the strategies we evaluated.

We developed an inventory of strategies, each consisting of several underlying actions (Appendix S2) as well as three combinations of strategies, as experts thought their combined benefits could be synergistic (Table 2). We also developed an enabling strategy that included increased monitoring and assessment of salmon CUs, which experts considered essential for effective implementation and evaluation of all other strategies. This enabling strategy was not included in the cost-effectiveness analysis because it would not directly result in recovery, but it was assumed that it would be implemented in addition to any other strategy.

2.2.4 | Estimate the costs and feasibility of actions

The costs and feasibility of each action within each strategy were quantified during the second workshop and finalized in follow-up communication (Appendix S1). The cost of labour, consumables and equipment, capital assets, overheads, monitoring and coordination were estimated for each action, drawing on salmon management reports and the experts' experience (Iacona et al., 2018). All costs are presented in Canadian dollars (2018 CAD). We included management and transaction costs, but did not include opportunity costs (i.e. foregone profits) in our estimates, and provide justification for this in Appendix S1 (Naidoo et al., 2006). The feasibility of each strategy was quantified as the product of the average probability of uptake and the average probability of success, across each action.

TABLE 2 Benefits, costs, feasibility and cost-effectiveness (E) of each strategy designed to conserve salmon on British Columbia's Central Coast. Strategies are ordered by cost-effectiveness from the ranking method. Detailed actions for each strategy are described in Appendix S2. Benefit is the difference between the probability of achieving the conservation objective (i.e. thriving conservation units [CUs] with green status) after 20 years under each strategy and the baseline scenario per CU group, averaged across experts and summed across all CU groups (Equation 1). Cost is the average annual present value divided across the 20-year time frame. Feasibility is the probability of uptake multiplied by the probability of success

	Conservation strategy	Summary of underlying actions	Benefit	Cost (CAD millions)	Feasibility	E	E rank
	Limit future industrial development	Restrict future industrial developments in salmon habitat, manage water pollution; assess risks of shipping traffic to salmon	1,490	0.15	0.74	362	1
	Predation control	Conduct experimental culls (or traditional First Nation harvest) of pinnipeds; reduce juvenile predation by trout & sculpin	1,241	0.11	0.46	253	2
	Remove barriers to fish passage	Remove significant barriers (e.g. culverts) to upstream adult migration and juvenile dispersal	1,395	0.21	0.72	239	3
	Salmon aquaculture management	Implement salmon aquaculture best practices; develop siting guidelines for aquaculture licenses; incentivize land-based aquaculture	772	0.09	0.34	147	4
	Watershed protection	Improve forestry practices; restore watershed vegetation to pre-logging conditions	1,407	0.54	0.7	92	5
	Marine and estuary habitat restoration	Restore and protect eelgrass and other important salmon rearing estuary habitats	1,293	0.79	0.56	46	6
	Sustainable commercial harvest	Reduce mixed-stock catches; improve enforcement of regulations; reduce bycatch of non-target salmon species	1,406	0.59	0.38	45	7
	Sustainable recreational harvest	Limit expansion of tourism and sport fishing operations; regulate size and daily catch limits; improve monitoring	707	0.99	0.73	26	9
	Supplement small populations with hatcheries	Set up hatcheries as a last resort for CUs in the red zone; monitor and improve effectiveness of hatcheries supplementing target CUs; fish ladders over natural barriers where appropriate	1,102	3.27	0.72	12	11
	Stream restoration	Fine-scale habitat monitoring; maintain and restore riparian habitat characteristics and processes	1,421	10.55	0.76	5	13
	Combined harvest strategy	Overarching harvest strategy; Sustainable commercial harvest strategy; Sustainable recreational harvest strategy	1,818	1.59	0.59	34	8
	Combined supplementation and predation strategy	Supplement small populations; Predator control	1,911	3.39	0.57	16	10
	Combined habitat strategy	Watershed protection and water management; Stream restoration; Remove barriers to fish passage	2,366	11.30	0.73	8	12
	All strategies combined		2,896	17.30	0.59	5	14
	Enabling strategy: Monitoring and assessment	Integrated status assessments for CUs; adult escapement and smolt abundance monitoring for CU indicator streams; monitor data-deficient CUs	NA	0.699	0.71	NA	NA

2.2.5 | Identify benefits of strategies

We used a structured expert elicitation approach (Hemming, Burgman, Hanea, McBride, & Wintle, 2018; Martin, Burgman, et al., 2012; Martin, Nally, et al., 2012) to elicit the benefits of each strategy for the nine groups of salmon CUs. We used a modified Delphi method whereby experts provided their initial benefit estimates, and then in a second round of scoring had the chance to adjust their estimates after seeing box plots summarizing anonymous estimates by other experts (Mukherjee et al., 2015). Fourteen experts completed round 1, and 10 experts (k) completed round 2. Estimates from round 2 were used to calculate the benefit of each strategy. In each round, experts individually estimated the likely response of each strategy (i) for each group of CUs (j), defined as the probability that on average each CU within the group would be thriving after 20 years (P_{ijk} , Figure S3), along with upper and lower limits and estimates of confidence on a scale of 0–100. Experts also estimated the probability of each CU group achieving this objective in 20 years if no additional management strategies were implemented, which served as a business-as-usual baseline scenario (P_{0jk}). Experts were asked to consider the effects of climate change and other emerging threats on freshwater and marine habitats, and existing and ongoing management in this baseline scenario. Experts agreed that it was biologically conservative to assume DD CUs are currently in the red status zone (i.e. of conservation concern and below their lower biological benchmark) to ensure estimates of future scenarios were precautionary. Further details on how benefits were estimated are described in Appendix S1.

Following Carwardine et al. (2019), the benefit (B_{ij}) of a strategy for a given CU group was calculated as the change in the probability of achieving the objective under the baseline scenario, compared with the estimated benefit if the strategy was implemented, averaged across experts (M_{ij} is the number of experts who made predictions for that CU–strategy combination).

$$B_{ij} = \frac{\sum_{k=1}^{M_{ij}} (P_{ijk} - P_{0jk})}{M_{ij}}. \quad (1)$$

2.2.6 | Cost-effectiveness calculation

First, using the simple ranking method, we ranked strategies based on their cost-effectiveness (E_i ; Carwardine et al., 2012), calculated as the sum of the cumulative benefits, multiplied by the feasibility per strategy (F_i), divided by the costs per strategy (C_i):

$$E_i = \frac{B_i F_i}{C_i}, \quad (2)$$

where the total benefit per strategy (B_i) is the sum of the benefits across all CU groups, weighted by the number of CUs occurring in that group (N_j):

$$B_i = \sum_{j=1}^N B_{ij} N_j. \quad (3)$$

We then used the complementarity approach to identify strategies and their combinations that are predicted to maximize the probability of achieving as many thriving salmon CUs as possible for a given budget (Chadés et al., 2015). For example, rather than choosing two cost-effective strategies that benefit the same CU group, the analysis would select one of those strategies, and another slightly more expensive strategy that benefits a different CU group, thus increasing the number of CUs conserved. To assess complementarity, arbitrary thresholds of ‘conservation’ must be defined to determine the number of CUs that have reached a satisfactory level of protection under any given budget. Here, we used conservation thresholds of 50%, 60% and 70% probability of achieving the objective—i.e. being a thriving CU with green status in 20 years.

Uncertainty in the experts’ assessments were assessed by conducting sensitivity analyses to test if the cost-effectiveness results from the simple ranking and complementarity methods were robust to upper and lower estimates of the costs and benefits of each strategy (Appendix S1).

3 | RESULTS

3.1 | Predicted outcomes if no or all strategies were implemented

Under a business-as-usual scenario (i.e. the baseline scenario with no additional investment), only one quarter of salmon CUs on BC’s Central Coast were predicted to have a >50% chance of achieving the conservation objective within the next 20 years (i.e. 6 coho, 5 pink and 8 ‘green-status lake-type sockeye’ salmon CUs of all 79 CUs, Table S1; Figure S3). No CUs were predicted to have a >70% probability of reaching this conservation objective under business-as-usual (Table S1; Figure 3; Figure S3). Without additional conservation strategies, declines were predicted for the CUs that currently have green status, such as the 8 CUs in the ‘green-status lake-type’ sockeye salmon group, which were estimated to have 62% probability of thriving in 20 years (Table S1). There was considerable variation in the experts’ best guess estimates of the probability of these CUs still thriving in 20 years (Figure S3). The lower estimates of expert predictions suggested that under the business-as-usual scenario, no CUs would have a >50% probability of thriving in 20 years and the upper estimates suggest that 19 CUs would have >70% probability of thriving.

Conversely, if all 10 strategies were implemented, all CUs (except the South Atnarko Lake Sockeye CU) were predicted to have >50% probability of thriving in 20 years (Figure 3). However, only 43% of all CUs—including 6 Chinook, 9 chum, 6 coho, 5 pink and 8 ‘green-status lake-type sockeye’ salmon—were estimated to reach >60% probability of thriving (Table S1). Amber, red or DD status sockeye CUs were predicted to have <60% probability of meeting the conservation objective even with the implementation of all strategies (Table S1). The experts’ lower estimates suggested that with all 10 strategies, only 5 pink and 8 ‘green-status lake-type sockeye’




FIGURE 3 The number of Pacific salmon conservation units (CUs) that would exceed three threshold probabilities (>50%, >60% and >70%) at different levels of annual investment if complementary sets of strategies were implemented (total = 79 CUs). For example, with an annual budget of \$11.3M, the ‘Combined Habitat Strategy’ (i.e. watershed protection, stream restoration and removal of barriers to fish passage) could be implemented, resulting in 75 CUs with >50% probability of recovery, 34 CUs with >60% probability of recovery and 19 CUs with >70% probability of recovery (shown in Table S1)

salmon CUs would reach >50% probability of achieving thriving status, while the upper estimates suggest that all CUs would reach the >60% threshold, and 34 CUs would have >70% probability of reaching the conservation objective.

The average annual equivalent cost of carrying out all strategies was predicted to be \$17.3M/year (min. and max. estimates = \$16.1–\$41.4M/year; Table 2). To support the successful implementation of strategies, an additional \$0.7M/year would be required to conduct the necessary monitoring and assessment of biological status for all salmon CUs (i.e. enabling strategy in Table 2).

3.2 | Cost-effectiveness of strategies

Not all strategies performed equally when comparing their expected benefit (benefit × feasibility) per dollar spent. The most cost-effective strategy predicted to conserve salmon CUs was to ‘Limit Future Industrial Development’ (Table 2; Figure S4). This strategy aimed to restrict industrial development (e.g. oil and gas infrastructure, aquaculture, wind farms) in critical areas of salmon spawning and rearing habitat by ensuring the integrity of these habitats is maintained and safeguarded from potential future degradation associated with development. This strategy had the highest benefit to species, and a relatively low cost of \$150,000/year for the next 20 years (compared to the average annual cost of individual strategies = \$1.73M). On average across all CU groups, ‘Limiting Future Industrial Development’ would result in an increase of 12.6% ($SD = 2.6\%$) probability of achieving a thriving CU compared to the baseline scenario (Table S1). The second most cost-effective strategy was predicted to be ‘Predation

Control’ (i.e. predator culls which may include—but not be limited to—revitalizing traditional First Nation harvests for abundant pin-nipeds and trapping of predatory fishes that consume juvenile salmon; Table 2; Figure S4), which also had a relatively low annual cost of \$0.11M.

The ‘Combined Habitat Strategy’—including ‘Watershed Protection’, ‘Stream Restoration’ and ‘Removal of Barriers to Fish Passage’—was predicted to be highly beneficial, but had a low cost-effectiveness rank due to higher costs of implementation relative to other strategies (Table 2; Figures S4 and S5). ‘Stream Restoration’ and ‘Supplement Small Populations’ with hatcheries and other means of enhancement were considered the least cost-effective strategies due primarily to their high costs (Table 2; Figure S4).

3.3 | Complementary strategies to maximize conservation success

We conducted a complementarity analysis to identify which strategies would achieve recovery (or maintain thriving status) for the largest number of CUs under various budgets. Two clear investment thresholds emerge that are consistent across all conservation objective thresholds (Figure 3; Table S1). The first investment threshold is \$0.15M/year, to fund the most cost-effective strategy overall ‘Limit Future Industrial Development’. This strategy was predicted to lead to 19 CUs reaching >60% probability of thriving and 13 CUs having >70% probability of thriving. Next, for a budget of \$11.3M, investing in the ‘Combined Habitat Strategy’ was predicted to deliver the highest number of CU’s conserved across all objective thresholds (34 CUs would reach >60% probability of thriving and 19 CUs would have >70% probability of thriving). Without the ‘Combined Habitat

Strategy', over half of the CUs were predicted to have <50% probability of recovery (Figure 3; Table S1). Further investment above \$11.3M would benefit only three river-type sockeye CUs by increasing their probability of recovery to >50% at an extra \$6M (Figure 3).

3.4 | Sensitivity analysis

There was variation among the experts in the predicted costs and benefits of different strategies. Using the lower cost estimates, 'Removal of Barriers to Fish Passage' became the most cost-effective strategy (Table S2) and resulted in the same probability of achieving the conservation objective as 'Limiting Future Industrial Development' for less money (\$69,900/year to remove five barriers over 20 years) under the 60% and 70% thresholds (Figure S7). In the upper-cost scenario, the 'Combined Habitat Strategy' was 213% more expensive than the best-guess estimate, raising the average annual cost from \$11M to \$35M (Table S2), but was still selected in the complementarity analysis as it was the best strategy at improving the probability of success to >50% for 95% CUs (Figure S7).

There was also wide uncertainty in the probability of each CU group achieving the objective under each strategy (Figures S3 and S8). Experts predicted that under the worst-case scenario (i.e. experts' lower benefit estimates), no CU would reach the >60% threshold under any strategy; the optimistic scenario (i.e. experts' upper benefit estimates) suggested that most CUs would achieve a >60% probability of recovery if 'Limiting Future Industrial Development' or 'Predation Control' strategies were implemented (Figure S8).

4 | DISCUSSION

The PTM framework we applied for BC's Central Coast is a decision-support tool that can help guide conservation investment to maximize the benefit to Pacific salmon. We integrated Indigenous local knowledge into the analysis, through the design of the objectives and strategies, as well as their expert knowledge on the cost, feasibility and benefits, which we believe produced more relevant, inclusive and legitimate findings (Ban et al., 2018). This process relied on existing relationships with First Nations, and benefited from coordinated efforts led by the Pacific Salmon Foundation and CCIRA. Our study also demonstrated that the PTM framework can be applied not only to groups of threatened taxa or species but also can include all levels of threat, and populations and ecotypes within species.

4.1 | Strategic planning in an era of high stakes for salmon

The strategic planning process revealed the urgency for conservation strategies and actions to support the recovery of Pacific salmon in Canada. Our analysis suggested a 'business-as-usual' baseline scenario will be insufficient to recover or maintain thriving salmon populations

along BC's Central Coast. Three quarters of CUs were estimated to have <50% chance of achieving this objective if no additional strategies are implemented (Table S1). However, if resources are allocated in a strategic manner, our analyses suggest there is substantial scope to improve the overall status of salmon on BC's Central Coast. In addition to the predicted benefits to salmon populations, the social, economic and ecological benefits of implementing these proposed strategies potentially include: job provision, secure and stable fishing opportunities, health and well-being for affected communities, increased habitat protection and greater opportunities for First Nation stewardship and governance.

The choice of decision-support framework is important because not all prioritization methods produce results that are equally cost-effective and beneficial (Giakoumi et al., 2013). If the goal is to conserve the most CUs per dollar spent, the associated costs, benefits and feasibility of strategies should be compared across multiple species and CUs in a complementarity exercise to avoid inefficient spending and maximize conservation outcomes (Martin et al., 2018). PTM has the potential to be a useful approach to guide the strategic and transparent allocation of new funding to maximize the return on investment for Pacific salmon.

At the regional level, PTM could serve as a model for agencies or First Nations interested in strategic planning and, at the national level, it could inform the implementation of existing legislation and policies, such as the Canadian Species at Risk Act and the Canadian WSP; both under restrictive budgets. There also is scope to explicitly include Indigenous social and cultural values of salmon in future PTM exercises by weighting different CU groups according to their relative importance to First Nation communities. This would provide a mechanism for ensuring that resources are preferentially directed to those CUs tied to First Nations cultures and communities.

4.2 | Cost-effective strategies for salmon conservation

The choice of strategy to maximize gains in salmon recovery on the Central Coast ultimately depends on the resources available (Figure 3). Our analyses suggest that there are a few relatively inexpensive strategies that could provide immediate—though marginal—benefits to many salmon CUs, regardless of their biological status. 'Limiting Future Industrial Development' in critical spawning, rearing and migration habitats was identified as a cost-effective option to safeguard and recover CUs, for an additional \$150,000/year over the next 20 years above existing funding (Figure 3; Figure S6). Regardless of the conservation strategy chosen, the cost of monitoring and status assessment of salmon CUs (\$0.7M/year) needs to be added to any budget, as this was considered necessary by experts. At budgets <\$2M/year, we consider it would be important for decision-makers to decide which target threshold probability to aim for (i.e. >50%, >60% or >70%), as the most cost-effective strategy depended on the chosen threshold (Figure 3).

If a larger budget of \$11.3M was available per year, the 'Combined Habitat Strategies' were predicted to be highly beneficial. These

actions include: (a) protecting and restoring watersheds and hydrology from forestry impacts, (b) restoring stream habitat to increase egg and juvenile survival and (c) removal of barriers that limit fish passage and migration (Table 2). While these habitat strategies also had the highest uncertainty regarding the extent of habitat restoration and barrier removal required—leading to highly variable costs—they consistently outperformed other strategies in the sensitivity analyses (Figure S7). Future research could refine these cost estimates.

Several uncertainties and caveats should be considered if implementing these strategies. It is inherently difficult for experts to predict the benefit of strategies into the future while accounting for multiple threats and underlying variable marine conditions that may reduce survival (e.g. Malick & Cox, 2016; Peterman & Dorner, 2012). This uncertainty is compounded by the logistical need to group together CUs that may have distinct (or unknown) biological status, life histories or threats. Management strategy evaluation frameworks (i.e. simulation modelling to quantify the predicted ability of strategies to meet multiple objectives) could be used to estimate expected outcomes while accounting for pervasive uncertainty (Punt, Butterworth, de Moor, De Oliveira, & Haddon, 2014), although pursuits for more knowledge should not be an excuse for delays in action (Martin, Burgman, et al., 2012; Martin, Nally, et al., 2012). Further consideration of the opportunity costs, economic benefits, social preferences and risks of these strategies, such as industrial development, sustainable commercial harvest and forestry, would provide a more complete assessment of the costs and benefits of actions within the broader conservation and development objectives in the region.

5 | CONCLUSIONS

The PTM framework is a systematic approach that can be used to quantify trade-offs between costs and benefits of a diverse suite of conservation strategies for Pacific salmon. The estimated benefits resulting from cost-effective strategies, combined with the cultural and economic importance of salmon to First Nations in this region, underscore the importance of prioritizing conservation strategies for Pacific salmon and their habitats in a systematic way. The status of salmon CUs in the Central Coast is relatively good compared to other regions such as southern BC, Washington and Oregon. This allows for a wider array of conservation management options, including proactive strategies that aim to safeguard CUs from the future risk of development. Even in regions with higher risks of local extirpation—where proposed strategies may be more expensive, less feasible and may result in lower probabilities of recovery—the PTM framework could provide a structured and transparent method to identify the most cost-effective options.

ACKNOWLEDGEMENTS

We thank CCIRA and Stewardship Offices' staff from the Heiltsuk, Nuxalk, Kitasoo/Xai'xais and Wuikinuxv Nations for their support. We thank the following individuals for providing valuable advice and feedback: P. Siwallace, M. Reid, D. Chan, D. Rolston, B. Edgar, D. Neasloss,

D. Dobson, D. Stewart and all experts. Thank you to the Salmon Watersheds Program staff at Pacific Salmon Foundation for use of the Pacific Salmon Explorer, K. Kellock for producing maps and E. Jones, J. Belzile, C. Stevenson and L. Chalifour for facilitating workshop discussions. We thank D. Braun, W. Atlas & J. Walkus for comments on the manuscript. Mitacs, the Central Coast Indigenous Resource Alliance, Pacific Salmon Foundation and the Tom Buell Endowment Fund funded this project. This work was supported by Environment and Climate Change Canada and a NSERC Discovery Grant to T.G.M. Icon credits: Dan Hetteix from the Noun Project (creek), Freepik from Flaticon.com (seal) and Microsoft Office Professional Plus 2019. Infographic was designed by Kate at Fuse Consulting.

AUTHORS' CONTRIBUTIONS

J.C.W., K.C., E.H., T.G.M., L.K., B.C. and J.D.R. conceived the ideas and designed methodology; J.C.W., K.C., E.H., T.G.M., L.K., B.C., J.H. and J.D.R. collected the data; J.C.W. and L.K. analysed the data; J.C.W. led the project and writing of the manuscript; K.C., E.H., B.C., M.J.B., C.F., A.F., J.W.M., M.H.H.P. and J.D.R. participated as experts in the elicitation process. All authors contributed critically to drafts and gave final approval for publication.

DATA AVAILABILITY STATEMENT

Data underlying the results are available via the Simon Fraser University Radar data repository <https://dx.doi.org/10.25314/8a9a45cf-7dfb-4180-adc3-5f33e5669415> (Reynolds et al., 2020). Please note that personal data are removed and summarized data across experts are provided.

ORCID

Jessica C. Walsh  <https://orcid.org/0000-0002-5284-4323>

Alejandro Frid  <https://orcid.org/0000-0003-4844-5562>

John D. Reynolds  <https://orcid.org/0000-0002-0459-0074>

REFERENCES

- Ban, N. C., Frid, A., Reid, M., Edgar, B., Shaw, D., & Siwallace, P. (2018). Incorporate Indigenous perspectives for impactful research and effective management. *Nature Ecology and Evolution*, 2(11), 1680–1683. <https://doi.org/10.1038/s41559-018-0706-0>
- Barnas, K. A., Katz, S. L., Hamm, D. E., Dias, M. C., & Jordan, C. E. (2015). Is habitat restoration targeting relevant ecological needs for endangered species? Using Pacific salmon as a case study. *Ecosphere*, 6(7), 110. <https://doi.org/10.1890/ES14-00466.1>
- Beechie, T., Pess, G., Roni, P., & Giannico, G. (2008). Setting river restoration priorities: A review of approaches and a general protocol for identifying and prioritizing actions. *North American Journal of Fisheries Management*, 28(3), 891–905. <https://doi.org/10.1577/M06-174.1>
- Bernhardt, E. S., Palmer, M. A., Allan, J. D., Alexander, G., Barnas, K., Brooks, S., ... Sudduth, E. (2005). Synthesizing U.S. river restoration efforts. *Science*, 308(5722), 636–637. <https://doi.org/10.1126/science.1109769>
- Campbell, S. K., & Butler, V. L. (2010). Archaeological evidence for resilience of Pacific northwest salmon populations and the socioecological system over the last ~7,500 years. *Ecology and Society*, 15(1), 17. Retrieved from <http://www.ecologyandsociety.org/vol15/iss1/art17/>

- Cannon, A., & Yang, D. Y. (2006). Early storage and sedentism on the Pacific Northwest Coast: Ancient DNA analysis of salmon remains from Namu, British Columbia. *American Antiquity*, 71(1), 123–140. <https://doi.org/10.2307/40035324>
- Carwardine, J., Martin, T. G., Firn, J., Reyes, R. P., Nicol, S., Reeson, A., ... Chadés, I. (2019). Priority Threat Management for biodiversity conservation: A handbook. *Journal of Applied Ecology*, 56(2), 481–490. <https://doi.org/10.1111/1365-2664.13268>
- Carwardine, J., O'Connor, T., Legge, S., Mackey, B., Possingham, H. P., & Martin, T. G. (2012). Prioritizing threat management for biodiversity conservation. *Conservation Letters*, 5(3), 196–204. <https://doi.org/10.1111/j.1755-263X.2012.00228.x>
- Carwardine, J., Wilson, K. A., Watts, M., Etter, A., Klein, C. J., & Possingham, H. P. (2008). Avoiding costly conservation mistakes: The importance of defining actions and costs in spatial priority setting. *PLoS ONE*, 3(7), e2586. <https://doi.org/10.1371/journal.pone.0002586>
- Chadés, I., Nicol, S., van Leeuwen, S., Walters, B., Firn, J., Reeson, A., ... Carwardine, J. (2015). Benefits of integrating complementarity into priority threat management. *Conservation Biology*, 29(2), 525–536. <https://doi.org/10.1111/cobi.12413>
- Connors, K., Jones, E., Kellock, K., Hertz, E., Honka, L., & Belzile, J. (2018). *BC Central Coast: A snapshot of salmon populations and their habitats. Technical report*. Vancouver, BC: Pacific Salmon Foundation. Retrieved from https://salmonwatersheds.ca/library/lib_442/
- Cultus Sockeye Recovery Team. (2009). National conservation strategy for Cultus Lake sockeye salmon (*Oncorhynchus nerka*). *Canadian Technical Report of Fisheries and Aquatic Sciences*, 2846, viii + 46 p.
- Evans, M. C., Tulloch, A. I. T., Law, E. A., Raiter, K. G., Possingham, H. P., & Wilson, K. A. (2015). Clear consideration of costs, condition and conservation benefits yields better planning outcomes. *Biological Conservation*, 191(November), 716–727. <https://doi.org/10.1016/j.biocon.2015.08.023>
- Fisheries and Oceans Canada. (2005). *Canada's policy for conservation of wild Pacific salmon*. Vancouver, BC. Retrieved from <http://www.pac.dfo-mpo.gc.ca/fm-gp/salmon-saumon/wsp-pss/policy-politique/index-eng.html>
- Fisheries and Oceans Canada. (2018). *Wild Salmon Policy 2018 to 2022 Implementation Plan*. Vancouver, BC. Retrieved from <https://www.pac.dfo-mpo.gc.ca/fm-gp/species-especes/salmon-saumon/wsp-pss/ip-pmo/intro-eng.html>
- Gardner, M. P., & Pinfold, T. A. (2011). *Performance review of the Wild Salmon Policy*. Sechelt, BC, Canada: Prepared by Gardner Pinfold Consultants for Fisheries and Oceans Canada, 67 pp.
- Garibaldi, A., & Turner, N. (2004). Cultural keystone species: Implications for ecological conservation and restoration. *Ecology and Society*, 9(3), 1–18. <http://www.ecologyandsociety.org/vol9/iss3/art1/>
- Giakoumi, S., Sini, M., Gerovasileiou, V., Mazor, T., Beher, J., Possingham, H. P., ... Katsanevakis, S. (2013). Ecoregion-based conservation planning in the Mediterranean: Dealing with large-scale heterogeneity. *PLoS ONE*, 8(10), e76449. <https://doi.org/10.1371/journal.pone.0076449>
- Good, T. P., Beechie, T. J., McElhany, P., McClure, M. M., & Ruckelshaus, M. H. (2007). Recovery planning for Endangered Species Act-listed Pacific salmon: Using science to inform goals and strategies. *Fisheries*, 32(9), 426–440. [https://doi.org/10.1577/1548-8446\(2007\)32\[426:RPFESL\]2.0.CO;2](https://doi.org/10.1577/1548-8446(2007)32[426:RPFESL]2.0.CO;2)
- Governor's Salmon Recovery Office. (2018). *2018 State of the salmon in watersheds*. Washington State Recreation and Conservation Office. Retrieved from <https://stateofsalmon.wa.gov/>
- Gustafson, R. G., Waples, R. S., Myers, J. M., Weitkamp, L. A., Bryant, G. J., Johnson, O. W., & Hard, J. J. (2007). Pacific salmon extinctions: Quantifying lost and remaining diversity. *Conservation Biology*, 21(4), 1009–1020. <https://doi.org/10.1111/j.1523-1739.2007.00693.x>
- Hemming, V., Burgman, M. A., Hanea, A. M., McBride, M. F., & Wintle, B. C. (2018). A practical guide to structured expert elicitation using the IDEA protocol. *Methods in Ecology and Evolution*, 9(1), 169–180. <https://doi.org/10.1111/2041-210X.12857>
- Hoekstra, J. M., Bartz, K. K., Ruckelshaus, M. H., Moslemi, J. M., & Harms, T. K. (2007). Quantitative threat analysis for management of an imperiled species: Chinook Salmon (*Oncorhynchus tshawytscha*). *Ecological Applications*, 17(7), 2061–2073. <https://doi.org/10.1890/06-1637.1>
- Holt, C. A., Cass, A., Holtby, B., & Riddell, B. (2009). *Indicators of status and benchmarks for conservation units in Canada's Wild Salmon Policy*. Nanaimo, Canada: Department of Fisheries and Oceans Canada, Canadian Science Advisory Secretariat Research Document 2009/058. viii + 74 pp. Retrieved from <https://waves-vagues.dfo-mpo.gc.ca/Library/339096.pdf>
- Iacona, G. D., Sutherland, W. J., Mappin, B., Adams, V. M., Armsworth, P. R., Coleshaw, T., ... Possingham, H. P. (2018). Standardized reporting of the costs of management interventions for biodiversity conservation. *Conservation Biology*, 32(5), 979–988. <https://doi.org/10.1111/cobi.13195>
- Kareiva, P., Marvier, M., & McClure, M. (2000). Recovery and management options for spring/summer Chinook salmon in the Columbia River Basin. *Science*, 290(5493), 977–979. <https://doi.org/10.1126/science.290.5493.977>
- Kehoe, L. J., Lund, J., Chalifour, L., Hinch, S. G., McDuffee, M., Richardson, J. S., ... Martin, T. G. (2020). Conservation in heavily urbanized biodiverse regions requires urgent management action and attention to governance. *Conservation Science and Practice*, in press.
- Levi, T., Darimont, C. T., MacDuffee, M., Mangel, M., Paquet, P., & Wilmers, C. C. (2012). Using grizzly bears to assess harvest-ecosystem tradeoffs in salmon fisheries. *PLoS Biology*, 10(4), e1001303. <https://doi.org/10.1371/journal.pbio.1001303>
- Malick, M. J., & Cox, S. P. (2016). Regional-scale declines in productivity of pink and chum salmon stocks in western North America. *PLoS ONE*, 11(1), e0146009. <https://doi.org/10.1371/journal.pone.0146009>
- Mantua, N. J., Taylor, N. G., Ruggerone, G. T., Myers, K. W., Preikshot, D., Augerot, X., ... Walters, C. J. (2009). The Salmon MALBEC project: A North Pacific-scale study to support salmon conservation planning. In *North Pacific Anadromous Fish Commission Bulletin Doc. No 1060*. Seattle, WA: School of Aquatic and Fishery Sciences, University of Washington, 49 pp. Retrieved from www.npafc.org
- Marine Planning Partnership Initiative. (2015). *Central Coast Marine Plan*. British Columbia, Canada: Marine Planning Partnership Initiative, Heiltsuk, Kitasoo/Xai'Xais, Nuxalk and Wuikinuxv Nations. Retrieved from http://mapocean.org/wp-content/uploads/2015/08/MarinePlan_CentralCoast_10082015.pdf
- Martin, T. G., Burgman, M. A., Fidler, F., Kuhnert, P. M., Low-Choy, S., McBride, M., & Mengersen, K. (2012). Eliciting expert knowledge in conservation science. *Conservation Biology*, 26(1), 29–38. <https://doi.org/10.1111/j.1523-1739.2011.01806.x>
- Martin, T. G., Kehoe, L., Mantyka-Pringle, C., Chadés, I., Wilson, S., Bloom, R. G., ... Smith, P. A. (2018). Prioritizing recovery funding to maximize conservation of endangered species. *Conservation Letters*, 11(6), e12604. <https://doi.org/10.1111/conl.12604>
- Martin, T. G., Nally, S., Burbidge, A. A., Arnall, S., Garnett, S. T., Hayward, M. W., ... Possingham, H. P. (2012). Acting fast helps avoid extinction. *Conservation Letters*, 5(4), 274–280. <https://doi.org/10.1111/j.1755-263X.2012.00239.x>
- McLaren, D., Fedje, D., Dyck, A., Mackie, Q., Gauvreau, A., & Cohen, J. (2018). Terminal Pleistocene epoch human footprints from the Pacific coast of Canada. *PLoS ONE*, 13(3), 12–15. <https://doi.org/10.1371/journal.pone.0193522>
- McLaren, D., Fedje, D., Mackie, Q., Davis, L. G., Erlandson, J., Gauvreau, A., & Vogelaar, C. (2020). Late pleistocene archaeological discovery models on the Pacific Coast of North America. *PaleoAmerica*, 6(1), 43–63. <https://doi.org/10.1080/20555563.2019.1670512>
- Mukherjee, N., Hugé, J., Sutherland, W. J., McNeill, J., Van Opstal, M., Dahdouh-Guebas, F., & Koedam, N. (2015). The Delphi technique

- in ecology and biological conservation: Applications and guidelines. *Methods in Ecology and Evolution*, 6(9), 1097–1109. <https://doi.org/10.1111/2041-210X.12387>
- Naidoo, R., Balmford, A., Ferraro, P. J., Polasky, S., Ricketts, T. H., & Rouget, M. (2006). Integrating economic costs into conservation planning. *Trends in Ecology & Evolution*, 21(12), 681–687. <https://doi.org/10.1016/j.tree.2006.10.003>
- Nelitz, M., Murray, C., & Wieckowski, K. (2008). *Returning salmon: Integrated planning and the Wild Salmon Policy in B.C.* Vancouver, Canada: ESSA Technologies for the David Suzuki Foundation.
- Ogden, A. D., Irvine, J. R., O'Brien, M., Komick, N., Brown, G., & Tompkins, A. (2014). *Canadian commercial catches and escapements of Chinook and coho salmon separated into hatchery- and wild-origin fish.* NPAFC Doc. 1531. Retrieved from <https://npafc.org/wp-content/uploads/Public-Documents/2014/1531Canada.pdf>
- Peterman, R. M., & Dorner, B. (2012). A widespread decrease in productivity of sockeye salmon (*Oncorhynchus nerka*) populations in western North America. *Canadian Journal of Fisheries and Aquatic Sciences*, 69(8), 1255–1260. <https://doi.org/10.1139/f2012-063>
- Price, M. H. H., English, K. K., Rosenberger, A. G., MacDuffee, M., & Reynolds, J. D. (2017). Canada's Wild Salmon Policy: An assessment of conservation progress in British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences*, 74(10), 1507–1518. <https://doi.org/10.1139/cjfas-2017-0127>
- Punt, A. E., Butterworth, D. S., de Moor, C. L., De Oliveira, J. A. A., & Haddon, M. (2014). Management strategy evaluation: Best practices. *Fish and Fisheries*, 17(2), 303–334. <https://doi.org/10.1111/faf.12104>
- Reynolds, J. D., Walsh, J. C., Connors, K., Hertz, E., Kehoe, L., Martin, T. G., ... Price, M. H. H. (2020). *Priority threat management for Pacific salmon conservation on the Central Coast of British Columbia: Expert elicitation data.* <https://doi.org/10.25314/8a9a45cf-7dfb-4180-adc3-5f33e5669415>
- Roni, P., Beechie, T. J., Bilby, R. E., Leonetti, F. E., Pollock, M. M., & Pess, G. R. (2002). A review of stream restoration techniques and a hierarchical strategy for prioritizing restoration in Pacific Northwest Watersheds. *North American Journal of Fisheries Management*, 22(1), 1–20. [https://doi.org/10.1577/1548-8675\(2002\)022<0001:AROSRT>2.0.CO;2](https://doi.org/10.1577/1548-8675(2002)022<0001:AROSRT>2.0.CO;2)
- Ruckelshaus, M. H., Levin, P., Johnson, J. B., & Kareiva, P. M. (2002). The Pacific salmon wars: What science brings to the challenge of recovering species. *Annual Review of Ecology and Systematics*, 33, 665–706. <https://doi.org/10.1146/annurev.ecolsys.33.010802.150504>
- Sakinaw Lake Sockeye Recovery Team. (2005). *National recovery strategy for sockeye salmon (Oncorhynchus nerka), Sakinaw Lake population, in British Columbia.* Ottawa, Canada: Recovery of Nationally Endangered Wildlife.
- Schoen, E. R., Wipfli, M. S., Trammell, E. J., Rinella, D. J., Floyd, A. L., Grunblatt, J., ... Witmer, F. D. W. (2017). Future of Pacific salmon in the face of environmental change: Lessons from one of the world's remaining productive salmon regions. *Fisheries*, 42(10), 538–553. <https://doi.org/10.1080/03632415.2017.1374251>
- Walsh, J. C., Connors, K., Hertz, E., Kehoe, L., Martin, T. G., Connors, B., ... Reynolds, J. D. (2020). *BC Central Coast: Prioritizing strategies for Pacific salmon recovery and persistence.* Vancouver, BC: Report produced for the Central Coast Indigenous Resource Alliance. Retrieved from https://salmonwatersheds.ca/library/lib_454/
- Walters, C., English, K., Korman, J., & Hilborn, R. (2019). The managed decline of British Columbia's commercial salmon fishery. *Marine Policy*, 101(March), 25–32. <https://doi.org/10.1016/j.marpol.2018.12.014>
- Waples, R. S. (1991). Pacific salmon, *Oncorhynchus* spp., and the definition of "species" under the Endangered Species Act. *Marine Fisheries Review*, 53(3), 11–22.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Walsh JC, Connors K, Hertz E, et al. Prioritizing conservation actions for Pacific salmon in Canada. *J Appl Ecol.* 2020;00:1–12. <https://doi.org/10.1111/1365-2664.13646>