

Impacts of snow on soil temperature observed across the circumpolar north

Yu Zhang, Artem B Sherstiukov, Budong Qian, Steven V Kokelj, & Trevor C. Lantz

2018

Faculty of Social Science

Faculty Publications

© Zhang, Sherstiukov, Qian, Kokelj, & Lantz. This is an open access article distributed under the terms of the Creative Commons CC BY 3.0 License: <http://creativecommons.org/licenses/by/3.0/>.

Original citation:

Zhang, Y., Sherstiukov, A. B., Qian, B., Kokelj, S. V., & Lantz, T. C. (2018). Impacts of snow on soil temperature observed across the circumpolar north. *Environmental Research Letters*, 13(4), 044012. <https://doi.org/10.1088/1748-9326/aab1e7>

Downloaded from UVicSpace Research & Learning Repository

dspace.library.uvic.ca



University
of Victoria

Libraries

Environmental Research Letters



LETTER

Impacts of snow on soil temperature observed across the circumpolar north

OPEN ACCESS

RECEIVED

9 November 2017

REVISED

7 February 2018

ACCEPTED FOR PUBLICATION

23 February 2018

PUBLISHED

3 April 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Yu Zhang^{1,6} , Artem B Sherstiukov², Budong Qian³, Steven V Kokelj⁴ and Trevor C Lantz⁵

¹ Canada Centre for Remote Sensing, Canada Centre for Mapping and Earth Observation, Natural Resources Canada, 560 Rochester Street, Ottawa, Ontario, K1A 0E4, Canada

² All-Russian Research Institute of Hydrometeorological Information-World Data Center, 6 Korolyov Street, Obninsk, 249035, Russia

³ Ottawa Research and Development Centre, Science and Technology Branch, Agriculture and Agri-Food Canada, 960 Carling Avenue, Ottawa, Ontario, K1A 0C6, Canada

⁴ Northwest Territories Geological Survey, PO Box 1320, 4601 B 52 Avenue, Yellowknife, Northwest Territories, X1A 2L9, Canada

⁵ School of Environmental Studies, University of Victoria, 3800 Finnerty Road, Victoria, British Columbia, V8P 5C2, Canada

⁶ Author to whom any correspondence should be addressed.

E-mail: yu.zhang@canada.ca

Keywords: soil temperature, soil-air temperature difference, snow, northern high latitudes

Abstract

Climate warming has significant impacts on permafrost, infrastructure and soil organic carbon at the northern high latitudes. These impacts are mainly driven by changes in soil temperature (T_S). Snow insulation can cause significant differences between T_S and air temperature (T_A), and our understanding about this effect through space and time is currently limited. In this study, we compiled soil and air temperature observations (measured at about 0.2 m depth and 2 m height, respectively) at 588 sites from climate stations and boreholes across the northern high latitudes. Analysis of this circumpolar dataset demonstrates the large offset between mean T_S and T_A in the low arctic and northern boreal regions. The offset decreases both northward and southward due to changes in snow conditions. Correlation analysis shows that the coupling between annual T_S and T_A is weaker, and the response of annual T_S to changes in T_A is smaller in boreal regions than in the arctic and the northern temperate regions. Consequently, the inter-annual variation and the increasing trends of annual T_S are smaller than that of T_A in boreal regions. The systematic and significant differences in the relationship between T_S and T_A across the circumpolar north is important for understanding and assessing the impacts of climate change and for reconstruction of historical climate based on ground temperature profiles for the northern high latitudes.

1. Introduction

Air temperatures (T_A) across northern high latitudes are increasing twice as fast as the global average (IPCC 2013), which could have significant impacts on permafrost (Chadburn *et al* 2017), infrastructure (Nelson *et al* 2001), and the large stock of organic carbon stored in frozen soils (Schuur *et al* 2015). These impacts are mainly driven by changes in soil temperature (T_S). Thus, understanding the response of T_S to T_A increase is critical to predict the magnitude of these impacts. Seasonal snow cover can cause significant differences between T_A and T_S ($dT_{SA} = T_S - T_A$), known as surface offset, through thermal insulation and reflection of solar radiation (Smith and Riseborough 2002, Zhang 2005). Since

snow conditions are highly variable, observations and modelling studies show that annual dT_{SA} differs not only spatially (Zhang 2005, Morse *et al* 2012, Palmer *et al* 2012, Throop *et al* 2012), but also varies with time (Zhang *et al* 2001, Stieglitz *et al* 2003, Zhang *et al* 2005, Isard *et al* 2007, Osterkamp 2007, Romanovsky *et al* 2007, Woodbury *et al* 2009, Lawrence and Slater 2010, Qian *et al* 2011, Park *et al* 2014, Wang *et al* 2017). Previous observation-based investigations of dT_{SA} are limited to local or regional scales (Zhang *et al* 2001, Beltrami and Kellman 2003, Stieglitz *et al* 2003, Isard *et al* 2007, Osterkamp 2007, Romanovsky *et al* 2007, Sherstyukov 2008, Woodbury *et al* 2009, Qian *et al* 2011, Morse *et al* 2012, Park *et al* 2014, Streletskiy *et al* 2015, Wang *et al* 2017). Modelling studies can consider the entire circumpolar region but the results

differ widely due to model structure and input data (Lawrence and Slater 2010, Wang *et al* 2016). This uncertainty limits our capacity to assess the impacts of climate change on permafrost and northern ecosystems and affects reconstruction of historical climate change from deep ground (>10 m) temperature profiles (Beltrami and Kellman 2003, Mann and Schmidt 2003, Bartlett *et al* 2005).

In this study, we compiled observations at 588 sites from climate stations and boreholes across the northern high latitudes. Based on this large number of site observations, we analyzed the spatial and temporal variations of near surface T_S (measured at about 0.2 m below surface), near surface T_A (measured at about 2 m above surface), dT_{SA} and the relationships with snow conditions. We use the data to explore the following questions. (1) What is the general range of mean dT_{SA} , and are there any broad scale spatial patterns evident across the circumpolar north? (2) How does annual dT_{SA} change with climate warming, and is there any evidence that changes in dT_{SA} affected the response of T_S to T_A ? (3) How are the spatial and temporal variations in dT_{SA} related to snow conditions at the circumpolar scale?

2. Data and processing


The data used in this study include observations from climate stations and boreholes across the northern high latitudes. We only included sites with latitudes greater than 45 °N. Daily data from Russian meteorological stations were obtained from the All-Russian Research Institute of Hydrometeorological Information-World Data Centre (Sherstiukov 2012a). Daily data from Canadian climate stations were provided by Environment and Climate Change Canada. Borehole ground temperature and T_A data were obtained from Global Terrestrial Network for Permafrost (GTN-P) and the literature.

The Russian climate station dataset includes 458 stations for T_S , 619 stations for snow depth, 599 stations for T_A and precipitation. In combination, 264 stations have measurements of T_S , T_A , snow depth, and precipitation. A detailed description of the T_S dataset is provided in Sherstiukov (2012a). T_S was measured at 12 depths (0.02, 0.05, 0.1, 0.15, 0.2, 0.4, 0.6, 0.8, 1.2, 1.6, 2.4, and 3.2 m). The records are sparse at the top four depths. Therefore, we used observations at 0.2 m as the near surface T_S for the analysis. The observations were mainly from 1985–2011, and a third to a quarter of the stations also had observations from 1963–1976 and 1984. Sherstiukov (2012b) conducted a series of quality checks on the T_S dataset. Each record was given a quality flag but no corrections. In this study, we corrected the obvious errors based on the quality flags but strove to avoid over-correction. Obvious errors corrected included sudden sign changes and significant value changes.

We performed the correction by comparing the flagged records with the records on the preceding and the following days. On average, about one to two records of T_S at 0.2 m depth were corrected for each station.

In Canada, daily T_S was measured at about 80 stations across the country by Environment and Climate Change Canada. Most of these stations are in the south of 60 °N. T_S was measured at depths of 0.05, 0.1, 0.2, 0.5, 1.0, and 1.5 m during various periods between 1958 and 2008, mainly 1964–1999. T_S was recorded twice daily, morning (0800 h) and afternoon (1500 h). We only used morning observations in the analysis as there were more missing data in the afternoon observations (Qian *et al* 2011). In total, 70 stations have at least one year of complete observations for both T_A and T_S . Among these, 69 stations also have at least one year of snow depth observations.

The Global Terrestrial Network for Permafrost provides a comprehensive database for permafrost monitoring parameters, including ground temperatures at various depths, T_A and active-layer thickness. We downloaded (performed in March 2017) data from sites with at least one year of complete observations of T_A and/or ground temperatures at about 0.2 m depth. Some boreholes are near climate stations, so we estimated T_A for these boreholes from the nearby climate station data if T_A was not measured at the site. In total we compiled 145 boreholes with at least one year of complete observations of T_A and ground temperatures at about 0.2 m depth (101 boreholes in Russia, 29 boreholes in Alaska, 15 boreholes in Canada). We also included 109 sites from recently published papers: 53 sites across the Mackenzie Valley Corridor in northwestern Canada (Duchesne *et al* 2014, Wolfe *et al* 2010); one site in Peel Plateau in northwestern Canada (averaged for two spruce forest observation sites) (O'Neill *et al* 2015); two sites for alluvial and uplands in the Kendall Island Bird Sanctuary in the outer Mackenzie Delta (averaged for seven and eight observation sites, respectively) (Morse *et al* 2012); three sites for black spruce forest, white spruce forest and open black spruce peatland, respectively, in the Great Slave Region (each site was averaged from three nearby observation sites) (Morse *et al* 2016); three sites for bog, fen and palsa, respectively, in the southern Hudson Bay Lowlands (each site was averaged from two nearby observation sites) (Ou *et al* 2016); 24 sites in Labrador and Quebec in eastern Canada (measurements at barren and rock sites were excluded in boreal regions and the sites very close to one another were averaged as one site) (Way and Lewkowicz 2018); 11 sites across permafrost regions in Canada (Throop *et al* 2012); one site in the Lena River Delta, Russia (Boike *et al* 2013); and 11 sites in southern Norway (Farbrot *et al* 2011). Together with observations at climate stations, we compiled 588 sites with at least one year of complete observations of both T_A and T_S . The longest dataset available is 38 years. There are 442 sites with at least five years of complete observations



although the years may not be continuous. Figure 1(a) shows the distribution of the sites and the mean T_A .

Using daily snow depth observations at climate stations, we calculated annual snow cover duration, winter mean snow depth (last December, January and February) and annual mean snow depth (sum of daily snow depths divided by the total number of days in a year). In the text, annual T_A or annual mean T_A (or other variables) is the average of a year at a site, and mean T_A or mean annual T_A (or other variables) is the average of all the years of observations at a site. Annual averages are based on calendar year except for the correlation analysis of annual snow conditions described in the last two paragraphs in section 3.4, where a year is defined from October 1st to September 30th in the following year. Correlation analysis is based on Pearson momentum correlation coefficients (R). Simple linear regressions were calculated based on least squares to estimate temporal trends and linear relations between two variables. We also calculated mean absolute deviation (MAD) to represent inter-annual variations for annual T_A and T_S . These statistics were calculated for sites with at least five years of complete observations. We used a marginal significance level (one-sided $p < 0.1$) in trends and correlation analysis to maximize the number of sites for spatial coverage. The general patterns are similar when we limited to sites with longer observations and higher statistical significance levels.

3. Results


3.1. Spatial distribution of mean dT_{SA}

Mean T_A decreases from south to north (figure 1(a)). While mean T_S shares a similar pattern, it exhibits

a greater variation than that of T_A due to local variations in snow, soil and vegetation conditions. Figure 1(b) shows the spatial distribution of mean dT_{SA} . Mean dT_{SA} ranges from about 0.0°C–12.9°C, with an average of 4.4°C. Mean dT_{SA} shows a non-linear relationship with mean T_A : It increases from less than 2°C to up to 12.9°C (with an average of 7.3°C) until mean T_A reaches about -10°C , then it decreases to about 2°C until mean T_A is about 7°C , after which it has no obvious trend (figure 2(a)). Mean dT_{SA} is greatest in the low arctic and northern boreal regions. It decreases both northward and southward (figures 1(b) and 2). Because of these different patterns, the spatial gradients of mean T_S and T_A vary among the different biomes. The north-south gradient of mean T_S is about one and a half times of the gradient of mean T_A in the arctic, but only about two thirds of that of mean T_A in boreal regions on average. In northern temperate regions, the spatial gradients of mean T_S and T_A are similar, especially in southern areas.

3.2. Temporal variations of annual T_S and dT_{SA} with T_A

To investigate the response of T_S to changes in T_A , we calculated simple linear regressions and Pearson correlation coefficients between annual T_S and annual T_A , and between annual dT_{SA} and annual T_A based on the time series data for each site (table 1). Since T_A is the primary driver of the variations in T_S , annual T_S is positively and significantly ($p < 0.1$) correlated with annual T_A for 83% of the sites. However, the relationship between annual T_S and T_A varies with biomes. The R -values tend to be lower in boreal regions than in the arctic and the northern temperate regions (figure 3(a) and table 1). Annual dT_{SA} is significantly




($p < 0.1$) but negatively correlated with T_A for 83% of the sites, and the correlation tends to be stronger (closer to -1) in boreal regions than in the arctic and the northern temperate regions (figure 3(b) and table 1). This result indicates a weaker coupling between annual T_S and T_A associated with a stronger correlation between annual dT_{SA} and T_A in boreal regions than in the arctic and the northern temperate regions.

Slope coefficients (K) of linear regressions between annual T_S and T_A are less than 1.0 for almost all the sites (only 11 sites are exceptions). The values of K in boreal regions are smaller than in the arctic and the northern temperate regions (figure 3(c), table 1). Conversely, the values of K between annual dT_{SA} and T_A are negative, and they are smaller (closer to -1) in boreal regions than in the arctic and the northern temperate regions (figure 3(d), table 1). This analysis suggests that annual T_S is less responsive to changes in annual T_A in boreal regions than in the arctic and the northern temperate regions. However, annual dT_{SA} is more sensitive to changes in T_A in boreal regions than in other regions.

3.3. Impacts of dT_{SA} on the inter-annual variations and long-term trends of annual T_S

A negative correlation between annual dT_{SA} and T_A or a smaller response of annual T_S to the changes in annual T_A has two implications. First, the inter-annual fluctuation of T_S will be smaller than that of T_A . Second, if there is a long-term increasing trend in annual T_A , the increase of annual T_S will be less than that of T_A . To confirm these patterns, we calculated MAD and linear temporal trends for annual T_S and T_A based on the time series data for each site with at least five years of observations. The ratio between MAD of annual T_S and MAD of annual T_A increases linearly with the K between annual T_S and T_A and with the R between annual dT_{SA} and T_A (figures 4(a) and (b)). The ratio between the trend of annual T_S and the trend of annual T_A also increases linearly with the K between annual T_S and T_A and with the R between annual dT_{SA} and T_A (figures 4(c) and (d)). This result indicates that the inter-annual variation and the trend of T_S are closely related to the response of T_S to changes in annual T_A and the correlation between dT_{SA} and T_A . When the response of annual T_S to



change in T_A is small (small K -values), the inter-annual variation and the trend of T_S are smaller than that of T_A . Similarly, when annual dT_{SA} and T_A are strongly correlated, the inter-annual variation and the trend of T_S are smaller than that of T_A .

For different biomes, the ratio between MAD of annual T_S and MAD of annual T_A tends to be smaller in boreal regions than in the arctic and the northern temperate regions (table 2). Similarly, the ratio between the trend of annual T_S and the trend of annual T_A tends to be smaller in boreal regions than in the arctic and the northern temperate regions (table 2). These results are consistent with the weaker coupling between annual T_S and T_A (stronger correlation between annual dT_{SA} and T_A) and smaller K -values between annual T_S and T_A in the boreal than in the arctic and the northern temperate regions (table 1).

3.4. The causes of the spatial and temporal differences in dT_{SA}

Across the northern high latitudes, mean annual dT_{SA} is closely correlated with mean winter dT_{SA} ($R = 0.894$, $N = 554$) but the correlation with mean summer dT_{SA} is poor and negative ($R = -0.265$, $N = 554$ sites), indicating that mean annual dT_{SA} is mainly determined by winter conditions. Mean winter dT_{SA} is negatively correlated with mean winter T_A ($R = -0.740$, $N = 554$) while the correlation between mean summer dT_{SA} and mean summer T_A is poor ($R = 0.358$, $N = 554$).

Observations of T_A , T_S and snow depth at climate stations in Russia and Canada indicate that mean winter dT_{SA} is correlated with mean winter snow depth ($R = 0.661$, $N = 333$ stations, figure 5(a)). Mean dT_{SA} is also closely correlated with mean annual snow depth ($R = 0.678$, $N = 329$ stations, figure 5(b)).




Figure 4. The distributions of the ratios between the mean absolute deviation (MAD) of annual T_S and MAD of T_A with (a) the K between annual T_S and T_A and (b) the R between annual dT_S and T_A . And the distributions of the ratios between the trend of annual T_S and the trend of annual T_A with (c) the K between annual T_S and T_A and (d) the R between annual dT_S and T_A . Each circle represents one site with significant correlations ($p < 0.1$) for the K between annual T_S and T_A and for the trends of annual T_S and T_A . They are calculated based on the time series of data for each site with at least five years of observations.

Table 1. The correlation between annual T_S and annual T_A and between annual dT_{SA} and annual T_A for sites with at least five years of observations.

Biomes	The arctic	Boreal regions	Northern temperate	Whole area
Number of sites	35	302	105	442
Average observation years	10.1	20.6	22.1	20.1
Correlation between annual T_S and annual T_A				
Average (median) R	0.66 (0.70)	0.56 (0.61)	0.74 (0.80)	0.61 (0.67)
Number (and %) of sites with a significant ($p < 0.1$) correlation	26 (74.3%)	243 (80.5%)	98 (93.3%)	367 (83.0%)
Average (median) K when $p < 0.1$	0.68 (0.63)	0.46 (0.64)	0.63 (0.61)	0.52 (0.48)
Correlation between annual dT_{SA} and annual T_A				
Average (median) R	-0.45 (-0.45)	-0.70 (-0.77)	-0.59 (-0.69)	-0.66 (-0.74)
Number (and %) of sites with a significant ($p < 0.1$) correlation	17 (48.5%)	269 (89.1%)	81 (77.1%)	367 (83.0%)
Average (median) K when $p < 0.1$	-0.56 (-0.51)	-0.65 (-0.63)	-0.46 (-0.47)	-0.60 (-0.59)

Table 2. The ratio between MAD of annual T_S and MAD of annual T_A , and the ratio between the trend of annual T_S and the trend of annual T_A when the trends are significant ($p < 0.1$) for both T_S and T_A .

	The arctic	Boreal regions	Northern temperate	Whole area
The ratios between MAD of annual T_S and MAD of annual T_A				
Number of sites	35	302	105	442
Average observation years	10.1	20.6	22.1	20.1
Average (median) ratios	1.00 (0.93)	0.73 (0.66)	0.83 (0.75)	0.77 (0.70)
% of sites with the ratio < 1	57.1	84.8	81.0	81.7
The ratios between the trend of annual T_S and the trend of annual T_A				
Number (and %) of sites with significant ($p < 0.1$) trends	11 (31.4%)	120 (39.7%)	58 (55.2%)	189 (42.8%)
Average observation years	13.5	25.7	26.5	25.2
Average (median) ratios	1.00 (1.05)	0.75 (0.70)	0.91 (0.82)	0.81 (0.79)
% of the sites with the ratio < 1	36.4	76.7	67.2	71.4

Table 3. Correlation analyses among annual mean snow depth, snow cover duration, annual T_A and dT_{SA} for climate stations with at least five years of complete observations.

Biomes	The arctic	Boreal regions	Northern temperate	Whole area
Number of sites	8	213	79	300
Average observation years	12.9	17.6	16.7	17.2
Correlation between annual mean snow depth and annual snow cover duration				
Average (median) R	0.48 (0.60)	0.57 (0.61)	0.66 (0.68)	0.59 (0.63)
Number (and %) of sites with a significant ($p < 0.1$) correlation	5 (62.5%)	187 (87.8%)	75 (94.9%)	267 (89.0%)
Correlation between annual mean snow depth and annual mean T_A				
Average (median) R	0.07 (−0.02)	−0.25 (−0.30)	−0.48 (−0.46)	−0.30 (−0.34)
Number (and %) of sites with a significant ($p < 0.1$) correlation	2 (25.0%)	112 (52.6%)	55 (69.6%)	169 (56.3%)
Average (median) K when $p < 0.1$ ($cm/^\circ C$)	4.05 (4.05)	−1.10 (−1.37)	−1.08 (−0.86)	−1.03 (−1.23)
Correlation between annual snow cover duration and annual mean T_A				
Average (median) R	−0.16 (−0.13)	−0.40 (−0.43)	−0.55 (−0.62)	−0.44 (−0.48)
Number (and %) of sites with a significant ($p < 0.1$) correlation	1 (12.5%)	147 (69.0%)	66 (83.5%)	214 (71.3%)
Average (median) K when $p < 0.1$ ($d/^\circ C$)	−4.9 (−4.9)	−6.4 (−5.7)	−9.1 (−8.7)	−7.2 (−6.2)
Correlation between annual dT_{SA} and annual mean snow depth				
Average (median) R	0.52 (0.55)	0.53 (0.60)	0.54 (0.60)	0.53 (0.60)
Number (and %) of sites with a significant ($p < 0.1$) correlation	7 (87.5%)	164 (77.0%)	62 (78.5%)	233 (77.7%)
Average (median) K when $p < 0.1$ ($^\circ C/cm$)	0.093 (0.089)	0.183 (0.150)	0.187 (0.156)	0.182 (0.150)
Correlation between annual dT_{SA} and annual snow cover duration				
Average (median) R	0.15 (0.17)	0.39 (0.45)	0.43 (0.51)	0.40 (0.46)
Number (and %) of sites with a significant ($p < 0.1$) correlation	2 (25.0%)	144 (67.6%)	55 (69.6%)	201 (67.0%)
Average (median) K when $p < 0.1$ ($^\circ C/d$)	0.055 (0.055)	0.039 (0.035)	0.018 (0.018)	0.034 (0.034)

For stations with three to eight months of snow cover in a year, mean dT_{SA} increases linearly with mean snow cover duration ($R = 0.797$, $N = 295$ stations) (figure 5(c)). Mean annual snow depth tends to be low when mean T_A is very low or very high (figure 6(a)), and snow cover duration decreases with the increase in mean T_A (figure 6(b)). Thus, the large dT_{SA} in the low arctic and northern boreal regions is mainly due to the long duration of relatively thick snow cover in this region. In the high arctic, thin snow cover due to low snowfall, wind redistribution (Morse *et al* 2012, Palmer *et al* 2012) and development of wind slab (Derksen *et al* 2014) reduce the insulation effects on the ground. In the northern temperate regions, short duration of snow cover reduces annual dT_{SA} .

The year-to-year variation of annual dT_{SA} is also positively correlated with snow depth and snow cover duration for most of the stations (table 3), indicating that snow condition has significant impacts on the temporal variation of annual dT_{SA} . Annual mean snow depth and annual snow cover duration (from October 1st to September 30th in the following year) are also positively correlated to each other. Snow effects on the response of T_S to changes in T_A depend on the changes in snow conditions with T_A (due to snowfall and snowmelt) and the consequent changes in dT_{SA} . Annual snow depths and snow cover durations generally have negative correlations with annual T_A , especially for snow cover durations (table 3) (except for some stations in Siberia and one station in northeast Canada where the correlations are positive). However, the strength of the correlations vary with biomes. In the arctic, the correlations between snow conditions (depth and duration) and annual T_A are generally not statistically significant. The correlations are stronger (more negative) in the northern temperate than in

boreal regions. However, annual dT_{SA} is more sensitive (higher K-values) to changes in snow depth in boreal regions than in the northern temperate regions (table 3). This snow depth effect on dT_{SA} probably is the main reason for the reduced response of annual T_S to changes in T_A in boreal regions.

Since the periods of observation are short, the temporal trends of dT_{SA} and snow conditions are not significant for most of the stations. For stations with significant trends ($p < 0.1$) for both annual dT_{SA} and snow conditions, the trends of annual dT_{SA} are positively correlated with the trends of snow conditions (with winter mean snow depth: $R = 0.865$, $N = 40$ stations; with annual mean snow depth: $R = 0.585$, $N = 46$ stations; with annual snow cover duration: $R = 0.669$, $N = 42$ stations).

4. Discussion

The effects of snow on ground temperature (the surface offset) have been recognized in previous studies (Smith and Riseborough 2002, Zhang 2005, Palmer *et al* 2012). Several observation-based studies have assessed the differences between T_S and T_A and their temporal dynamics but only for limited sites and areas. Modelling studies can cover large areas, but their results vary widely due to model structure, resolutions and inputs (Lawrence and Slater 2010, Wang *et al* 2016). This study synthesizes a large number of observations across the northern high latitudes. The data show that there is a large offset between T_S and T_A in the low arctic and northern boreal regions across the circumpolar, but small offsets in the high arctic, southern boreal and northern temperate regions. In boreal regions, a thick and relatively long snow cover duration and




Figure 5. Scatter graphs (a) between mean winter dT_{SA} and mean winter snow depth, (b) between mean annual dT_{SA} and mean annual snow depth, and (c) between mean annual dT_{SA} and mean annual snow cover duration. Each circle represents one site (averaged for all the years with data available).

its strong impacts on annual dT_{SA} variation decouple T_S and T_A and dampen the response of T_S to climate warming (increase in T_A).

The results of this study agree qualitatively with previous observational analysis and modelling studies, which have shown that snow is the major factor differentiating mean T_S from T_A in permafrost regions (e.g. Smith and Riseborough 2002). The insulation effect of snow on annual T_S is at least twice that of the shading and cooling effects of vegetation in summer (Matyshak *et al* 2015). When snow cover is thin or absent in winter, the annual mean dT_{SA} can be very small (Lacelle *et al* 2016). Based on observations at




Figure 6. The distribution of mean T_A with (a) mean annual snow depth, and (b) mean annual snow cover duration. Each circle represents one site (averaged for all the years with data available). The data are from 329 climate stations with T_S observations in Russia and Canada.

ten sites across Canadian permafrost regions, Throop *et al* (2012) indicate that dT_{SA} is the smallest in the arctic sites, and the largest in the boreal of the Mackenzie Valley. A modelling study by Lawrence and Slater (2010) shows that the warming of T_S is less than that of T_A for most of the northern high latitudes during 1950–2100. The ratio of the warming between T_S and T_A is smaller in the boreal than in the arctic and the northern temperate regions (figure 1(c) in Lawrence and Slater 2010).

Climate stations account for about half of the sites reported in this study. They are usually located in flat areas with short vegetation, which can be very different from natural conditions, especially in forest regions. In boreal regions, vegetation is denser and higher than in the arctic, organic layers and mosses are usually deep and wet, and ground ice is common in near surface permafrost. Thus, T_S in natural boreal areas may be even lower and less responsive to changes in T_A than measured at climate stations. On the other hand, arctic regions have low and sparse vegetation, and organic and mineral deposits are usually thin. Thus, the response of T_S to changes in T_A is likely to be direct and rapid (Throop *et al* 2012).

The results of this study are useful for understanding the current distribution of permafrost and its

response to climate warming, for assessing the impacts of climate change on ecosystems and soil organic carbon, for improving models about snow effects on T_S , and for reconstruction of historical climate based on ground temperature profiles at the northern high latitudes. For example, figure 2 and table 2 indicate that the increase of annual T_S with time is smaller than that of T_A in boreal regions. Thus, the warming and degradation of permafrost in boreal regions will be less than the estimates directly from T_A . In contrast, near surface T_S in arctic regions is highly responsive to changes in T_A , therefore these regions can be very sensitive to climate warming. On the other hand, reconstructions of historical climate from deep ground temperature profiles would likely underestimate the changes in T_A in boreal regions. For instance, the estimated warming of near-surface T_S during 1950–2000 in Canada (Beltrami *et al* 2003) is smaller than that of T_A observed at climate stations, especially in boreal regions (Vincent *et al* 2015, Zhang *et al* 2000).

Our results show the insulative effects of snow on T_S at the circumpolar scale. The data also show strong site variations due to local and regional ground and climate conditions. Significant changes in snowfall could directly affect snow depth, thus altering dT_{SA} and the response of T_S to T_A . Changes in soil and vegetation conditions due to fire disturbances and gradual responses to climate warming could also affect T_S and its response to T_A .

5. Conclusions

This study compiled T_S and T_A observations from 588 climate stations and boreholes across the northern high latitudes. This broad observational data coverage analyzed in this paper shows a large offset between mean T_S and T_A in the low arctic and northern boreal regions across the circumpolar north. The offset decreases both northward and southward. Thus, the north-south gradient of mean T_S is about one and a half times of that of T_A in the arctic, but only about two-thirds of T_A in boreal regions. Further south, the spatial gradients of mean T_S and T_A are similar. This pattern is closely related to snow conditions. Thick and persistent snow cover and its strong impacts on annual dT_{SA} in boreal regions weaken the coupling between annual T_S and T_A , and the response of annual T_S to variation in annual T_A is smaller in the boreal than in the arctic and the northern temperate regions. Consequently, the inter-annual variation and the trends of T_S are smaller than that of T_A in boreal regions. This systematic and significant differences in the relationship between T_S and T_A across the circumpolar north is important for understanding and assessing the impacts of climate change and for reconstruction of historical climate based on ground temperature profiles for the northern high latitudes.

Acknowledgements

The authors thank Steve Wolfe and Peter Morse for stimulating discussion and their careful review of the earlier versions of the paper. This study was funded by Polar Knowledge Canada Science and Technology Program (project 186). This study also contributes to a project affiliated to the Arctic Boreal Vulnerability Experiment (ABOVE), a NASA Terrestrial Ecology program.

ORCID iDs

Yu Zhang  <https://orcid.org/0000-0001-5377-8017>

References

- Bartlett M G, Chapman D S and Harris R N 2005 Snow effect on North American ground temperatures, 1950–2002 *J. Geophys. Res.* **110** F03008
- Beltrami H, Gosselin C and Mareschal J C 2003 Ground surface temperatures in Canada: spatial and temporal variability *Geophys. Res. Lett.* **30** 1499
- Beltrami H and Kellman L 2003 An examination of short- and long-term air-ground temperature coupling *Glob. Planet. Change* **38** 291–303
- Boike J *et al* 2013 Baseline characteristics of climate, permafrost and land cover from a new permafrost observatory in the Lena River Delta, Siberia 1998–2011 *Biogeosciences* **10** 2105–28
- Brandt J P 2009 The extent of the North American boreal zone *Environ. Rev.* **17** 101–61
- Brown J, Ferrians O, Heginbottom J A and Melnikov E 2002 *Circum-Arctic Map of Permafrost and Ground-Ice Conditions, Version 2* (Boulder, CO: National Snow and Ice Data Center)
- Chadburn S E, Burke E J, Cox P M, Friedlingstein P, Hugelius G and Westermann S 2017 An observation-based constraint on permafrost loss as a function of global warming *Nat. Clim. Change* **7** 340–44
- Derksen C, Lemmetyinen J, Toose P, Silis A, Pulliainen J and Sturm M 2014 Physical properties of Arctic versus subarctic snow: implications for high latitude passive microwave snow water equivalent retrievals *J. Geophys. Res. Atmos.* **119** 7254–70
- Duchesne C, Riseborough D W and Smith S L 2014 Air and near surface ground temperatures, indices and summary statistics from 1993–2012 for the Mackenzie Valley Corridor, Northwest Territories *Geological Survey of Canada Open File* vol 7392 revised (Ottawa: Natural Resources Canada Canada) p 85
- Farbrot H, Hipp T F, Etzelmüller B, Isaksen K, Ódegård R S, Schuler T V and Humlum O 2011 Air and ground temperature variations observed along elevation and continentality gradients in southern Norway *Permafrost Periglac. Process.* **22** 343–60
- IPCC 2013 Climate change 2013: the physical science basis *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker (Cambridge: Cambridge University Press) p 1535
- Isard S A, Schaetzl R J and Andresen J A 2007 Soils cool as climate warms in the Great Lakes region: 1951–2000 *Ann. Assoc. Am. Geogr.* **97** 467–76
- Lacelle D, Lapalme C, Davila A F, Pollard W, Marinova M, Heldmann J and McKay C P 2016 Solar radiation and air and ground temperature relations in the cold and hyper-arid Quartermain Mountains, McMurdo Dry Valleys of Antarctica *Permafrost Periglac. Process.* **27** 163–76

- Lawrence D M and Slater A G 2010 The contribution of snow condition trends to future ground climate *Clim. Dyn.* **34** 969–81
- Mann M E and Schmidt G A 2003 Ground vs. surface air temperature trends: implications for borehole surface temperature reconstructions *Geophys. Res. Lett.* **30** 1607
- Matyshak G V, Goncharova O Y, Moskalenko N G, Walker D A, Epstein H E and Shur Y 2015 Contrasting soil thermal regimes in the forest-tundra transition near Nadym, West Siberia, Russia *Permafrost Periglac. Process.* **28** 108–18
- Morse P D, Burn C R and Kokelj S V 2012 Influence of snow on near-surface ground temperatures in upland and alluvial environments of the outer Mackenzie Delta, Northwest Territories *Can. J. Earth Sci.* **49** 895–913
- Morse P D, Wolfe S A, Kokelj S V and Gaanderse A J R 2016 The occurrence and thermal disequilibrium state of permafrost in forest ecotopes of the Great Slave Region, Northwest Territories, Canada *Permafrost Periglac. Process.* **27** 145–62
- Nelson F E, Anisimov O A and Shiklomanov N I 2001 Subsidence risk from thawing permafrost *Nature* **410** 889–90
- O'Neill H B, Burn C R, Kokelj S V and Lantz T C 2015 'Warm' tundra: atmospheric and near-surface ground temperature inversions across an alpine treeline in continuous permafrost, western Arctic, Canada *Permafrost Periglac. Process.* **26** 103–18
- Osterkamp T 2007 Causes of warming and thawing of permafrost in Alaska *EOS* **88** 522–3
- Ou C, Leblon B, Zhang Y, LaRocquea A, Webster K and McLaughlin J 2016 Modelling and mapping permafrost at high spatial resolution using Landsat and Radarsat images in northern Ontario, Canada: part 1—model calibration *Int. J. Remote Sens.* **37** 2727–50
- Palmer M J, Burn C R and Kokelj S V 2012 Factors influencing permafrost temperatures across tree line in the uplands east of the Mackenzie Delta, 2004–2010 *Can. J. Earth Sci.* **49** 877–94
- Park H, Sherstiukov A B, Fedorov A N, Polyakov I V and Walsh J E 2014 An observation-based assessment of the influences of air temperature and snow depth on soil temperature in Russia *Environ. Res. Lett.* **9** 064026
- Potapov P, Hansen M C, Stehman S V, Loveland T R and Pittman K 2008 Combining MODIS and Landsat imagery to estimate and map boreal forest cover loss *Remote Sens. Environ.* **112** 3708–19
- Qian B, Gregorich E G, Gameda S, Hopkins D W and Wang X L 2011 Observed soil temperature trends associated with climate change in Canada *J. Geophys. Res.* **116** D02106
- Romanovsky V E, Sazonova T S, Balobaev V T, Shender N I and Sergueev D O 2007 Past and recent changes in air and permafrost temperatures in eastern Siberia *Glob. Planet. Change* **56** 399–413
- Schuur E A G *et al* 2015 Climate change and the permafrost carbon feedback *Nature* **520** 171–9
- Sherstyukov A B 2008 Correlation of soil temperature with air temperature and snow cover depth in Russia (in Russian) *Earth Cryosphere* **12** 79–87
- Sherstiukov A 2012a Data set of daily soil temperature at depths to 320 cm at meteorological stations of Russian RIHMI-WDC **176** 233–56
- Sherstiukov A 2012b Statistical quality control of a daily soil temperature dataset *RIHMI-WDC* **176** 224–32
- Smith M W and Riseborough D W 2002 Climate and the limits of permafrost: a zonal analysis *Permafrost Periglac. Process.* **13** 1–15
- Stieglitz M, Déry S J, Romanovsky V E and Osterkamp T E 2003 The role of snow cover in the warming of arctic permafrost *Geophys. Res. Lett.* **30** 1721
- Streletskiy D A, Sherstiukov A B, Frauenfeld O W and Nelson F E 2015 Changes in the 1963–2013 shallow ground thermal regime in Russian permafrost regions *Environ. Res. Lett.* **10** 125005
- Throop J, Lewkowicz A G and Smith S L 2012 Climate and ground temperature relations at sites across the continuous and discontinuous permafrost zones, northern Canada *Can. J. Earth Sci.* **49** 865–76
- Vincent L A, Zhang X, Brown R D, Feng Y, Mekis E, Milewska E J, Wan H and Wang X L 2015 Observed trends in Canada's climate and influence of low-frequency variability modes *J. Clim.* **28** 4545–60
- Wang W *et al* 2016 Evaluation of air–soil temperature relationships simulated by land surface models during winter across the permafrost region *Cryosphere* **10** 1721–37
- Wang Y, Hu Z Z and Yan F 2017 Spatiotemporal variations of differences between surface air and ground temperatures in China *J. Geophys. Res. Atmos.* **122** 7990–9
- Way R G and Lewkowicz A G 2018 Environmental controls on ground temperature and permafrost in Labrador, northeast Canada *Permafrost Periglac. Process.* **2018** 1–13
- Wolfe S A, Smith S L, Chartrand J, Kokelj S, Palmer M and Stevens C W 2010 Geotechnical database and descriptions of permafrost monitoring sites established 2006–10 in the northern Mackenzie Corridor, Northwest Territories *Geol. Survey Canada* vol 6677 (Ottawa: Natural Resources Canada) p 76 1 CD-ROM (<https://doi.org/10.4095/287167>)
- Woodbury A D, Bhuiyan A K M H, Hanesiak J and Akinremi O O 2009 Observations of northern latitude ground-surface and surface-air temperatures *Geophys. Res. Lett.* **36** L07703
- Zhang T 2005 Influence of the seasonal snow cover on the ground thermal regime: an overview *Rev. Geophys.* **43** RG4002
- Zhang T, Barry R G, Gilichinsky D, Bykhovets S S, Sorokovikov V A and Ye J P 2001 An amplified signal of climatic change in soil temperatures during the last century at Irkutsk, Russia *Clim. Change* **49** 41–76
- Zhang X, Vincent L A, Hogg W D and Niitsoo A 2000 Temperature and precipitation trends in Canada during the 20th century *Atmos. Ocean* **38** 395–429
- Zhang Y, Chen W, Smith S L, Riseborough D W and Cihlar J 2005 Soil temperature in Canada during the 20th century: complex responses to atmospheric climate change *J. Geophys. Res.* **110** D03112