

---

Faculty of Science

Faculty Publications

---

Some Generalized Steffensen's Inequalities via a New Identity for Local Fractional Integrals

Tuba Tunç, Mehmet Zeki Sarikaya, and H. M. Srivastava

2017

© 2017 Srivastava et al. This is an open access article distributed under the terms of the Creative Commons Attribution License. <http://creativecommons.org/licenses/by/4.0>

This article was originally published at:

<http://www.etamaths.com/index.php/ijaa/article/view/879>

---

Citation for this paper:

Tunç, T.; Sarikaya, M. Z.; & Srivastava, H. M. (2017). Some generalized Steffensen's inequalities via a new identity for local fractional integrals. *International Journal of Analysis and Applications*, 13(1), 98-107.

## SOME GENERALIZED STEFFENSEN'S INEQUALITIES VIA A NEW IDENTITY FOR LOCAL FRACTIONAL INTEGRALS

TUBA TUNÇ<sup>1,\*</sup>, MEHMET ZEKI SARIKAYA<sup>1</sup> AND H. M. SRIVASTAVA<sup>2,3</sup>

ABSTRACT. In this study, we first give an identity for local fractional integrals. We then make use of this identity in order to derive several generalizations of the celebrated Steffensen's inequality associated with local fractional integrals. Relevant connections of the results presented in this paper with those that were proven in earlier works are also pointed out.

### 1. INTRODUCTION

As long ago as 1919, Steffensen [25] established the following result which is known in the literature as Steffensen's inequality.

**Theorem 1.1.** *Let  $a$  and  $b$  be real numbers such that  $a < b$ . Also let  $f, g : [a, b] \rightarrow \mathbb{R}$  be integrable functions such that  $f$  is nonincreasing and, for every  $x \in [a, b]$ ,  $0 \leq g(x) \leq 1$ . Then*

$$\int_{b-\lambda}^b f(x)dx \leq \int_a^b f(x)g(x)dx \leq \int_a^{a+\lambda} f(x)dx, \quad (1.1)$$

where

$$\lambda = \int_a^b g(x)dx.$$

Steffensen's inequality (1.1) happens to be the most basic inequality which deals with the comparison between integrals over a whole interval  $[a, b]$  and integrals over a subset of  $[a, b]$ . In fact, the inequality (1.1) has attracted considerable attention and interest from mathematicians and researchers. In this connection, the interested reader is referred to a number of works (see, for example, [3], [4], [7]-[10], [13], [14], [17] and [26]) for various related integral inequalities.

Recently, Wu and Srivastava [27] proved the following inequality which is a weighted version of the inequality (1.1).

**Theorem 1.2.** *Let  $f$ ,  $g$  and  $h$  be integrable functions defined on  $[a, b]$  with  $f$  nonincreasing. Also let  $0 \leq g(x) \leq h(x)$  for all  $x \in [a, b]$ . Then the following inequalities hold true:*

$$\begin{aligned} \int_{b-\lambda}^b f(x)h(x)dx &\leq \int_{b-\lambda}^b (f(x)h(x) - [f(x) - f(b - \lambda)][h(x) - g(x)]) dx \\ &\leq \int_a^b f(x)g(x)dx \\ &\leq \int_a^{a+\lambda} (f(x)h(x) - [f(x) - f(a + \lambda)][h(x) - g(x)]) dx \\ &\leq \int_a^{a+\lambda} f(x)h(x)dx, \end{aligned}$$

where  $\lambda$  is given by

$$\int_a^{a+\lambda} h(x)dx = \int_a^b g(x)dx = \int_{b-\lambda}^b h(x)dx.$$

---

Received 12<sup>th</sup> August, 2016; accepted 7<sup>th</sup> October, 2016; published 3<sup>rd</sup> January, 2017.

2010 *Mathematics Subject Classification.* 26D15, 26D99, 26A33.

*Key words and phrases.* Steffensen's inequality; local fractional integrals; fractal space; Fink's identity.

©2017 Authors retain the copyrights of  
 their papers, and all open access articles are distributed under the terms of the Creative Commons Attribution License.

We recall the following identity given by Fink [1]:

$$\begin{aligned} & \frac{1}{n} \left( f(x) + \sum_{k=1}^{n-1} F_k(x) \right) - \frac{1}{b-a} \int_a^b f(t) dt \\ &= \frac{1}{n!(b-a)} \int_a^b (x-t)^{n-1} \kappa(t,x) f^{(n)}(t) dt, \end{aligned} \quad (1.2)$$

where

$$F_k(x) = \left( \frac{n-k}{k!} \right) \left( \frac{f^{(k-1)}(a)(x-a)^k - f^{(k-1)}(b)(x-b)^k}{b-a} \right) \quad (1.3)$$

and

$$\kappa(t,x) = \begin{cases} t-a & (a \leq t \leq x \leq b) \\ t-b & (a \leq x < t \leq b) \end{cases}$$

Pečarić *et al.* [15] gave generalizations of Steffensen's inequality (1.1) via Fink's identity (1.2). Subsequently, Pečarić *et al.* [16] derived several new identities related to various generalizations of Steffensen's inequality (1.1).

## 2. DEFINITIONS, NOTATIONS AND PRELIMINARIES

The concepts of fractional calculus [8] and local fractional calculus (also called fractal calculus) (see, for details, [28] and [32]) are becoming increasingly useful in a wide variety of problems in mathematical, physical and engineering sciences (see, for example, the recent works [29] to [39]).

With a view to introducing the definition of the local fractional derivative and the local fractional integral, we need the following notations and preliminaries (see [28] and [32]).

For  $0 < \alpha \leq 1$ , we have the following  $\alpha$ -type sets of elements:

$\mathbb{Z}^\alpha$  : The  $\alpha$ -type set of integers defined by

$$\mathbb{Z}^\alpha := \{0^\alpha, \pm 1^\alpha, \pm 2^\alpha, \dots, \pm n^\alpha, \dots\}.$$

$\mathbb{Q}^\alpha$  : The  $\alpha$ -type set of the rational numbers defined by

$$\mathbb{Q}^\alpha := \left\{ m^\alpha : m^\alpha = \left( \frac{p}{q} \right)^\alpha \quad (p, q \in \mathbb{Z}; q \neq 0) \right\}.$$

$\mathbb{J}^\alpha$  : The  $\alpha$ -type set of the irrational numbers defined by

$$\mathbb{J}^\alpha := \left\{ m^\alpha : m^\alpha \neq \left( \frac{p}{q} \right)^\alpha \quad (p, q \in \mathbb{Z}; q \neq 0) \right\}.$$

$\mathbb{R}^\alpha$  : The  $\alpha$ -type set of the real line numbers defined by

$$\mathbb{R}^\alpha := \mathbb{Q}^\alpha \cup \mathbb{J}^\alpha.$$

**Proposition.** *Let  $a^\alpha, b^\alpha$  and  $c^\alpha$  belong to the set  $\mathbb{R}^\alpha$  of real line numbers. Then*

- (1)  $a^\alpha + b^\alpha$  and  $a^\alpha b^\alpha$  belong to the set  $\mathbb{R}^\alpha$ ;
- (2)  $a^\alpha + b^\alpha = b^\alpha + a^\alpha = (a+b)^\alpha = (b+a)^\alpha$ ;
- (3)  $a^\alpha + (b^\alpha + c^\alpha) = (a+b)^\alpha + c^\alpha$ ;
- (4)  $a^\alpha b^\alpha = b^\alpha a^\alpha = (ab)^\alpha = (ba)^\alpha$ ;
- (5)  $a^\alpha (b^\alpha c^\alpha) = (a^\alpha b^\alpha) c^\alpha$ ;
- (6)  $a^\alpha (b^\alpha + c^\alpha) = a^\alpha b^\alpha + a^\alpha c^\alpha$ ;
- (7)  $a^\alpha + 0^\alpha = 0^\alpha + a^\alpha = a^\alpha$  and  $a^\alpha 1^\alpha = 1^\alpha a^\alpha = a^\alpha$ .

The definitions of the local fractional derivative and the local fractional integral can now be given as follows.

**Definition 2.1.** (see [28] and [32]) *A non-differentiable function  $f : \mathbb{R} \rightarrow \mathbb{R}^\alpha$  ( $x \rightarrow f(x)$ ) is said to be local fractional continuous at  $x = x_0$  if, for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that*

$$|f(x) - f(x_0)| < \varepsilon^\alpha$$

holds true for

$$|x - x_0| < \delta \quad (\varepsilon, \delta \in \mathbb{R}).$$

If the function  $f(x)$  is local continuous on the interval  $(a, b)$ , we denote this property as follows:

$$f(x) \in C_\alpha(a, b).$$

**Definition 2.2.** (see [28] and [32]) The local fractional derivative of  $f(x)$  of order  $\alpha$  ( $0 < \alpha \leq 1$ ) at  $x = x_0$  is defined by

$$f^{(\alpha)}(x_0) = \left. \frac{d^\alpha f(x)}{dx^\alpha} \right|_{x=x_0} = \lim_{x \rightarrow x_0} \frac{\Delta^\alpha (f(x) - f(x_0))}{(x - x_0)^\alpha}, \quad (2.1)$$

where

$$\Delta^\alpha (f(x) - f(x_0)) \cong \Gamma(\alpha + 1)(f(x) - f(x_0)).$$

If there exists

$$f^{(k+1)\alpha}(x) = \overbrace{D_x^\alpha \cdots D_x^\alpha}^{k+1 \text{ times}} f(x) \quad (x \in I \subseteq \mathbb{R}),$$

then we write

$$f \in D_{(k+1)\alpha}(I) \quad (k \in \mathbb{N}_0 := \{0, 1, 2, \dots\} = \mathbb{N} \cup \{0\}).$$

**Definition 2.3.** (see [28] and [32]) Let  $f(x) \in C_\alpha[a, b]$ . Then the local fractional integral of  $f(x)$  of order  $\alpha$  ( $0 < \alpha \leq 1$ ) is defined by

$${}_a I_b^\alpha f(x) = \frac{1}{\Gamma(1 + \alpha)} \int_a^b f(t)(dt)^\alpha = \frac{1}{\Gamma(1 + \alpha)} \lim_{\Delta t \rightarrow 0} \sum_{j=0}^{N-1} f(t_j)(\Delta t_j)^\alpha \quad (2.2)$$

with

$$\Delta t_j = t_{j+1} - t_j \quad \text{and} \quad \Delta t = \max \{ \Delta t_1, \Delta t_2, \dots, \Delta t_{N-1} \},$$

where  $[t_j, t_{j+1}]$  ( $j = 0, 1, \dots, N-1$ ) and

$$a = t_0 < t_1 < \dots < t_{N-1} < t_N = b$$

is a partition of the interval  $[a, b]$ .

Clearly, we find from Definition 2.3 that

$${}_a I_b^\alpha f(x) = \begin{cases} 0 & (a = b) \\ -{}_b I_a^\alpha f(x) & (a < b). \end{cases} \quad (2.3)$$

If, for any  $x \in [a, b]$ , there exists  ${}_a I_x^\alpha f(x)$ , then we denote it simply as follows:

$$f(x) \in I_x^\alpha [a, b].$$

**Lemma 2.1.** (see [28] and [32]) It is asserted that

- (i)  $\frac{d^\alpha x^{k\alpha}}{dx^\alpha} = \frac{\Gamma(1 + k\alpha)}{\Gamma(1 + (k-1)\alpha)} x^{(k-1)\alpha}$ ;
- (ii)  $\frac{1}{\Gamma(1 + \alpha)} \int_a^b x^{k\alpha} (dx)^\alpha = \frac{\Gamma(1 + k\alpha)}{\Gamma(1 + (k+1)\alpha)} (b^{(k+1)\alpha} - a^{(k+1)\alpha}) \quad (k \in \mathbb{R})$ .

**Lemma 2.2.** (see [28] and [32]) Each of the following assertions holds true.

(1) Local fractional integration is anti-differentiation: Suppose that

$$f(x) = g^{(\alpha)}(x) \in C_\alpha [a, b].$$

Then

$${}_a I_b^\alpha f(x) = g(b) - g(a).$$

(2) Local fractional integration by parts: Suppose that

$$f(x), g(x) \in D_\alpha [a, b] \quad \text{and} \quad f^{(\alpha)}(x), g^{(\alpha)}(x) \in C_\alpha [a, b].$$

Then

$${}_a I_b^\alpha f(x)g^{(\alpha)}(x) = f(x)g(x)|_a^b - {}_a I_b^\alpha f^{(\alpha)}(x)g(x).$$

We next recall that Sarikaya *et al.* [23] proved the following generalized Steffensen inequality for local fractional integrals.

**Theorem 2.1.** Let  $f(x), g(x) \in I_x^\alpha [a, b]$  such that the function  $f(x)$  is non-increasing and  $0 \leq g(x) \leq 1$  on  $[a, b]$  ( $a < b$ ). Then

$${}_{b-\lambda}I_b^\alpha f(x) \leq {}_aI_b^\alpha f(x)g(x) \leq {}_aI_{a+\lambda}^\alpha f(x), \quad (2.4)$$

where

$$\lambda^\alpha = \Gamma(1 + \alpha) {}_aI_b^\alpha g(x).$$

Sarikaya *et al.* [23] also stated the following identities which we shall use in order to prove our main results in this paper.

$$\begin{aligned} & \frac{1}{\Gamma(\alpha + 1)} \int_a^{a+\lambda} f(x)(dx)^\alpha - {}_aI_b^\alpha f(x)g(x) \\ &= \frac{1}{\Gamma(\alpha + 1)} \int_a^{a+\lambda} [f(x) - f(a + \lambda)] [1 - g(x)] (dx)^\alpha \\ & \quad + \frac{1}{\Gamma(\alpha + 1)} \int_{a+\lambda}^b [f(a + \lambda) - f(x)] g(x)(dx)^\alpha \end{aligned} \quad (2.5)$$

and

$$\begin{aligned} & {}_aI_b^\alpha f(x)g(x) - \frac{1}{\Gamma(\alpha + 1)} \int_{b-\lambda}^b f(x)(dx)^\alpha \\ &= \frac{1}{\Gamma(\alpha + 1)} \int_a^{b-\lambda} [f(x) - f(b - \lambda)] g(x)(dx)^\alpha \\ & \quad + \frac{1}{\Gamma(\alpha + 1)} \int_{b-\lambda}^b [f(b - \lambda) - f(x)] [1 - g(x)] (dx)^\alpha. \end{aligned} \quad (2.6)$$

The interested reader is referred to several other related works including (for example) [2], [5], [6], [11], [12], [18] to [24] and [28] to [39] for the theory and applications of local fractional calculus.

In Section 3 of this paper, we give several inequalities which provide generalizations Steffensen's inequality (1.1) for local fractional integrals.

### 3. MAIN RESULTS

We start with the following important identity for our work. Throughout this paper,  $T_k(x)$  is defined by

$$T_k(x) = \frac{(n-1-k)^\alpha}{\Gamma(1+k\alpha)} \left( \frac{f^{(k\alpha)}(a)(x-a)^{k\alpha} - f^{(k\alpha)}(b)(x-b)^\alpha}{(b-a)^\alpha} \right). \quad (3.1)$$

**Lemma 3.1.** Let  $f^{(n-1)\alpha}(t)$  be absolutely continuous on  $[a, b]$  with  $f^{(n)\alpha} \in I_x^\alpha [a, b]$ . Then

$$\begin{aligned} & \frac{1}{n^\alpha} \left( \frac{f(x)}{\Gamma(1+\alpha)} + \sum_{k=1}^{n-1} F_k \right) - \frac{1}{(b-a)^\alpha \Gamma(1+\alpha)} \int_a^b f(y)(dy)^\alpha \\ &= \frac{1}{n^\alpha (b-a)^\alpha \Gamma(1+(n-1)\alpha) [\Gamma(1+\alpha)]^2} \\ & \quad \times \int_a^b (x-t)^{(n-1)\alpha} \kappa_\alpha(t, x) f^{(n)\alpha}(t) dt, \end{aligned}$$

where

$$F_k(x) = \frac{(n-k)^\alpha}{\Gamma(1+k\alpha)} \left( \frac{f^{(k-1)\alpha}(a)(x-a)^{k\alpha} - f^{(k-1)\alpha}(b)(x-b)^{k\alpha}}{(b-a)^\alpha} \right)$$

and

$$\kappa_\alpha(t, x) = \begin{cases} (t-a)^\alpha & (a \leq t \leq x \leq b) \\ (t-b)^\alpha & (a \leq x < t \leq b). \end{cases}$$

*Proof.* We begin by recalling the following local fractional Taylor's formula:

$$f(x) = f(y) + \sum_{k=1}^{n-1} \frac{f^{(k\alpha)}(y)(x-y)^{k\alpha}}{\Gamma(1+k\alpha)} + \frac{1}{\Gamma(1+\alpha)} \int_a^b \frac{f^{(n\alpha)}(t)(x-t)^{(n-1)\alpha}}{\Gamma(1+(n-1)\alpha)} (dt)^\alpha,$$

which, upon integration with respect to  $y$  from  $y = a$  to  $y = b$ , yields

$$\begin{aligned} \int_a^b (dy)^\alpha \int_y^x (dt)^\alpha &= \int_a^x (dy)^\alpha \int_y^x (dt)^\alpha + \int_x^b (dy)^\alpha \int_y^x (dt)^\alpha \\ &= \int_a^x (dt)^\alpha \int_a^t (dy)^\alpha - \int_x^b (dy)^\alpha \int_x^y (dt)^\alpha \\ &= \int_a^x (dt)^\alpha \int_a^t (dy)^\alpha - \int_x^b (dt)^\alpha \int_t^b (dy)^\alpha. \end{aligned}$$

Thus, by evaluating the last integral, we have

$$\begin{aligned} \frac{f(x)(b-a)^\alpha}{\Gamma(1+\alpha)} &= \frac{1}{\Gamma(1+\alpha)} \int_a^b f(y)(dy)^\alpha + \sum_{k=1}^{n-1} I_k + \frac{1}{\Gamma(1+(n-1)\alpha)[\Gamma(1+\alpha)]^2} \\ &\quad \times \int_a^b f^{(n\alpha)}(t)(x-t)^{(n-1)\alpha} \kappa_\alpha(t, x)(dt)^\alpha, \end{aligned} \quad (3.2)$$

where

$$I_k(x) = \frac{1}{\Gamma(1+\alpha)} \int_a^b \frac{f^{(k\alpha)}(y)(x-y)^{k\alpha}}{\Gamma(1+k\alpha)} (dy)^\alpha.$$

By local fractional integration by parts in the above expression for  $I_k(x)$ , we get

$$I_k(x) = I_{k-1}(x) - (b-a)^\alpha F_k(x)(n-k)^{-\alpha} \quad (1 \leq k \leq n-1).$$

Hence

$$(n-k)^\alpha [I_k(x) - I_{k-1}(x)] = -(b-a)^\alpha F_k(x). \quad (3.3)$$

The sum from  $k = 1$  to  $k = n-1$  in (3.3) is given by

$$\sum_{k=1}^{n-1} I_k = -(b-a)^\alpha \sum_{k=1}^{n-1} F_k(x) + (n-1)^\alpha I_0. \quad (3.4)$$

Substituting from (3.4) into (3.2) and rearranging the resulting equation, we have desired inequality asserted by Lemma 3.1.  $\square$

**Theorem 3.1.** *Let  $f : [a, b] \mapsto \mathbb{R}^\alpha$  be such that  $f^{(n-1)\alpha}$  is absolutely continuous for some  $n \in \mathbb{N} \setminus \{1\}$ . Suppose that the functions  $g, h \in I_x^\alpha[a, b]$  are such that  $h$  is positive and  $0^\alpha \leq g \leq 1^\alpha$  on  $[a, b]$ . Also let*

$${}_a I_{a+\lambda} h(t) = {}_a I_b g(t) h(t)$$

and the function  $S_1$  be defined by

$$S_1(x) = \begin{cases} \frac{1}{\Gamma(1+\alpha)} \int_a^x [1-g(t)] h(t)(dt)^\alpha & (x \in [a, a+\lambda]) \\ \frac{1}{\Gamma(1+\alpha)} \int_x^b g(t) h(t)(dt)^\alpha & (x \in [a+\lambda, b]). \end{cases} \quad (3.5)$$

Then

$$\begin{aligned}
& \frac{1}{\Gamma(1+\alpha)} \int_a^{a+\lambda} f(t)h(t)(dt)^\alpha - \frac{1}{\Gamma(1+\alpha)} \int_a^b f(t)g(t)h(t)(dt)^\alpha \\
& \quad - \Gamma(1+\alpha) \sum_{k=0}^{n-2} \frac{1}{\Gamma(1+\alpha)} \int_a^b S_1(x)T_k(x)(dx)^\alpha \\
& = -\frac{1}{(b-a)^\alpha \Gamma(1+(n-2)\alpha) [\Gamma(1+\alpha)]^2} \\
& \quad \times \int_a^b \left( \int_a^b S_1(x)(x-t)^{(n-2)\alpha} \kappa_\alpha(t,x)(dx)^\alpha \right) f^{(n\alpha)}(t)(dt)^\alpha. \tag{3.6}
\end{aligned}$$

*Proof.* It is easily seen that

$$\begin{aligned}
& \frac{1}{\Gamma(1+\alpha)} \int_a^{a+\lambda} f(t)h(t)(dt)^\alpha - \frac{1}{\Gamma(1+\alpha)} \int_a^b f(t)g(t)h(t)(dt)^\alpha \\
& = \frac{1}{\Gamma(1+\alpha)} \int_a^{a+\lambda} [f(t) - f(a+\lambda)] [1-g(t)] h(t)(dt)^\alpha \\
& \quad + \frac{1}{\Gamma(1+\alpha)} \int_{a+\lambda}^b [f(a+\lambda) - f(t)] g(t)h(t)(dt)^\alpha \\
& = \left( \frac{1}{\Gamma(1+\alpha)} \int_a^t [1-g(x)] h(x)(dx)^\alpha \right) [f(t) - f(a+\lambda)] \Big|_a^{a+\lambda} \\
& \quad - \frac{1}{\Gamma(1+\alpha)} \int_a^{a+\lambda} \left( \frac{1}{\Gamma(1+\alpha)} \int_a^t [1-g(x)] h(x)(dx)^\alpha \right) (df(t))^\alpha \\
& \quad + \left( \frac{1}{\Gamma(1+\alpha)} \int_b^t g(x)h(x)(dx)^\alpha \right) [f(a+\lambda) - f(t)] \Big|_{a+\lambda}^b \\
& \quad - \frac{1}{\Gamma(1+\alpha)} \int_{a+\lambda}^b \left( \frac{1}{\Gamma(1+\alpha)} \int_t^b g(x)h(x)(dx)^\alpha \right) (df(t))^\alpha \\
& = -\frac{1}{\Gamma(1+\alpha)} \int_a^{a+\lambda} \left( \frac{1}{\Gamma(1+\alpha)} \int_a^t [1-g(x)] h(x)(dx)^\alpha \right) (df(t))^\alpha \\
& \quad - \frac{1}{\Gamma(1+\alpha)} \int_{a+\lambda}^b \left( \frac{1}{\Gamma(1+\alpha)} \int_t^b g(x)h(x)(dx)^\alpha \right) (df(t))^\alpha \\
& = -\frac{1}{\Gamma(1+\alpha)} \int_a^b S_1(t) d(f(t))^\alpha \\
& = -\frac{1}{\Gamma(1+\alpha)} \int_a^b S_1(x) f^{(\alpha)}(x)(dx)^\alpha.
\end{aligned}$$

By applying Lemma 3.1 with  $f^{(\alpha)}$  and replacing  $n$  by  $n-1$  ( $n \in \mathbb{N} \setminus \{1\}$ ), we obtain

$$\begin{aligned}
f^{(\alpha)}(x) & = -\Gamma(1+\alpha) \sum_{k=0}^{n-2} T_k(x) + \frac{1}{(b-a)^\alpha \Gamma(1+(n-2)\alpha) \Gamma(1+\alpha)} \\
& \quad \times \int_a^b (x-t)^{(n-2)\alpha} \kappa_\alpha(t,x) f^{(n\alpha)}(t)(dt)^\alpha. \tag{3.7}
\end{aligned}$$

Moreover, by using the equation (3.7), we get

$$\begin{aligned}
& \frac{1}{\Gamma(1+\alpha)} \int_a^b S_1(x) f^{(\alpha)}(x) (dx)^\alpha \\
&= -\Gamma(1+\alpha) \sum_{k=0}^{n-2} \frac{1}{\Gamma(1+\alpha)} \int_a^b S_1(x) T_k(x) (dx)^\alpha \\
&\quad + \frac{1}{(b-a)^\alpha \Gamma(1+(n-2)\alpha) [\Gamma(1+\alpha)]^2} \\
&\quad \times \int_a^b S_1(x) \left( \int_a^b (x-t)^{(n-2)\alpha} \kappa_\alpha(t,x) f^{(n\alpha)}(t) (dt)^\alpha \right) (dx)^\alpha. \tag{3.8}
\end{aligned}$$

Finally, by applying Fubini's theorem for local fractional double integrals in the last term in (3.8), we arrive at the assertion (3.6) of Theorem 3.1.  $\square$

**Theorem 3.2.** *Let  $f : [a, b] \mapsto \mathbb{R}^\alpha$  be such that  $f^{(n-1)\alpha}$  is absolutely continuous for some  $n \in \mathbb{N} \setminus \{1\}$ . Suppose that the functions  $g, h \in I_x^\alpha[a, b]$  are such that  $h$  is positive and  $0^\alpha \leq g \leq 1^\alpha$  on  $[a, b]$ . Also let*

$${}_{b-\lambda}I_b h(t) = {}_aI_b g(t)h(t)$$

and the function  $S_2$  be defined by

$$S_2(x) = \begin{cases} \frac{1}{\Gamma(1+\alpha)} \int_a^x g(t)h(t) (dt)^\alpha & (x \in [a, b-\lambda]) \\ \frac{1}{\Gamma(1+\alpha)} \int_x^b [1-g(t)]h(t)(dt)^\alpha & (x \in [b-\lambda, b]). \end{cases} \tag{3.9}$$

Then

$$\begin{aligned}
& \frac{1}{\Gamma(1+\alpha)} \int_a^b f(t)g(t)h(t)(dt)^\alpha - \frac{1}{\Gamma(1+\alpha)} \int_{b-\lambda}^b f(t)h(t)(dt)^\alpha \\
&\quad - \Gamma(1+\alpha) \sum_{k=0}^{n-2} \frac{1}{\Gamma(1+\alpha)} \int_a^b S_2(x) T_k(x) (dx)^\alpha \\
&= -\frac{1}{(b-a)^\alpha \Gamma(1+(n-2)\alpha) [\Gamma(1+\alpha)]^2} \\
&\quad \times \int_a^b \left( \int_a^b S_2(x) (x-t)^{(n-2)\alpha} \kappa_\alpha(t,x) (dx)^\alpha \right) f^{(n\alpha)}(t) (dt)^\alpha. \tag{3.10}
\end{aligned}$$

*Proof.* We observe that

$$\begin{aligned}
& \frac{1}{\Gamma(1+\alpha)} \int_a^b f(t)g(t)h(t)(dt)^\alpha - \frac{1}{\Gamma(1+\alpha)} \int_{b-\lambda}^b f(t)h(t)(dt)^\alpha \\
&= \frac{1}{\Gamma(1+\alpha)} \int_a^{b-\lambda} [f(t) - f(b-\lambda)]g(t)h(t)(dt)^\alpha \\
&\quad + \frac{1}{\Gamma(1+\alpha)} \int_{b-\lambda}^b [f(b-\lambda) - f(t)][1-g(t)]h(t)(dt)^\alpha \\
&= \left( \frac{1}{\Gamma(1+\alpha)} \int_a^t g(x)h(x)(dx)^\alpha \right) [f(t) - f(b-\lambda)] \Big|_a^{b-\lambda} \\
&\quad - \frac{1}{\Gamma(1+\alpha)} \int_a^{b-\lambda} \left( \frac{1}{\Gamma(1+\alpha)} \int_a^t g(x)h(x)(dx)^\alpha \right) (df(t))^\alpha \\
&\quad + \left( \frac{1}{\Gamma(1+\alpha)} \int_b^t [1-g(x)]h(x)(dx)^\alpha \right) [f(b-\lambda) - f(t)] \Big|_{b-\lambda}^b \\
&\quad - \frac{1}{\Gamma(1+\alpha)} \int_{b-\lambda}^b \left( \frac{1}{\Gamma(1+\alpha)} \int_t^b [1-g(x)]h(x)(dx)^\alpha \right) (df(t))^\alpha \\
&= -\frac{1}{\Gamma(1+\alpha)} \int_a^{b-\lambda} \left( \frac{1}{\Gamma(1+\alpha)} \int_a^t g(x)h(x)(dx)^\alpha \right) (df(t))^\alpha \\
&\quad - \frac{1}{\Gamma(1+\alpha)} \int_{b-\lambda}^b \left( \frac{1}{\Gamma(1+\alpha)} \int_t^b [1-g(x)]h(x)(dx)^\alpha \right) (df(t))^\alpha \\
&= -\frac{1}{\Gamma(1+\alpha)} \int_a^b S_2(t)d(f(t))^\alpha \\
&= -\frac{1}{\Gamma(1+\alpha)} \int_a^b S_2(x)f^{(\alpha)}(x)(dx)^\alpha.
\end{aligned}$$

By making use of Lemma 3.1 with  $f^{(\alpha)}$  and replacing  $n$  by  $n-1$  ( $n \in \mathbb{N} \setminus \{1\}$ ), we obtain

$$\begin{aligned}
f^{(\alpha)}(x) &= -\Gamma(1+\alpha) \sum_{k=0}^{n-2} T_k(x) + \frac{1}{(b-a)^\alpha \Gamma(1+(n-2)\alpha) \Gamma(1+\alpha)} \\
&\quad \times \int_a^b (x-t)^{(n-2)\alpha} \kappa_\alpha(t,x) f^{(n\alpha)}(t)(dt)^\alpha.
\end{aligned} \tag{3.11}$$

Thus, by using this last equation (3.11), we get

$$\begin{aligned}
& \frac{1}{\Gamma(1+\alpha)} \int_a^b S_2(x)f^{(\alpha)}(x)(dx)^\alpha \\
&= -\Gamma(1+\alpha) \sum_{k=0}^{n-2} \frac{1}{\Gamma(1+\alpha)} \int_a^b S_2(x)T_k(x)(dx)^\alpha \\
&\quad + \frac{1}{(b-a)^\alpha \Gamma(1+(n-2)\alpha) [\Gamma(1+\alpha)]^2} \\
&\quad \times \int_a^b S_2(x) \left( \int_a^b (x-t)^{(n-2)\alpha} \kappa_\alpha(t,x) f^{(n\alpha)}(t)(dt)^\alpha \right) (dx)^\alpha.
\end{aligned} \tag{3.12}$$

Finally, by applying Fubini's theorem of local fractional double integrals in the last term in (3.12), we deduce the result asserted by Theorem 3.2.  $\square$

**Theorem 3.3.** Let  $f : [a, b] \mapsto \mathbb{R}^\alpha$  be such that  $f^{(n-1)\alpha}$  is absolutely continuous for some  $n \in \mathbb{N} \setminus \{1\}$ . Suppose that the functions  $g, h \in I_x^\alpha[a, b]$  are such that  $h$  is positive and  $0^\alpha \leq g \leq 1^\alpha$  on  $[a, b]$ . Also let

$${}_a I_{a+\lambda} h(t) = {}_a I_b g(t)h(t)$$

and the function  $S_1$  be defined by (3.5). If the function  $f$  is  $n$ -convex for local fractional calculus and

$$\int_a^b S_1(x)(x-t)^{(n-2)\alpha} \kappa_\alpha(t,x)(dx)^\alpha \leq 0 \quad (t \in [a, b]),$$

then the following inequality holds true:

$$\begin{aligned} \frac{1}{\Gamma(1+\alpha)} \int_a^b f(t)g(t)h(t)(dt)^\alpha &\leq \frac{1}{\Gamma(1+\alpha)} \int_a^{a+\lambda} f(t)h(t)(dt)^\alpha \\ &- \Gamma(1+\alpha) \sum_{k=0}^{n-2} \frac{1}{\Gamma(1+\alpha)} \int_a^b S_1(x)T_k(x)(dx)^\alpha. \end{aligned}$$

*Proof.* Since the function  $f$  is  $n$ -convex, we can suppose that  $f$  is  $n$  times differentiable and  $f^{(n\alpha)} \geq 0$ . Using this property and the assumption (3.6) of Theorem 3.1, we get the required inequality asserted by Theorem 3.3.  $\square$

**Theorem 3.4.** Let  $f : [a, b] \mapsto \mathbb{R}^\alpha$  be such that  $f^{(n-1)\alpha}$  is absolutely continuous for some  $n \in \mathbb{N} \setminus \{1\}$ . Suppose that the functions  $g, h \in I_x^\alpha[a, b]$  are such that  $h$  is positive and  $0^\alpha \leq g \leq 1^\alpha$  on  $[a, b]$ . Also let

$${}_{b-\lambda}I_b h(t) = {}_aI_b g(t)h(t)$$

and the function  $S_2$  be defined by (3.9). If the function  $f$  is  $n$ -convex for local fractional calculus and

$$\int_a^b S_2(x)(x-t)^{(n-2)\alpha} \kappa_\alpha(t,x)(dx)^\alpha \leq 0 \quad (t \in [a, b]),$$

then the following inequality holds true:

$$\begin{aligned} \frac{1}{\Gamma(1+\alpha)} \int_a^b f(t)g(t)h(t)(dt)^\alpha &\geq \frac{1}{\Gamma(1+\alpha)} \int_{b-\lambda}^b f(t)h(t)(dt)^\alpha \\ &+ \Gamma(1+\alpha) \sum_{k=0}^{n-2} \frac{1}{\Gamma(1+\alpha)} \int_a^b S_2(x)T_k(x)(dx)^\alpha. \end{aligned}$$

*Proof.* Since the function  $f$  is  $n$ -convex, we can suppose that  $f$  is  $n$  times differentiable and  $f^{(n\alpha)} \geq 0$ . Using this property and the assumption (3.10) of Theorem 3.2, we obtain the required inequality asserted by Theorem 3.4.  $\square$

#### 4. CONCLUDING REMARKS AND OBSERVATIONS

The present investigation is motivated essentially by widespread applications of fractional calculus and local fractional calculus (also called fractal calculus) in a large variety of problems in mathematical, physical and engineering sciences. Here, in this paper, we have first derived an identity for local fractional integrals. We then make use of this identity in order to derive several generalizations of the celebrated Steffensen's inequality associated with local fractional integrals. We have also briefly considered relevant connections of the results presented in this paper with the results which were proven in earlier works.

#### REFERENCES

- [1] A. M. Fink, Bounds of the deviation of a function from its averages, Czechoslovak Math. J. 42 (1992), 289–310.
- [2] H. Budak, M. Z. Sarikaya and H. Yildirim, New inequalities for local fractional integrals, Iranian J. Sci. Tech. Trans. A Sci. (in press).
- [3] S. Abramovich, M. K. Bakula, M. Matić and J.E. Pečarić, A variant of Jensen-Steffensen's inequality and quasi-arithmetic means, J. Math. Anal. Appl. 307 (2005), 370–386.
- [4] J. Bergh, A generalization of Steffensen's inequality, J. Math. Anal. Appl. 41 (1973), 187–191.
- [5] G.-S. Chen, Generalizations of Hölder's and some related integral inequalities on fractal space, J. Funct. Spaces Appl. 2013 (2013), Article ID 198405.
- [6] S. Erden and M. Z. Sarikaya, Generalized Pompeiu type inequalities for local fractional integrals and its applications, Appl. Math. Comput. 274 (2016), 282–291.
- [7] H. Gauchman, On a further generalization of Steffensen's inequality, J. Inequal. Appl. 5 (2000), 505–513.
- [8] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, Theory and Applications of Fractal Differential Equations, North-Holland Mathematical Studies, Vol. 204, Elsevier (North-Holland) Science Publishers, Amsterdam, 2006.

- [9] D. S. Mitrinović, *Analytic Inequalities*, Springer-Verlag, New-York, Heidelberg and Berlin, 1970.
- [10] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and New Inequalities in Analysis*, East European Series on Mathematical Analysis, Vol. 61, Kluwer Academic Publishers, Dordrecht, Boston and New York, 1993.
- [11] H. Mo, Generalized Hermite-Hadamard inequalities involving local fractional integral [arXiv:1410.1062].
- [12] H. Mo, X. Sui and D. Yu, Generalized convex functions on fractal sets and two related inequalities, *Abstr. Appl. Anal.* 2014 (2014), Article ID 636751.
- [13] J. E. Pečarić, On the Bellman generalization of Steffensen's inequality, *J. Math. Anal. Appl.* 88 (1982), 505–507.
- [14] J. E. Pečarić, On the Bellman generalization of Steffensen's inequality. II, *J. Math. Anal. Appl.* 104 (1984), 432–434.
- [15] J. E. Pečarić, A. Perušić and A. Vukelić, Generalizations of Steffensen's inequality via Fink identity and related results, *Adv. Inequal. Appl.* 2014 (2014), Article ID 9.
- [16] J. E. Pečarić, A. Perušić and A. Vukelić, Generalizations of Steffensen's inequality via Fink identity and related results. II, *Rend. Ist. Mat. Univ. Trieste* 47 (2015), 1–19.
- [17] F. Qi and B.-N. Guo, On Steffensen pairs, *J. Math. Anal. Appl.* 271 (2002), 534–541.
- [18] M. Z. Sarikaya and H. Budak, Generalized Ostrowski type inequalities for local fractional integrals, *Proc. Amer. Math. Soc.* (in press).
- [19] M. Z. Sarikaya and H. Budak, On generalized Hermite-Hadamard inequality for generalized convex function, *RGMIA Res. Rep. Collect.* 18 (2015), Article ID 64.
- [20] M. Z. Sarikaya, H. Budak and S. Erden, On new inequalities of Simpson's type for generalized convex functions, *RGMIA Res. Rep. Collect.* 18 (2015), Article ID 66.
- [21] M. Z. Sarikaya, S. Erden and H. Budak, Some integral inequalities for local fractional integrals, *RGMIA Res. Rep. Collect.* 18 (2015), Article ID 65.
- [22] M. Z. Sarikaya, S. Erden and H. Budak, Some generalized Ostrowski type inequalities involving local fractional integrals and applications, *Adv. Inequal. Appl.* 2016 (2016), Article ID 6.
- [23] M. Z. Sarikaya, T. Tunç and S. Erden, Generalized Steffensen inequalities for local fractional integrals, *RGMIA Res. Rep. Collect.* 18 (2015), Article ID 85.
- [24] M. Z. Sarikaya, T. Tunç and H. Budak, On generalized some integral inequalities for local fractional integrals, *Appl. Math. Comput.* 276 (2016), 316–323.
- [25] J. F. Steffensen, On certain inequalities and methods of approximation, *J. Inst. Actuar.* 51 (1919), 274–297.
- [26] U. M. Ozkan and H. Yildirim, Steffensen's integral inequality on time scales, *J. Inequal. Appl.* 2007 (2007), Article ID 46524.
- [27] S.-H. Wu and H. M. Srivastava, Some improvements and generalizations of Steffensen's integral inequality, *Appl. Math. Comput.* 192 (2007), 422–428.
- [28] X.-J. Yang, *Advanced Local Fractional Calculus and Its Applications*, World Science Publisher, New York, London and Hong Kong, 2012.
- [29] X.-J. Yang, Local fractional integral equations and their applications, *Adv. Comput. Sci. Appl.* 1 (2012), 234–239.
- [30] X.-J. Yang, Generalized local fractional Taylor's formula with local fractional derivative, *J. Expert Syst.* 1 (2012), 26–30.
- [31] X.-J. Yang, Local fractional Fourier analysis, *Adv. Mech. Engrg. Appl.* 1 (2012), 12–16.
- [32] X.-J. Yang, D. Baleanu and H. M. Srivastava, *Local Fractional Integral Transforms and Their Applications*, Academic Press (Elsevier Science Publishers), Amsterdam, Heidelberg, London and New York, 2016.
- [33] Y.-J. Yang, D. Baleanu and X.-J. Yang, Analysis of fractal wave equations by local fractional Fourier series method, *Adv. Math. Phys.* 2013 (2013), Article ID 632309.
- [34] X.-J. Yang, D. Baleanu and H. M. Srivastava, Local fractional similarity solution for the diffusion equation defined on Cantor sets, *Appl. Math. Lett.* 47 (2015), 54–60.
- [35] X.-J. Yang, J. A. T. Machado and H. M. Srivastava, A new numerical technique for solving the local fractional diffusion equation: Two-dimensional extended differential transform approach, *Appl. Math. Comput.* 274 (2016), 143–151.
- [36] X.-J. Yang and H. M. Srivastava, An asymptotic perturbation solution for a linear operator of free damped vibrations in fractal medium described by local fractional derivatives, *Commun. Nonlinear Sci. Numer. Simulat.* 29 (2015), 499–504.
- [37] X.-J. Yang, H. M. Srivastava and D. Baleanu, Initial-boundary value problems for local fractional Laplace equation arising in fractal electrostatics, *J. Appl. Nonlinear Dyn.* 4 (2015), 349–356.
- [38] X.-J. Yang, H. M. Srivastava and C. Cattani, Local fractional homotopy perturbation method for solving fractal partial differential equations arising in mathematical physics, *Romanian Rep. Phys.* 67 (2015), 752–761.
- [39] Y. Zhang, H. M. Srivastava and M.-C. Baleanu, Local fractional variational iteration algorithm II for non-homogeneous model associated with the non-differentiable heat flow, *Adv. Mech. Engrg.* 7 (10) (2015), 1–5.

<sup>1</sup>DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY

<sup>2</sup>DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF VICTORIA, VICTORIA, BRITISH COLUMBIA V8W 3R4, CANADA

<sup>3</sup>CENTER FOR GENERAL EDUCATION (DEPARTMENT OF SCIENCE AND TECHNOLOGY), CHINA MEDICAL UNIVERSITY, TAICHUNG 40402, TAIWAN, REPUBLIC OF CHINA

\*CORRESPONDING AUTHOR: tubatunc@duzce.edu.tr