
Faculty of Science

Faculty Publications

Some Higher-Degree Lacunary Fractional Splines in the Approximation of Fractional Differential Equations

Hari Mohan Srivastava, Pshtiwan Othman Mohammed, Juan L. G. Guirao, & Y. S. Hamed

March 2021

© 2021 H. M. Srivastava et al. This is an open access article distributed under the terms of the Creative Commons Attribution License <https://creativecommons.org/licenses/by/4.0/>

This article was originally published at:

<https://doi.org/10.3390/sym13030422>

Citation for this paper:

Srivastava, H. M., Mohammed, P. O., Guirao, J. L. G., & Hamed, Y. S. (2021). Some Higher-Degree Lacunary Fractional Splines in the Approximation of Fractional Differential Equations. *Symmetry*, 13(3), 1-13. <https://doi.org/10.3390/sym13030422>.

Article

Some Higher-Degree Lacunary Fractional Splines in the Approximation of Fractional Differential Equations

Hari Mohan Srivastava ^{1,2,3,4} , Pshtiwan Othman Mohammed ^{5,*} , Juan L. G. Guirao ^{6,*}  and Y. S. Hamed ⁷ 

¹ Department of Mathematics and Statistics, University of Victoria, Victoria, BC V8W 3R4, Canada; harimsri@math.uvic.ca

² Department of Medical Research, China Medical University Hospital, China Medical University, Taichung 40402, Taiwan

³ Department of Mathematics and Informatics, Azerbaijan University, 71 Jeyhun Hajibeyli Street, AZ1007 Baku, Azerbaijan

⁴ Section of Mathematics, International Telematic University Uninettuno, I-00186 Rome, Italy

⁵ Department of Mathematics, College of Education, University of Sulaimani, Sulaimani 46001, Kurdistan Region, Iraq

⁶ Departamento de Matemática Aplicada y Estadística, Universidad Politécnica de Cartagena, Campus de la Muralla, 30203 Cartagena, Murcia, Spain

⁷ Department of Mathematics and Statistics, College of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia; yasersalah@tu.edu.sa

* Correspondence: pshtiwansangawi@gmail.com (P.O.M.); juan.garcia@upct.es (J.L.G.G.)



Citation: Srivastava, H.M.; Mohammed, P.O.; Guirao, J.L.G.; Hamed, Y.S. Some Higher-Degree Lacunary Fractional Splines in the Approximation of Fractional Differential Equations. *Symmetry* **2021**, *13*, 422. <https://doi.org/10.3390/sym13030422>

Academic Editor: Palle E. T. Jorgensen

Received: 23 February 2021

Accepted: 4 March 2021

Published: 5 March 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: In this article, we begin by introducing two classes of lacunary fractional spline functions by using the Liouville–Caputo fractional Taylor expansion. We then introduce a new higher-order lacunary fractional spline method. We not only derive the existence and uniqueness of the method, but we also provide the error bounds for approximating the unique positive solution. As applications of our fundamental findings, we offer some Liouville–Caputo fractional differential equations (FDEs) to illustrate the practicability and effectiveness of the proposed method. Several recent developments on the theory and applications of FDEs in (for example) real-life situations are also indicated.

Keywords: FDEs; lacunary fractional spline; Riemann–Liouville fractional derivative; Liouville–Caputo fractional derivative; fractional Taylor's expansion; error bounds

1. Introduction

In recent years, the subject of fractional calculus (that is, the calculus of integrals and derivatives of any arbitrary real or complex order) has gained considerable popularity and importance due mainly to its demonstrated applications in the mathematical modelling of numerous seemingly diverse and widespread real-life problems in the fields of mathematical, physical, engineering and statistical sciences. Such operators of fractional-order derivatives as (for example) the Riemann–Liouville fractional derivative and the Liouville–Caputo fractional derivative are found to be potentially useful in the mathematical modelling of many of these problems (see, for example, [1–7]).

Motivated essentially by some recent developments (see, for example, [8]; see also [9,10] as well as the references to the related earlier works cited therein), we introduce and investigate here two classes of higher-order lacunary fractional spline functions based upon the Liouville–Caputo fractional derivative. For this purpose, we make use of a (presumably new) lacunary fractional spline method in order to investigate the above-mentioned classes of higher-order lacunary fractional spline functions by applying the Liouville–Caputo fractional Taylor expansion. We not only prove the existence and uniqueness of the method on each of the classes, but we also find the error bounds of the method via the modulus of continuity. Furthermore, with a view to illustrating our theoretical results, we successfully solve several Liouville–Caputo fractional differential equations (FDEs) by using the method

which we have introduced in this paper. Furthermore, finally, we graphically illustrate the numerical solutions which are presented here.

2. Definitions and Preliminaries

In this section, we revisit and recall all of the needed concepts and results involving fractional calculus and integral equations. We choose to divide this section into the following subsections.

2.1. Fractional Calculus

There are many definitions of fractional integrals and fractional derivatives, so it is always necessary to specify which definition is being used by us. In this article, we need the left Riemann–Liouville (L-RL) fractional integral and fractional derivative, as well as the left Liouville–Caputo (L-LC) derivative, which are defined as follows.

Definition 1 ([11–13]). For any L^1 function Y defined on $[b_1, b_2]$, the v th L-RL fractional integral of $Y(\eta)$ is defined for $\Re(v) > 0$ as follows:

$${}^{\text{RL}}\mathcal{I}_{b_1+}^v Y(\eta) := \frac{1}{\Gamma(v)} \int_{b_1}^{\eta} (\eta - \xi)^{v-1} Y(\xi) d\xi \quad (\eta \in [b_1, b_2]). \tag{1}$$

For any C^n function Y defined on $[b_1, b_2]$, the v th L-RL and L-LC fractional derivatives of $Y(\eta)$ are defined for $n - 1 \leq \Re(v) < n$ ($n \in \mathbb{N}$) as follows:

$${}^{\text{RL}}\mathcal{D}_{b_1+}^v Y(\eta) := \frac{d^n}{d\eta^n} {}^{\text{RL}}\mathcal{I}_{b_1+}^{n-v} Y(\eta) \quad (\eta \in [b_1, b_2]) \tag{2}$$

and

$${}^{\text{LC}}\mathcal{D}_{b_1+}^v Y(\eta) := {}^{\text{RL}}\mathcal{I}_{b_1+}^{n-v} \frac{d^n}{d\eta^n} Y(\eta) \quad (\eta \in [b_1, b_2]), \tag{3}$$

respectively, \mathbb{N} being the set of positive integers.

Lemma 1 (see [11–14]). Let $\mu > -1, \eta > b_1$ and $\Re(v) > 0$. Then the L-RL fractional integral of the power function satisfies the following result:

$${}^{\text{RL}}\mathcal{I}_{b_1+}^v (\eta - b_1)^\mu = \frac{\Gamma(\mu + 1)}{\Gamma(\mu + 1 + v)} (\eta - b_1)^{\mu+v}.$$

Moreover, the L-RL and L-LC fractional derivatives of a constant \mathfrak{C} are given by

$${}^{\text{RL}}\mathcal{D}_{b_1+}^v \mathfrak{C} = \mathfrak{C} \frac{\Gamma(\mu + 1)}{\Gamma(1 - v)} (\eta - b_1)^{-v} \quad (0 < v \leq 1)$$

and

$${}^{\text{LC}}\mathcal{D}_{b_1+}^v \mathfrak{C} = 0 \quad (0 < v \leq 1),$$

respectively.

Definition 2 (see [15,16]). Let $\Omega \subset \mathbb{R}$ denote an interval with $b_1 \in \Omega$ and $b_1 \leq \eta$ for all $\eta \in \Omega$. Furthermore, let $\delta \in \mathbb{R}^+$. We then define the sets ${}_{b_1}\mathbf{I}_\delta$ and ${}_{b_1}\mathbf{D}_\delta$ of functions as follows:

$${}_{b_1}\mathbf{I}_\delta = \left\{ Y : Y(\eta) \in C(\Omega) \text{ and } {}^{\text{RL}}\mathcal{I}_{b_1+}^\delta Y(\eta) \text{ exists and is finite in } \Omega \right\}$$

and

$${}_{b_1}D_\delta = \left\{ Y : Y(\eta) \in C(\Omega) \text{ and } {}^{LC}D_{b_1+}^\delta Y(\eta) \text{ exists and is finite in } \Omega \right\}.$$

Theorem 1 (see [15,16]). Let $\delta \in (0, 1]$ and $\ell \in \mathbb{N}$. Furthermore, let $Y(\eta)$ be a continuous function on $[b_1, b_2]$ satisfying the following conditions:

- (1) ${}^{LC}D_{b_1+}^{m\delta} Y \in C([b_1, b_2])$ and ${}^{LC}D_{b_1+}^{m\delta} Y \in {}_{b_1}I_\delta([b_1, b_2])$ for all $m = 1, 2, \dots, \ell$.
- (2) ${}^{LC}D_{b_1+}^{(\ell+1)\delta} Y(\eta)$ is continuous on $[b_1, b_2]$.

Then, for each $\eta \in [b_1, b_2]$,

$$Y(\eta) = \sum_{m=0}^{\ell} {}^{LC}D_{b_1+}^{m\delta} Y(a) \frac{(\eta - b_1)^{m\delta}}{\Gamma(m\delta + 1)} + R_\ell(\eta, b_1),$$

where

$$R_\ell(\eta, b_1) = \frac{(\eta - b_1)^{(\ell+1)\delta}}{\Gamma((\ell + 1)\delta + 1)} {}^{LC}D_{b_1+}^{(\ell+1)\delta} Y(\xi) \quad (a \leq \xi \leq \eta).$$

Definition 3 (see [17]). Let $Y : \mathcal{I} \subset \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function. Then, the modulus of a continuity of Y is defined by

$$\omega_\delta(h) := \omega(\delta; Y) = \sup_{|h| < \delta} |Y(\eta + h) - Y(\eta)|, \quad \forall \eta \in \mathcal{I}.$$

2.2. Formulation of the Problem

Given the mesh points: $\Delta : 0 = \eta_0 < \eta_1 < \dots < \eta_n = 1$, with $\eta_i = \frac{i}{n}$, $i = 0, 1, \dots, n - 1$, $h = \frac{1}{n}$, and the real numbers $\left\{ Y_i, {}^{LC}D_{0+}^{1/2} Y_i, \left({}^{LC}D_{0+}^{1/2} \right)^2 Y_i, \left({}^{LC}D_{0+}^{1/2} \right)^4 Y_i \right\}_{i=0}^n$ associated with the knots, our problem is to find s in a suitable class such that

$$\left({}^{LC}D_{0+}^{1/2} \right)^m s_\Delta(\eta_i) = \left({}^{LC}D_{0+}^{1/2} \right)^m Y_i \quad (i = 0, 1, \dots, n; m = 0, 1, 2, 4). \tag{4}$$

3. The First Class of Lacunary Fractional Spline

This section is being divided into the following subsections.

3.1. Existence and Uniqueness

Here, in this subsection, we define the class $S_{n,6}^{2\delta}$ of lacunary fractional spline functions as detailed below.

Definition 4. We say that s_Δ in $S_{n,6}^{2\delta}$ if the following conditions are fulfilled:

$${}^{LC}D_{0+}^{\delta m} s_\Delta \in C([0, 1]) \quad \left(m = 0, 1; \delta = \frac{1}{2} \right) \tag{5}$$

and

$$s_\Delta = \begin{cases} \sum_{i=0}^6 a_i (\eta - \eta_i)^{i/2} & (\eta \in [\eta_i, \eta_{i+1}] \forall i = 0, 1, \dots, n - 2) \\ \sum_{i=0}^7 a_i (\eta - \eta_i)^{i/2} & (\eta \in [\eta_{n-1}, \eta_n]). \end{cases} \tag{6}$$

We construct s_Δ in order that it is a solution of (4) for functions $(D)^{\delta m} S_\Delta \in C([0, 1])$ for $m = 0, 1, \dots, 6$ and $\delta = \frac{1}{2}$. For this purpose, we set

$$s_\Delta = \begin{cases} s_i(\eta) & (\eta_i \leq \eta \leq \eta_{i+1}; \forall i = 0, 1, \dots, n-2) \\ s_{n-1}(\eta) & (\eta_{n-1} \leq \eta \leq \eta_n). \end{cases}$$

Owing to (4) and (6), we can write

$$s_i(\eta) = \sum_{m=0}^2 \frac{1}{\Gamma(\frac{1}{2}m + 1)} (\eta - \eta_i)^{\frac{1}{2}m} (\text{LC}\mathcal{D}_{0+}^{1/2})^m Y_i + \frac{4}{3\sqrt{\pi}} (\eta - \eta_i)^{3/2} a_{i, \frac{3}{2}} + \frac{1}{2} (\eta - \eta_i)^2 (\text{LC}\mathcal{D}_{0+}^{1/2})^4 Y_i + \sum_{m=5}^6 \frac{1}{\Gamma(\frac{1}{2}m + 1)} (\eta - \eta_i)^{\frac{1}{2}m} a_{i, \frac{m}{2}}, \quad (7)$$

and

$$s_{n-1}(\eta) = \sum_{m=0}^2 \frac{1}{\Gamma(\frac{1}{2}m + 1)} (\eta - \eta_{n-1})^{\frac{1}{2}m} (\text{LC}\mathcal{D}_{0+}^{1/2})^m Y_{n-1} + \frac{4}{3\sqrt{\pi}} (\eta - \eta_{n-1})^{3/2} a_{n-1, \frac{3}{2}} + \frac{1}{2} (\eta - \eta_{n-1})^2 (\text{LC}\mathcal{D}_{0+}^{1/2})^4 Y_{n-1} + \sum_{m=5}^7 \frac{1}{\Gamma(\frac{1}{2}m + 1)} (\eta - \eta_{n-1})^{\frac{1}{2}m} a_{n-1, \frac{m}{2}}. \quad (8)$$

Then the coefficients are determined by

$$\begin{aligned} a_{i, \frac{3}{2}} &= \frac{\sqrt{\pi}(415\beta_i^1 + 128\beta_i^2 - 543\beta_i^3)}{4(60\pi + 287)}, \\ ha_{i, \frac{5}{2}} &= -\frac{\frac{225}{2}(\beta_i^3 - \beta_i^1)\pi + 240\beta_i^1 + 240\beta_i^2 - 128\beta_i^3}{\sqrt{\pi}(60\pi + 287)}, \\ h^{\frac{3}{2}}a_{i, 3} &= 6\beta_i^3 - 5\beta_i^1 + \frac{4(415\beta_i^1 + 128\beta_i^2 - 543\beta_i^3)}{60\pi + 287}, \\ a_{n-1, \frac{3}{2}} &= \frac{2776}{3485}\beta_{n-1}^1 + \frac{8}{697}\beta_{n-1}^2 - \frac{5612}{3485}\beta_{n-1}^3 + \frac{2796}{3485}\beta_{n-1}^4, \\ ha_{n-1, \frac{5}{2}} &= \frac{1}{697}(-2064\beta_{n-1}^1 - 90\beta_{n-1}^2 + 4960\beta_{n-1}^3 - 2806\beta_{n-1}^4), \\ h^{\frac{3}{2}}a_{n-1, 3} &= \frac{32}{697\sqrt{\pi}}(-3\beta_{n-1}^1 + 10\beta_{n-1}^2 - 9\beta_{n-1}^3 + 4858\beta_{n-1}^4), \\ h^2a_{n-1, \frac{7}{2}} &= \frac{1}{697}(3168\beta_{n-1}^1 - 105\beta_{n-1}^2 - 7224\beta_{n-1}^3 + 4858\beta_{n-1}^4), \end{aligned}$$

where

$$\begin{aligned} h^{\frac{3}{2}}\beta_i^1 &= 6\left(Y_{i+1} - Y_i - \frac{2}{\sqrt{\pi}} \text{LC}\mathcal{D}_{0+}^{1/2}Y_i - h(\text{LC}\mathcal{D}_{0+}^{1/2})^2Y_i - \frac{1}{2}h^2(\text{LC}\mathcal{D}_{0+}^{1/2})^4Y_i\right), \\ h\beta_i^2 &= \frac{15\sqrt{\pi}}{8}\left(\text{LC}\mathcal{D}_{0+}^{1/2}Y_{i+1} - \text{LC}\mathcal{D}_{0+}^{1/2}Y_i - h^{1/2}\frac{2}{\sqrt{\pi}}(\text{LC}\mathcal{D}_{0+}^{1/2})^2Y_i - \frac{4}{3\sqrt{\pi}}h^{3/2}(\text{LC}\mathcal{D}_{0+}^{1/2})^4Y_i\right), \\ h^{\frac{1}{2}}\beta_i^3 &= 2\left((\text{LC}\mathcal{D}_{0+}^{1/2})^2Y_{i+1} - (\text{LC}\mathcal{D}_{0+}^{1/2})^2Y_i - \frac{2}{\sqrt{\pi}}h^{1/2}(\text{LC}\mathcal{D}_{0+}^{1/2})^4Y_i\right), \end{aligned}$$

$$\begin{aligned}
 h^{\frac{3}{2}}\beta_{n-1}^1 &= \frac{105\sqrt{\pi}}{16} \left(Y_n - Y_{n-1} - h^{1/2} \frac{2}{\sqrt{\pi}} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^2 Y_{n-1} - \frac{1}{2} h^2 \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_{n-1} \right), \\
 h\beta_{n-1}^2 &= 6 \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} Y_n - {}^{\text{LC}}\mathcal{D}_{0+}^{1/2} Y_{n-1} - h^{1/2} \frac{2}{\sqrt{\pi}} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^2 Y_{n-1} \right. \\
 &\quad \left. - \frac{4}{3\sqrt{\pi}} h^{3/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_{n-1} \right), \\
 h^{\frac{1}{2}}\beta_{n-1}^3 &= \frac{15\sqrt{\pi}}{8} \left(\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^2 Y_n - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^2 Y_{n-1} - \frac{2}{\sqrt{\pi}} h^{1/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_{n-1} \right),
 \end{aligned}$$

and

$$\beta_{n-1}^4 = \frac{3\sqrt{\pi}}{4} h^{\frac{1}{2}} \left(\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_n - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_{n-1} \right).$$

Hence, clearly, we have proved Theorem 2 below.

Theorem 2. For a uniform partition Δ of the interval $I = [0, 1]$, there exists a unique fractional spline function $s_\Delta \in S_{n,6}^{2\delta}$ which is a solution of the problem involving (4).

3.2. Error Bounds

In this subsection, we first give the following main lemmas to work on our error bound theorem.

Lemma 2. Assume that Y satisfies the hypothesis of Theorem 1 for $m = j = 3, 5, 6$. For each $i = 0, 1, \dots, n - 2$, it is asserted that

$$\begin{aligned}
 \left| a_{i,\frac{j}{2}} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^j Y(\eta_i) \right| &\leq c_{i,\frac{j}{2}} h^{3-\frac{j}{2}} \omega_{6\delta}(h) \quad (j = 3, 5), \\
 \left| a_{i,3} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_i) \right| &\leq 6 \left(\frac{60\pi + 694}{60\pi + 287} \right) \omega_{6\delta}(h) \quad (\eta_i \leq \eta \leq \eta_{i+1}),
 \end{aligned}$$

where

$$c_{i,\frac{3}{2}} = \frac{543\sqrt{\pi}}{4(60\pi + 287)}$$

and

$$c_{i,\frac{5}{2}} = \frac{225\pi + 960}{2\sqrt{\pi}(60\pi + 287)}.$$

Proof. Suppose that $\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y \in C(I)$. Then, by using Theorem 1, we have

$$\begin{aligned}
 \beta_i^1 &= \frac{8}{\sqrt{\pi}} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_i + \frac{16h}{5\sqrt{\pi}} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_i + h^{3/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\xi_i), \\
 \beta_i^2 &= \frac{15\sqrt{\pi}}{8} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_i + \frac{15\sqrt{\pi}h}{16} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_i + h^{3/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_i),
 \end{aligned}$$

and

$$\beta_i^3 = \frac{4}{\sqrt{\pi}} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_i + \frac{8h}{3\sqrt{\pi}} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_i + h^{3/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_i),$$

where $\eta_i < \xi_i, \eta_i, \theta_i < \eta_{i+1}$ for $i = 0, 1, \dots, n-2$. We thus find that

$$\begin{aligned} \left| a_{i, \frac{3}{2}} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_i \right| &= \left| \frac{\sqrt{\pi}}{4(60\pi + 287)} \left[415\beta_i^1 + 128\beta_i^2 - 543\beta_i^3 \right] - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_i \right| \\ &= \frac{\sqrt{\pi}}{4(60\pi + 287)} h^{3/2} \left| 415 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\xi_i) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_i) \right] \right. \\ &\quad \left. + 128 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_i) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_i) \right] \right| \leq \frac{543\sqrt{\pi}}{4(60\pi + 287)} h^{3/2} \omega_{6\delta}(h), \end{aligned}$$

$$\begin{aligned} \left| a_{i, \frac{5}{2}} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_i \right| &= \frac{h^{3/2} h^{-1}}{\sqrt{\pi}(60\pi + 287)} \left| \frac{225}{2} \pi \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_i) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\xi_i) \right] \right. \\ &\quad \left. + 240 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\xi_i) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_i) \right] + 240 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_i) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_i) \right] \right| \\ &\leq \frac{225\pi + 960}{2\sqrt{\pi}(60\pi + 287)} h^{1/2} \omega_{6\delta}(h), \end{aligned}$$

and, finally, we get

$$\begin{aligned} \left| a_{i,3} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y_i \right| &= \left| 6\beta_i^3 - 5\beta_i^1 + \frac{4(415\beta_i^1 + 128\beta_i^2 - 543\beta_i^3)}{60\pi + 287} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y_i \right| \\ &\leq 5 \left| \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_i) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\xi_i) \right| + \frac{1660}{60\pi + 287} \left| \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\xi_i) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_i) \right| \\ &\quad + \frac{512}{60\pi + 287} \left| \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_i) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_i) \right| \leq 6 \left(\frac{60\pi + 694}{60\pi + 287} \right) \omega_{6\delta}(h), \end{aligned}$$

The proof of Lemma 2 is completed. \square

Lemma 3. Assume that Y satisfies the hypothesis of Theorem 1 for $m = 0, 1, \dots, 6$. The following estimates are valid:

$$\begin{aligned} \left| a_{n-1, \frac{j}{2}} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^j Y(\eta_{n-1}) \right| &\leq c_{n-1, \frac{j}{2}} h^{3-\frac{j}{2}} \omega_{6\delta}(h) \quad (j = 3, 5), \\ \left| a_{n-1,3} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta) \right| &\leq \left(\frac{1072}{697} \right) \omega_{6\delta}(h) \quad (\eta_{n-1} \leq \eta \leq \eta_n), \\ \left| a_{n-1, \frac{7}{2}} \right| &\leq \left(\frac{14217\sqrt{\pi}}{1394} \right) h^{-1/2} \omega_{6\delta}(h), \end{aligned}$$

where

$$c_{n-1, \frac{3}{2}} = \frac{63135\sqrt{\pi}}{41820}$$

and

$$c_{n-1, \frac{5}{2}} = \frac{4650\sqrt{\pi}}{697}.$$

Proof. Let $\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^m Y \in C(I)$ for $m = 0, 1, \dots, 6$. Then, by applying Theorem 1, we obtain

$$\begin{aligned} \beta_{n-1}^1 &= \frac{35}{4} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_{n-1} + \frac{7}{2} h \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_{n-1} + \frac{105\sqrt{\pi}}{96} h^{3/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\xi_{n-1}), \\ \beta_{n-1}^2 &= 6 \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_{n-1} + 3h \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_{n-1} + \frac{16}{5\sqrt{\pi}} h^{3/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_{n-1}), \end{aligned}$$

$$\beta_{n-1}^3 = \frac{15}{4} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_{n-1} + \frac{5}{2} h \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_{n-1} + \frac{15\sqrt{\pi}}{16} h^{3/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_{n-1}),$$

and

$$\beta_{n-1}^4 = \frac{3}{2} h \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_{n-1} + \frac{3\sqrt{\pi}}{4} h^{3/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\lambda_{n-1}),$$

where $\eta_{n-1} < \xi_{n-1}, \eta_{n-1}, \theta_{n-1}, \lambda_{n-1} < \eta_n$. We thus find that

$$\begin{aligned} \left| a_{n-1, \frac{3}{2}} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_{n-1} \right| &= \left| \frac{1}{3485} \left[2776\beta_{n-1}^1 + 40\beta_{n-1}^2 - 5612\beta_{n-1}^3 + 2796\beta_{n-1}^4 \right] \right. \\ &\quad \left. - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_{n-1} \right| \\ &\leq \frac{\sqrt{\pi}}{(3485)(12)} h^{3/2} \left| 36435 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\xi_{n-1}) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_{n-1}) \right] \right. \\ &\quad \left. + 1536 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_{n-1}) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_{n-1}) \right] \right. \\ &\quad \left. + 25164 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_{n-1}) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_{n-1}) \right] \right| \leq \frac{63135\sqrt{\pi}}{41820} h^{3/2} \omega_{6\delta}(h), \end{aligned}$$

$$\begin{aligned} \left| a_{n-1, \frac{5}{2}} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_{n-1} \right| &= \frac{h^{1/2}}{697} \frac{\sqrt{\pi}}{16} \left| -36120 \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\xi_{n-1}) \right. \\ &\quad \left. - 4608 \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_{n-1}) + 74400 \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_{n-1}) \right. \\ &\quad \left. - 33672 \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\lambda_{n-1}) \right| \leq \frac{4650\sqrt{\pi}}{697} h^{1/2} \omega_{6\delta}(h), \end{aligned}$$

$$\begin{aligned} \left| a_{n-1, 3} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y_{n-1} \right| &= \left| \frac{32}{697\sqrt{\pi}} \left[-3\beta_{n-1}^1 + 10\beta_{n-1}^2 - 9\beta_{n-1}^3 + 2\beta_{n-1}^4 \right] \right. \\ &\quad \left. - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y_{n-1} \right| \\ &\leq \frac{1}{697} \left| 105 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_{n-1}) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\xi_{n-1}) \right] \right. \\ &\quad \left. + 276 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_{n-1}) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\theta_{n-1}) \right] \right. \\ &\quad \left. + 673 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_{n-1}) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_{n-1}) \right] \right. \\ &\quad \left. + 24 \left[\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\lambda_{n-1}) - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y(\eta_{n-1}) \right] \right| \leq \frac{1072}{697} \omega_{6\delta}(h), \end{aligned}$$

and, finally, we get

$$\left| a_{n-1, \frac{7}{2}} \right| = \frac{h^{-2}}{697} \left| 3168\beta_{n-1}^1 - 105\beta_{n-1}^2 - 7224\beta_{n-1}^3 + 4858\beta_{n-1}^4 \right| \leq \frac{14217\sqrt{\pi}}{1349} h^{-1/2} \omega_{6\delta}(h).$$

This completes the proof of Lemma 3. \square

We now state and proof our result on the error bound.

Theorem 3 (Error bound). Let $s_\Delta \in S_{n,6}^{2\delta}$ be the solution of the problem involving (4). Then, for any

$$\left({}^{\text{LC}}\mathcal{D}_{0+}^\delta \right)^m Y \in C([0,1]) \quad \left(m = 0, 1, \dots; \delta = \frac{1}{2} \right),$$

the following error bound holds true:

$$\left| \left({}^{\text{LC}}\mathcal{D}_{0+}^\delta \right)^m s_i - \left({}^{\text{LC}}\mathcal{D}_{0+}^\delta \right)^m Y(\eta) \right| \leq c_{i,\delta m} h^{3-\delta m} \omega_{6\delta}(h) \quad (i = 0, 1, \dots, n - 1), \quad (9)$$

where $\omega_{6\delta}(h)$ is the modulus of continuity of $\left({}^{\text{LC}}\mathcal{D}_{0+}^\delta \right)^6 Y(\eta)$ and the constants $c_{i,\delta m}$ are given in Table 1.

Proof. Let us first assume that $\eta_i \leq \eta \leq \eta_{i+1}$ for $i = 0, 1, \dots, n - 2$. Then, from Equation (7) and by using Theorem 1, we have

$$s_i^{(m)}(\eta) - Y^{(m)}(\eta) = \begin{cases} \left(\frac{(\eta-\eta_i)^{3-\frac{m}{2}}}{\Gamma(4-\frac{m}{2})} \left[a_{i,\frac{3}{2}} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^3 Y_i \right] + \frac{(\eta-\eta_i)^{5-\frac{m}{2}}}{\Gamma(6-\frac{m}{2})} \left[a_{i,\frac{5}{2}} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_i \right] \right. \\ \left. + \frac{(\eta-\eta_i)^{6-\frac{m}{2}}}{\Gamma(7-\frac{m}{2})} \left[a_{i,3} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y_i \right] \right) & (m = 0, 1, \dots, 3); \\ \left(\frac{(\eta-\eta_i)^{5-\frac{m}{2}}}{\Gamma(6-\frac{m}{2})} \left[a_{i,\frac{5}{2}} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^5 Y_i \right] + \frac{(\eta-\eta_i)^{6-\frac{m}{2}}}{\Gamma(7-\frac{m}{2})} \left[a_{i,3} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y_i \right] \right) & (m = 4, 5); \\ a_{i,3} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^6 Y_i & (m = 6). \end{cases}$$

Now, by making use of the estimates asserted by Lemma 2, we can obtain the result for $i = 0, 1, \dots, n - 2$. In addition, by making use of Lemma 3, we can similarly obtain the rest of the results for $\eta_{n-1} \leq \eta \leq \eta_n$. This completes the proof of Theorem 3. \square

Table 1. The constants $c_{i,\delta m}$.

	$c_{i,0}$	$c_{i,\frac{1}{2}}$	$c_{i,1}$	$c_{i,\frac{3}{2}}$	$c_{i,2}$	$c_{i,\frac{5}{2}}$	$c_{i,3}$
$0 \leq i \leq n - 2$	$\frac{3909}{491}$	$\frac{707}{704}$	$\frac{1549}{145}$	$\frac{8776}{889}$	$\frac{1997}{163}$	$\frac{4378}{323}$	$\frac{3775}{339}$
$i = n - 1$	$\frac{716}{97}$	$\frac{2211}{122}$	$\frac{2308}{107}$	$\frac{8861}{934}$	$\frac{19,850}{697}$	$\frac{22,051}{697}$	$\frac{15,289}{697}$

4. The Second Class of Lacunary Fractional Spline

Here, in this connection, we consider the class of the lacunary fractional spline functions $S_{n,7}^{(2\delta,g)}$ and we denote this class of functions by $G_\Delta(\eta)$ such that

$$\left(D \right)^{\delta m} G_\Delta \in C([0,1]) \quad \left(m = 0, 1, 2; \delta = \frac{1}{2} \right), \quad (10)$$

$$G_\Delta = \sum_{i=0}^7 a_i (\eta - \eta_i)^{i/2} \quad (\eta \in [\eta_i, \eta_{i+1}] \forall i = 0, 1, 2, \dots, n - 1), \quad (11)$$

$$\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 G_\Delta(\eta_i + 0) = G_\Delta(\eta_i - 0) \quad (i = 0, 1, \dots, n - 1). \quad (12)$$

We construct G_Δ such that it is a solution of the problem involving (4) for

$$\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^m Y \in C([0,1]) \quad (m = 0, 1, \dots, 7).$$

For this purpose, we set

$$G_\Delta = G_i(\eta) \quad (i = 0, 1, \dots, n - 1).$$

Owing to (4) and (11), we can write

$$G_i(\eta) = \sum_{m=0}^2 \frac{1}{\Gamma(\frac{1}{2}m + 1)} (\eta - \eta_i)^{\frac{1}{2}m} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^m Y_i + \frac{4}{3\sqrt{\pi}} (\eta - \eta_i)^{3/2} b_{i,\frac{3}{2}} + \frac{1}{2} (\eta - \eta_i)^2 \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_i + \sum_{m=5}^7 \frac{1}{\Gamma(\frac{1}{2}m + 1)} (\eta - \eta_i)^{\frac{1}{2}m} b_{i,\frac{m}{2}}. \tag{13}$$

The constants can be determined, after using the conditions (10) to (12), as follows:

$$\begin{aligned} b_{i,\frac{3}{2}} &= \frac{1}{3485} (2776 \beta_i^1 + 40 \beta_i^2 - 5612 \beta_i^3 + 2796 \beta_i^4), \\ hb_{i,\frac{5}{2}} &= \frac{1}{697} (-2064 \beta_i^1 - 90 \beta_i^2 + 4960 \beta_i^3 - 2806 \beta_i^4), \\ h^{\frac{3}{2}} b_{i,3} &= \frac{32}{697\sqrt{\pi}} (-3 \beta_i^1 + 10 \beta_i^2 - 9 \beta_i^3 + 4858 \beta_i^4), \\ h^2 b_{i,\frac{7}{2}} &= \frac{1}{697} (3168 \beta_i^1 - 105 \beta_i^2 - 7224 \beta_i^3 + 4858 \beta_i^4), \end{aligned}$$

where

$$\begin{aligned} h^{\frac{3}{2}} \beta_i^1 &= \frac{105\sqrt{\pi}}{16} \left(Y_{i+1} - Y_i - h^{1/2} \frac{2}{\sqrt{\pi}} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^2 Y_i - \frac{1}{2} h^2 \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_i \right), \\ h\beta_i^2 &= 6 \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} Y_{i+1} - {}^{\text{LC}}\mathcal{D}_{0+}^{1/2} Y_i - h^{1/2} \frac{2}{\sqrt{\pi}} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^2 Y_i - \frac{4}{3\sqrt{\pi}} h^{3/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_i \right), \\ h^{\frac{1}{2}} \beta_i^3 &= \frac{15\sqrt{\pi}}{8} \left(\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^2 Y_{i+1} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^2 Y_i - \frac{2}{\sqrt{\pi}} h^{1/2} \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_i \right), \\ \beta_i^4 &= \frac{3\sqrt{\pi}}{4} h^{\frac{1}{2}} \left(\left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_{i+1} - \left({}^{\text{LC}}\mathcal{D}_{0+}^{1/2} \right)^4 Y_i \right). \end{aligned}$$

One can now observe that G_Δ is a *unique* element of $S_{n,7}^{(2\delta,g)}$ which is the solution of the interpolation problem (4). Then, by following the same technique which we used for the previous class, we are easily led the following theorem.

Theorem 4 (Error bound). *Let $s_\Delta \in S_{n,6}^{(2\delta,g)}$ be the solution of the interpolation problem involving (4). Then, for any*

$$\left({}^{\text{LC}}\mathcal{D}_{0+}^\delta \right)^m Y \in C([0,1]), \quad m = 0, 1, \dots, 7, \quad \delta = \frac{1}{2}, \quad \text{and } i = 0, 1, \dots, n - 1,$$

we have

$$\left| \left({}^{\text{LC}}\mathcal{D}_{0+}^\delta \right)^m s_i - \left({}^{\text{LC}}\mathcal{D}_{0+}^\delta \right)^m Y(\eta) \right| \leq c_{i,\delta m} h^{3.5-\delta m} \omega_{6\delta}(h), \tag{14}$$

where $\omega_{7\delta}(h)$ is the modulus of continuity of $\left({}^{\text{LC}}\mathcal{D}_{0+}^\delta \right)^7 Y(\eta)$, and the constants $c_{i,\delta m}$ are given in Table 2.

Table 2. The constants $c_{i,\delta m}$.

	$c_{i,0}$	$c_{i,\frac{1}{2}}$	$c_{i,1}$	$c_{i,\frac{3}{2}}$	$c_{i,2}$	$c_{i,\frac{5}{2}}$	$c_{i,3}$	$c_{i,\frac{7}{2}}$
$0 \leq i \leq n - 1$	$\frac{213}{49}$	$\frac{2894}{403}$	$\frac{5719}{530}$	$\frac{7565}{514}$	$\frac{5951}{350}$	$\frac{5277}{278}$	$\frac{808}{61}$	$\frac{8026}{697}$

5. Applications

In what follows, we provide some numerical examples to verify the validity of the proposed method.

Example 1. Consider the following FDE [8]:

$${}^{\text{LC}}\mathcal{D}_{0+}^2 Y(\eta) + 2 {}^{\text{LC}}\mathcal{D}_{0+}^\delta Y(\eta) + Y(\eta) = 2\eta + \frac{4}{\Gamma(4-\delta)}\eta^{3-\delta} + \frac{1}{3}\eta^3 \quad (0 < \delta \leq 1), \quad (15)$$

subject to the initial condition given by

$$Y(0) = Y'(0) = 0.$$

It is easily verified that the exact solution of this problem is

$$Y(\eta) = \frac{1}{3}\eta^3.$$

The maximal absolute errors and their fractional derivatives, which are obtained for $0 \leq \eta \leq 1$, are shown in Table 3. We note that

$$|e^{m\delta}(\eta)| = \left| \left({}^{\text{LC}}\mathcal{D}_{0+}^\delta\right)^m S_i(\eta) - \left({}^{\text{LC}}\mathcal{D}_{0+}^\delta\right)^m Y(\eta) \right|,$$

where

$$m = 0, 1, \dots, 4, \quad i = 0, 1, \dots, n-1, \quad \text{and} \quad \delta = \frac{1}{2}.$$

Figure 1a shows the exact and approximation solutions for $h = 0.05$. Also, the absolute error is shown in Figure 1b for $h = 0.05$.

Table 3. The observed maximum absolute errors for Example 1.

Methods	h	$ e(\eta) $	$ e^\delta(\eta) $	$ e^{2\delta}(\eta) $	$ e^{3\delta}(\eta) $	$ e^{4\delta}(\eta) $
Our method	0.1	4865×10^{-2}	9856×10^{-1}	1211×10^0	1145×10^0	2100×10^0
	0.01	1001×10^{-6}	9987×10^{-5}	6413×10^{-4}	1120×10^{-2}	8997×10^{-2}
	0.001	3544×10^{-9}	8770×10^{-8}	9948×10^{-6}	5189×10^{-5}	1102×10^{-3}
Method in [8]	0.1	4211×10^{-1}	1139×10^0	3832×10^0	1560×10^0	2546×10^0
	0.01	4211×10^{-4}	3603×10^{-3}	3832×10^{-2}	4936×10^{-2}	2546×10^{-1}
	0.001	1331×10^{-6}	1139×10^{-5}	3832×10^{-4}	1560×10^{-3}	2546×10^{-2}

Example 2. Consider the following FDE [8]:

$${}^{\text{LC}}\mathcal{D}_{0+}^\delta Y(\eta) = \eta^4 - \frac{1}{2}\eta^3 + \frac{24}{\Gamma(4-\delta)}\eta^{3-\delta} + \frac{3}{\Gamma(5-\delta)}\eta^{4-\delta} - Y(\eta) \quad (0 < \delta < 1), \quad (16)$$

With the initial condition $Y(0) = 0$. The exact solution is given by

$$Y(\eta) = \eta^4 - \frac{1}{2}\eta^3.$$

Together with

$$|e^{m\delta}(\eta)| = \left| \left({}^{\text{LC}}\mathcal{D}_{0+}^\delta\right)^m S_i(\eta) - \left({}^{\text{LC}}\mathcal{D}_{0+}^\delta\right)^m Y(\eta) \right|,$$

where

$$m = 0, 1, \dots, 4, \quad i = 0, 1, \dots, n-1 \quad \text{and} \quad \delta = \frac{1}{2},$$


we have derived the maximal absolute errors and their fractional derivatives. These results are shown in Table 4.

Figure 2a shows the exact and approximation solutions for $h = 0.05$. Also, the absolute error is shown in Figure 2b for $h = 0.05$.


All of the tables, which we have presented in this paper, show that our method is more accurate than the existing fractional spline methods.

Table 4. The observed maximum absolute errors for Example 2.

Methods	h	$ e(\eta) $	$ e^\delta(\eta) $	$ e^{2\delta}(\eta) $	$ e^{3\delta}(\eta) $	$ e^{4\delta}(\eta) $
Our method	0.1	9929×10^{-2}	5186×10^{-1}	2464×10^0	1063×10^1	3883×10^1
	0.01	9929×10^{-6}	1640×10^{-4}	2464×10^{-3}	3361×10^{-2}	3883×10^{-1}
	0.001	9929×10^{-10}	5186×10^{-8}	2464×10^{-6}	1063×10^{-4}	3883×10^{-3}
Method in [8]	0.1	7412×10^0	1487×10^0	3659×10^0	8398×10^1	2700×10^1
	0.01	7412×10^{-3}	4702×10^{-2}	3659×10^{-1}	2655×10^0	2700×10^0
	0.001	7412×10^{-6}	1487×10^{-4}	3659×10^{-3}	8398×10^{-2}	2700×10^{-1}




(a) Exact and approximation solutions for $h = 0.05$.




(b) Absolute errors for $h = 0.05$.

Figure 1. Plot illustrations for Example 1.



(a) Exact and approximation solutions for $h = 0.05$.



(b) Absolute errors for $h = 0.05$.

Figure 2. Plot illustrations for Example 2.

6. Conclusions

The content of this article can be summarized as follows:

1. Two classes of higher-order lacunary fractional spline functions are introduced.
2. A new lacunary fractional spline method is obtained for the above-mentioned classes by using the Liouville–Caputo fractional Taylor expansion.
3. The existence and uniqueness of the method on each of the classes is proved.

4. The error bounds of the method is shown via the modulus of continuity.
5. Some Liouville–Caputo FDEs are solved by using the new method in order to illustrate our theoretical results.
6. The numerical solutions are also illustrated graphically.

In conclusion, we remark that we have chosen to include some recent works (see, for example, [9,10,18–21] which will attract researchers and motivate them for further developments along these lines.

Author Contributions: Conceptualization, H.M.S., P.O.M. and Y.S.H.; methodology, H.M.S., P.O.M. and J.L.G.G.; software, P.O.M., J.L.G.G. and Y.S.H.; validation, H.M.S., P.O.M. and J.L.G.G.; formal analysis, P.O.M. and J.L.G.G.; investigation, P.O.M.; resources, H.M.S., P.O.M. and Y.S.H.; data curation, P.O.M. and J.L.G.G.; writing-original draft preparation, J.L.G.G.; writing-review and editing, H.M.S., J.L.G.G. and P.O.M.; visualization, H.M.S. and J.L.G.G.; supervision, H.M.S., P.O.M., J.L.G.G. and Y.S.H. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: This work was supported by the Taif University Researchers Supporting Project (No. TURSP-2020/155), Taif University, Taif, Saudi Arabia, and it is partially supported by Ministerio de Ciencia, Innovación y Universidades grant number PGC2018-097198-B-I00 and Fundación Séneca de la Región de Murcia grant number 20783/PI/18.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Mainardi, F. *Fractional Calculus and Waves in Linear Viscoelasticity: An Introduction to Mathematical Models*; Imperial College Press: Singapore, 2010.
2. Magin, R.L. *Fractional Calculus in Bioengineering*; Begell House Publishers Incorporated: Danbury, CT, USA, 2006.
3. Mohammed, P.O. A Generalized Uncertain Fractional Forward Difference Equations of Riemann–Liouville Type. *J. Math. Res.* **2019**, *11*, 43–50. [[CrossRef](#)]
4. Mohammed, P.O.; Abdeljawad, T. Discrete generalized fractional operators defined using h-discrete Mittag-Leffler kernels and applications to AB fractional difference systems. *Math. Meth. Appl. Sci.* **2020**, 1–26. [[CrossRef](#)]
5. Mohammed, P.O.; Abdeljawad, T.; Jarad, F.; Chu, Y.-M. Existence and uniqueness of uncertain fractional backward difference equations of Riemann–Liouville type. *Math. Prob. Eng.* **2020**, *2020*, 6598682. [[CrossRef](#)]
6. Alqudah, M.A.; Mohammed, P.O.; Abdeljawad, T. Solution of singular integral equations via Riemann–Liouville fractional integrals. *Math. Prob. Eng.* **2020**, *2020*, 1250970. [[CrossRef](#)]
7. Martinez, M.; Mohammed, P.O.; Valdes, J.E.N. Non-conformable fractional Laplace transform. *Kragujevac J. Math.* **2020**, *46*, 341–354.
8. Hamasalh, F.K.; Mohammed, P.O. Generalized quartic fractional spline interpolation with applications. *Internat. J. Open Probl. Compt. Math.* **2015**, *8*, 67–80. [[CrossRef](#)]
9. Tassaddiq1, A.; Yaseen, M.; Yousaf, A.; Srivastava, R. A cubic B-spline collocation method with new approximation for the numerical treatment of the heat equation with classical and non-classical boundary conditions. *Phys. Scr.* **2021**, *96*, 045212. [[CrossRef](#)]
10. Khabir, M.H.; Farah, R.A. 2017 Cubic B-spline collocation method for one-dimensional heat equation. *Pure Appl. Math.* **2017**, *6*, 51–58. [[CrossRef](#)]
11. Kilbas, A.A.; Srivastava, H.M.; Trujillo, J.J. *Theory and Applications of Fractional Differential Equations*; North-Holland Mathematics Studies; Elsevier Sci. B.V.: Amsterdam, The Netherlands, 2006; Volume 204.
12. Hilfer, R. *Applications of Fractional Calculus in Physics*; World Scientific Publishing Company: Singapore, 2000.
13. Podlubny, I. *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications*; Mathematics in Science and Engineering; Academic Press: New York, NY, USA, 1999; Volume 198.
14. Srivastava, H.M. Fractional-order derivatives and integrals: Introductory overview and recent developments. *Kyungpook Math. J.* **2020**, *60*, 73–116.
15. Usero, D. Fractional Taylor Series for Caputo Fractional Derivatives: Construction of Numerical Schemes. Preprint 2008. Available online: http://www.fdi.ucm.es/profesor/lvazquez/calfrac/docs/paper_Usero.pdf (accessed on 2 March 2018).

16. Fernandez, A.; Baleanu, D. The mean value theorem and Taylor's theorem for fractional derivatives with Mittag-Leffler kernel. *Adv. Differ. Equ.* **2018**, *2018*, 86. [[CrossRef](#)] [[PubMed](#)]
17. Steffens, K.G. *The History of Approximation Theory*; Birkhäuser: Basel, Switzerland, 2006.
18. Srivastava, H.M. Diabetes and its resulting complications: Mathematical modeling via fractional calculus. *Public Health Open Access* **2020**, *4*, 1–5.
19. Srivastava, H.M.; Deniz, D.; Saad, K.M. An efficient semi-analytical method for solving the generalized regularized long wave equations with a new fractional derivative operator. *J. King Saud Univ. Sci.* **2021**, *33*, 101345. [[CrossRef](#)]
20. Singh, H.; Srivastava, H.M.; Hammouch, Z.; Nisar, K.S. Numerical simulation and stability analysis for the fractional-order dynamics of COVID-19. *Results Phys.* **2021**, *20*, 103722. [[CrossRef](#)] [[PubMed](#)]
21. Srivastava, H.M.; Mohammed, P.O. A correlation between solutions of uncertain fractional forward difference equations and their paths. *Front. Phys.* **2020**, *8*, 280. [[CrossRef](#)]